

KOS Diagnostic Lab (A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. VEENA SETH

AGE/ GENDER : 78 YRS/FEMALE **PATIENT ID** : 1534349

COLLECTED BY : SURJESH REG. NO./LAB NO. :012407010026

REFERRED BY **REGISTRATION DATE** : 01/Jul/2024 09:35 AM BARCODE NO. :01512291 **COLLECTION DATE** : 01/Jul/2024 09:47AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 01/Jul/2024 11:42AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Value Unit **Biological Reference interval** Test Name

CLINICAL PATHOLOGY

MICROALBUMIN/CREATININE RATIO - RANDOM URINE

MICROALBUMIN: RANDOM URINE mg/L 0 - 25 94.15^H by SPECTROPHOTOMETRY CREATININE: RANDOM URINE 41.71 mg/dl 20 - 320

by SPECTROPHOTOMETRY MICROALBUMIN/CREATININE RATIO -0 - 30225.73H mg/g

RANDOM URINE

by SPECTROPHOTOMETRY

INTERPRETATION:-

| PHYSIOLOGICALLY NORMAL: | mg/L | 0 - 30 |
|-------------------------|------|----------|
| MICROALBUMINURIA: | mg/L | 30 - 300 |
| GROSS PROTEINURIA: | mg/L | > 300 |

Long standing un-treated Diabetes and Hypertension can lead to renal dysfunction.

2. Diabetic nephropathy or kidney disease is the most common cause of end stage renal disease(ERSD) or kidney failure.

3. Presence of Microalbuminuria is an early indicator of onset of compromised renal function in these patients.

4. Microalbuminuria is the condition when urinary albumin excretion is between 30-300 mg & above this it is called as macroalbuminuria, the presence of which indicates serious kidney disease.

5.Microalbuminuria is not only associated with kidney disease but of cardiovascular disease in patients with dibetes & hypertension.

6. Microalbuminuria reflects vascular damage & appear to be a marker of of early arterial disease & endothelial dysfunction.

NOTE:- IF A PATIENT HAS = 1+ PROTEINURIA (30 mg/dl OR 300 mg/L) BY URINE DIPSTICK (URINEANALYSIS), OVERT PROTEINURIA IS PRESENT AND TESTING FOR MICROALBUMIN IS INAPPROPIATE. IN SUCH A CASE, URINE PROTEIN: CREATININE RATIO OR 24 HOURS TOTAL URINE MICROPROTEIN IS **APPROPIATE**

*** End Of Report ***



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

