

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mrs. PANA SHARMA
AGE/ GENDER : 53 YRS/FEMALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 01512417
CLIENT CODE. : KOS DIAGNOSTIC LAB
CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

PATIENT ID : 1536906
REG. NO./LAB NO. : 012407030010
REGISTRATION DATE : 03/Jul/2024 08:20 AM
COLLECTION DATE : 03/Jul/2024 08:24 AM
REPORTING DATE : 03/Jul/2024 10:07 AM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

IMMUNOPATHOLOGY/SEROLOGY

ANTI THYROID PEROXIDASE (TPO/AMA) ANTIBODIES

ANTI TPO/AMA ANTIBODIES: SERUM by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)	1.3	IU/mL	0.00 - 10.0 DIABETES (II): < 25.0
---------------------------------------------------------------------------	-----	-------	--------------------------------------

INTERPRETATION:

- Thyroperoxidase (TPO) is an enzyme involved in thyroid hormone synthesis, catalyzing the oxidation of iodide on tyrosine residues in thyroglobulin for the synthesis of triiodothyronine and thyroxine (tetraiodothyronine).
- TPO is a membrane-associated hemo glycoprotein expressed only in thyrocytes and is one of the most important thyroid gland antigens.
- Anti-TPO is technically superior and a more specific method for measuring thyroid auto-antibodies, It is especially useful in patients presenting with subclinical hypothyroidism where TSH is elevated but Free T4 levels are normal.

INCREASED LEVELS (Autoimmune thyroid disease):

- Hashimoto thyroiditis.
- Idiopathic myxedema.
- Graves disease
- Post-partum thyroiditis.
- Primary hypothyroidism due to Hashimoto thyroiditis.

NOTE:

- The highest TPO antibody levels are observed in patients suffering from Hashimoto thyroiditis. In this disease, the prevalence of TPO antibodies is about 90% of cases, confirming the autoimmune origin of the disease.
- These auto-antibodies also frequently occur (60%-80%) in the course of Graves disease.
- In patients with subclinical hypothyroidism, the presence of TPO antibodies is associated with an increased risk of developing overt hypothyroidism.

*** End Of Report ***



DR. VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR. YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)

