

**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. ALKA GUPTA	<b>PATIENT ID</b>	: 1131930
<b>AGE/ GENDER</b>	: 37 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: <b>012407120058</b>
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 12/Jul/2024 07:09 PM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 12/Jul/2024 07:11 PM
<b>BARCODE NO.</b>	: 01513006	<b>REPORTING DATE</b>	: 12/Jul/2024 09:19 PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

**VITAMINS**

**VITAMIN D/25 HYDROXY VITAMIN D3**

<b>VITAMIN D (25-HYDROXY VITAMIN D3): SERUM</b> by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)	16.3 <sup>L</sup>	ng/mL	<b>DEFICIENCY:</b> < 20.0 <b>INSUFFICIENCY:</b> 20.0 - 30.0 <b>SUFFICIENCY:</b> 30.0 - 100.0 <b>TOXICITY:</b> > 100.0
--	-------------------	-------	--

**INTERPRETATION:**

<b>DEFICIENT:</b>	< 20	ng/mL
<b>INSUFFICIENT:</b>	21 - 29	ng/mL
<b>PREFERRED RANGE:</b>	30 - 100	ng/mL
<b>INTOXICATION:</b>	> 100	ng/mL

- Vitamin D compounds are derived from dietary ergocalciferol (from plants, Vitamin D2), or cholecalciferol (from animals, Vitamin D3), or by conversion of 7- dihydrocholecalciferol to Vitamin D3 in the skin upon Ultraviolet exposure.
- 25-OH--Vitamin D represents the main body resevoir and transport form of Vitamin D and transport form of Vitamin D, being stored in adipose tissue and tightly bound by a transport protein while in circulation.
- Vitamin D plays a primary role in the maintenance of calcium homeostatis. It promotes calcium absorption, renal calcium absorption and phosphate reabsorption, skeletal calcium deposition, calcium mobilization, mainly regulated by parathyroid hormone (PTH).
- Severe deficiency may lead to failure to mineralize newly formed osteoid in bone, resulting in rickets in children and osteomalacia in adults.

**DECREASED:**

- Lack of sunshine exposure.
- Inadequate intake, malabsorption (celiac disease)
- Depressed Hepatic Vitamin D 25- hydroxylase activity
- Secondary to advanced Liver disease
- Osteoporosis and Secondary Hyperparathroidism (Mild to Moderate deficiency)
- Enzyme Inducing drugs: anti-epileptic drugs like phenytoin, phenobarbital and carbamazepine, that increases Vitamin D metabolism.

**INCREASED:**

- Hypervitaminosis D is Rare, and is seen only after prolonged exposure to extremely high doses of Vitamin D. When it occurs, it can result in severe hypercalcemia and hyperphosphatemia.

**CAUTION:** Replacement therapy in deficient individuals must be monitored by periodic assessment of Vitamin D levels in order to prevent hypervitaminosis D

**NOTE:-**Dark coloured individuals as compare to whites, is at higher risk of developing Vitamin D deficiency due to excess of melanin pigment which interfere with Vitamin D absorption.

\*\*\* End Of Report \*\*\*



*Chopra*

**DR.VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

*Chopra*

**DR.YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.