

Dr. Vinay Chopra  
MD (Pathology & Microbiology)  
Chairman & Consultant Pathologist

Dr. Yugam Chopra  
MD (Pathology)  
CEO & Consultant Pathologist

<b>NAME</b>	: Mr. KULWANT SINGH	<b>PATIENT ID</b>	: 1548660
<b>AGE/ GENDER</b>	: 57 YRS/MALE	<b>REG. NO./LAB NO.</b>	: <b>012407140004</b>
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 14/Jul/2024 08:03 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 14/Jul/2024 08:09 AM
<b>BARCODE NO.</b>	: 01513097	<b>REPORTING DATE</b>	: 14/Jul/2024 09:58 AM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

**CLINICAL CHEMISTRY/BIOCHEMISTRY**

**LIPID PROFILE : BASIC**

CHOLESTEROL TOTAL: SERUM <i>by CHOLESTEROL OXIDASE PAP</i>	155.22	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SERUM <i>by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)</i>	94.92	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (DIRECT): SERUM <i>by SELECTIVE INHIBITION</i>	56.3	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	97.94	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTEROL: SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	98.92	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL: SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	18.98	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	423.36	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL RATIO: SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	2.76	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0
LDL/HDL RATIO: SERUM	1.74	RATIO	LOW RISK: 0.50 - 3.0



DR. VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR. YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

Dr. Vinay Chopra  
MD (Pathology & Microbiology)  
Chairman & Consultant Pathologist

Dr. Yugam Chopra  
MD (Pathology)  
CEO & Consultant Pathologist

<b>NAME</b>	: Mr. KULWANT SINGH	<b>PATIENT ID</b>	: 1548660
<b>AGE/ GENDER</b>	: 57 YRS/MALE	<b>REG. NO./LAB NO.</b>	: <b>012407140004</b>
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 14/Jul/2024 08:03 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 14/Jul/2024 08:09AM
<b>BARCODE NO.</b>	: 01513097	<b>REPORTING DATE</b>	: 14/Jul/2024 09:58AM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
<i>by CALCULATED, SPECTROPHOTOMETRY</i>			
<b>TRIGLYCERIDES/HDL RATIO: SERUM</b>	1.69 <sup>L</sup>	RATIO	MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0 <b>3.00 - 5.00</b>
<i>by CALCULATED, SPECTROPHOTOMETRY</i>			

**INTERPRETATION:**

1. Measurements in the same patient can show physiological & analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.
2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.
3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.
4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogenic lipoproteins such as LDL, VLDL, IDL, Lp(a), Chylomicron remnants) along with LDL-cholesterol as co-primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL & Non HDL.
5. Additional testing for Apolipoprotein B, hsCRP, Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR. VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR. YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra  
MD (Pathology & Microbiology)  
Chairman & Consultant Pathologist

Dr. Yugam Chopra  
MD (Pathology)  
CEO & Consultant Pathologist

<b>NAME</b>	: Mr. KULWANT SINGH	<b>PATIENT ID</b>	: 1548660
<b>AGE/ GENDER</b>	: 57 YRS/MALE	<b>REG. NO./LAB NO.</b>	: <b>012407140004</b>
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 14/Jul/2024 08:03 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 14/Jul/2024 08:09AM
<b>BARCODE NO.</b>	: 01513097	<b>REPORTING DATE</b>	: 14/Jul/2024 09:58AM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

**CALCIUM**

CALCIUM: SERUM	9.44	mg/dL	8.50 - 10.60
----------------	------	-------	--------------

*by ARSENAZO III, SPECTROPHOTOMETRY*

**INTERPRETATION:-**

1. Serum calcium (total) estimation is used for the diagnosis and monitoring of a wide range of disorders including diseases of bone, kidney, parathyroid gland, or gastrointestinal tract.
2. Calcium levels may also reflect abnormal vitamin D or protein levels.
3. The calcium content of an adult is somewhat over 1 kg (about 2% of the body weight). Of this, 99% is present as calcium hydroxyapatite in bones and <1% is present in the extra-osseous intracellular space or extracellular space (ECS).
4. In serum, calcium is bound to a considerable extent to proteins (approximately 40%), 10% is in the form of inorganic complexes, and 50% is present as free or ionized calcium.

**NOTE:-**Calcium ions affect the contractility of the heart and the skeletal musculature, and are essential for the function of the nervous system. In addition, calcium ions play an important role in blood clotting and bone mineralization.

**HYPOCALCEMIA (LOW CALCIUM LEVELS) CAUSES :-**

1. Due to the absence or impaired function of the parathyroid glands or impaired vitamin-D synthesis.
2. Chronic renal failure is also frequently associated with hypocalcemia due to decreased vitamin-D synthesis as well as hyperphosphatemia and skeletal resistance to the action of parathyroid hormone (PTH).
3. **NOTE:-** A characteristic symptom of hypocalcemia is latent or manifest tetany and osteomalacia.

**HYPERCALCEMIA (INCREASE CALCIUM LEVELS) CAUSES:-**

1. Increased mobilization of calcium from the skeletal system or increased intestinal absorption.
2. Primary hyperparathyroidism (pHPT)
3. Bone metastasis of carcinoma of the breast, prostate, thyroid gland, or lung.

**NOTE:-**Severe hypercalcemia may result in cardiac arrhythmia.

\*\*\* End Of Report \*\*\*



DR. VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR. YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.