

Dr. Vinay Chopra  
MD (Pathology & Microbiology)  
Chairman & Consultant Pathologist

Dr. Yugam Chopra  
MD (Pathology)  
CEO & Consultant Pathologist

NAME : Mrs. ANU  
AGE/ GENDER : 26 YRS/FEMALE  
COLLECTED BY :  
REFERRED BY :  
BARCODE NO. : 01513180  
CLIENT CODE. : KOS DIAGNOSTIC LAB  
CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

PATIENT ID : 1549150  
REG. NO./LAB NO. : 012407150034  
REGISTRATION DATE : 15/Jul/2024 10:05 AM  
COLLECTION DATE : 15/Jul/2024 10:07 AM  
REPORTING DATE : 15/Jul/2024 11:32 AM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## IMMUNOPATHOLOGY/SEROLOGY

### ANTI THYROID PEROXIDASE (TPO/AMA) ANTIBODIES

ANTI TPO/AMA ANTIBODIES: SERUM	< 1.00	IU/mL	0.00 - 10.0
by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)			DIABETES (II): < 25.0

#### INTERPRETATION:

1. Thyroperoxidase (TPO) is an enzyme involved in thyroid hormone synthesis, catalyzing the oxidation of iodide on tyrosine residues in thyroglobulin for the synthesis of triiodothyronine and thyroxine (tetraiodothyronine).
2. TPO is a membrane-associated hemo glycoprotein expressed only in thyrocytes and is one of the most important thyroid gland antigens.
3. Anti-TPO is technically superior and a more specific method for measuring thyroid auto-antibodies, It is especially useful in patients presenting with subclinical hypothyroidism where TSH is elevated but Free T4 levels are normal.

#### INCREASED LEVELS (Autoimmune thyroid disease):

1. Hashimoto thyroiditis.
2. Idiopathic myxedema.
3. Graves disease
4. Post-partum thyroiditis.
5. Primary hypothyroidism due to Hashimoto thyroiditis.

#### NOTE:

1. The highest TPO antibody levels are observed in patients suffering from Hashimoto thyroiditis. In this disease, the prevalence of TPO antibodies is about 90% of cases, confirming the autoimmune origin of the disease.
2. These auto-antibodies also frequently occur (60%-80%) in the course of Graves disease.
3. In patients with subclinical hypothyroidism, the presence of TPO antibodies is associated with an increased risk of developing overt hypothyroidism.

\*\*\* End Of Report \*\*\*



DR. VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR. YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)

