



Test Name	Value	Unit	Biological Reference interval
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANT		
CLIENT ADDRESS	. C240/1 NICHOLCON DOAD AMDALA CANT	νŢ	
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPORTING DATE	: 15/Jul/2024 11:26AM
BARCODE NO.	: 01513194	COLLECTION DATE	: 15/Jul/2024 11:22AM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	REGISTRATION DATE	: 15/Jul/2024 11:20 AM
COLLECTED BY	:	REG. NO./LAB NO.	: 012407150048
AGE/ GENDER	: 33 YRS/FEMALE	PATIENT ID	: 1549190
NAME	: Mrs. MAMTA		
	Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Patholog		(Pathology)
\$0 9001 : 2008 CERT	IFIED LAB	EXCELLENCE IN HEALTHCARE	& DIAGNOSTICS

HAEMATOLOGY

BLOOD GROUP (ABO) AND RH FACTOR TYPING

ABO GROUP by SLIDE AGGLUTINATION RH FACTOR TYPE by SLIDE AGGLUTINATION

А

POSITIVE



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com



Page 1 of 5





	Dr. Vinay Ch MD (Pathology & Chairman & Cons	Microbiology)		(Pathology)
IAME	: Mrs. MAMTA			
AGE/ GENDER	: 33 YRS/FEMALE		PATIENT ID	: 1549190
COLLECTED BY	:		REG. NO./LAB NO.	: 012407150048
REFERRED BY	: LOOMBA HOSPITAL (AMBAL	A CANTT)	REGISTRATION DATE	: 15/Jul/2024 11:20 AM
BARCODE NO.	: 01513194		COLLECTION DATE	: 15/Jul/2024 11:22AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		REPORTING DATE	: 15/Jul/2024 01:11PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, A	AMBALA CANT	Т	
Test Name		Value	Unit	Biological Reference interval
ΗΕΡΔΤΙΤΙς Ο ΔΝΤΙΒΟ			(HCV) ANTIBODY: TOT	
	DY (HCV) TOTAL: SERUM	0.08	S/CO	NEGATIVE: < 1.00
HEPATITIS C ANTIBO		NON - R	EACTIVE	POSITIVE: > 1.00
RESULT		NON		
	ESCENT MICROPARTICLE IMMUNOAS	SSAY)		
INTERPRETATION:-	SULT (INDEX)		REMARKS	
KI NI	< 1.00		NON - REACTIVE/NOT - DE	TECTED
	> =1.00		ASYMPTOMATIC/INFECTIVE S	
needle punctures in I compared to HAV & I	nealthcare workers, dialysis patie	ents and rarely occurs in 85 %	from mother to infant. 10 % of infected individuals. In hid	ntation, injection drug abusers, accidental of new cases show sexual transmission. As gh risk population, the predictive value of Anti
1. Indicator of past or 2. Routine screening NOTE:	present infection, but does not c of low and high prevelance popu	ilation including	g blood donors.	ved Infection.

1. False positive results are seen in Auto-immune disease, Rheumatoid Factor, HYpergammaglobulinemia, Paraproteinemia, Passive antibody transfer, Anti-idiotypes and Anti-superoxide dismutase.

2. False negative results are seen in early Acute infection, Immunosuppression and Immuno-incompetence.

3. HCV-RNĂ PCR recommended in all reactive results to differentiate between past and present infection.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







	Dr. Vinay Chopr MD (Pathology & Mic Chairman & Consulta	robiology)		(Pathology)
NAME	: Mrs. MAMTA			
AGE/ GENDER	: 33 YRS/FEMALE		PATIENT ID	: 1549190
COLLECTED BY	:		REG. NO./LAB NO.	: 012407150048
REFERRED BY	: LOOMBA HOSPITAL (AMBALA C	ANTT)	REGISTRATION DATE	: 15/Jul/2024 11:20 AM
BARCODE NO.	: 01513194		COLLECTION DATE	: 15/Jul/2024 11:22AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		REPORTING DATE	: 15/Jul/2024 01:38PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AME	BALA CANTT		
CLIENT ADDRESS Test Name	: 6349/1, NICHOLSON ROAD, AME	BALA CANTT	Unit	Biological Reference interval
Test Name	: 6349/1, NICHOLSON ROAD, AME	Value		
Test Name ANT HIV 1/2 AND P24 AN		Value / VIRUS (HI 0.14		
Test Name ANT HIV 1/2 AND P24 AN by CMIA (CHEMILUMIN HIV 1/2 AND P24 AN	HUMAN IMMUNODEFICIENCY	Value / VIRUS (HI 0.14) NON - REA	IV) DUO ULTRA WITH S/CO	(P-24 ANTIGEN DETECTION) NEGATIVE: < 1.00
Test Name ANT HIV 1/2 AND P24 AN by CMIA (CHEMILUMIN HIV 1/2 AND P24 AN by CMIA (CHEMILUMIN INTERPRETATION:- RESUL	I HUMAN IMMUNODEFICIENC ITIGEN: SERUM <i>IESCENT MICROPARTICLE IMMUNOASSAY</i> ITIGEN RESULT	Value / VIRUS (HI 0.14) NON - REA	IV) DUO ULTRA WITH S/CO	(P-24 ANTIGEN DETECTION) NEGATIVE: < 1.00

exposed to HIV 1/2 infection or the sample has been tested during the "window phase" i.e. before the development of detectable levels of antibodies. Hence a Non Reactive result does not exclude the possibility of exposure or infection with HIV 1/2. **RECOMMENDATIONS:**

Results to be clinically correlated
 Rarely falsenegativity/positivity may occur.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com







		hopra & Microbiology) nsultant Patholog		(Pathology)
NAME	: Mrs. MAMTA			
AGE/ GENDER	: 33 YRS/FEMALE		PATIENT ID	: 1549190
COLLECTED BY	:		REG. NO./LAB NO.	: 012407150048
REFERRED BY	: LOOMBA HOSPITAL (AMBA	ALA CANTT)	REGISTRATION DATE	: 15/Jul/2024 11:20 AM
BARCODE NO.	:01513194	<i>,</i>	COLLECTION DATE	: 15/Jul/2024 11:22AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		REPORTING DATE	: 15/Jul/2024 01:11PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD		۳	
			1	
	. 00 10/ 1, 110110110110110110	, multiner or in i	1	
Test Name		Value	Unit	Biological Reference interval
		Value	Unit	
Test Name	HEPAT	Value ITIS B SURFAC	Unit CE ANTIGEN (HBsAg) UL	TRA
Test Name HEPATITIS B SURFA		Value	Unit	
Test Name HEPATITIS B SURFA SERUM	HEPAT	Value ITIS B SURFAC 0.01	Unit CE ANTIGEN (HBsAg) UL	TRA NEGATIVE: < 1.0
Test Name HEPATITIS B SURFA SERUM by CMIA (CHEMILUMII	HEPAT CE ANTIGEN (HBsAg):	Value ITIS B SURFAC 0.01 ASSAY)	Unit CE ANTIGEN (HBsAg) UL	TRA NEGATIVE: < 1.0
Test Name HEPATITIS B SURFA SERUM <i>by CMIA (CHEMILUMII</i> HEPATITIS B SURFA RESULT	HEPAT CE ANTIGEN (HBSAg): NESCENT MICROPARTICLE IMMUNO, CE ANTIGEN (HBSAg)	Value ITIS B SURFAC 0.01 ASSAY) NON - R	Unit CE ANTIGEN (HBsAg) UL S/CO	TRA NEGATIVE: < 1.0
Test Name HEPATITIS B SURFA SERUM <i>by CMIA (CHEMILUMII</i> HEPATITIS B SURFA RESULT <i>by CMIA (CHEMILUMII</i>	HEPAT CE ANTIGEN (HBsAg): Nescent Microparticle IMMUNO	Value ITIS B SURFAC 0.01 ASSAY) NON - R	Unit CE ANTIGEN (HBsAg) UL S/CO	TRA NEGATIVE: < 1.0
Test Name HEPATITIS B SURFA SERUM <i>by CMIA (CHEMILUMII</i> HEPATITIS B SURFA RESULT <i>by CMIA (CHEMILUMII</i> <u>INTERPRETATION:</u>	HEPAT CE ANTIGEN (HBSAg): NESCENT MICROPARTICLE IMMUNO, CE ANTIGEN (HBSAg) NESCENT MICROPARTICLE IMMUNO,	Value ITIS B SURFAC 0.01 ASSAY) NON - R	Unit CE ANTIGEN (HBsAg) UL S/CO EACTIVE	TRA NEGATIVE: < 1.0
Test Name HEPATITIS B SURFA SERUM <i>by CMIA (CHEMILUMII</i> HEPATITIS B SURFA RESULT <i>by CMIA (CHEMILUMII</i> <u>INTERPRETATION:</u> RESU	HEPAT CE ANTIGEN (HBSAg): NESCENT MICROPARTICLE IMMUNO, CE ANTIGEN (HBSAg)	Value ITIS B SURFAC 0.01 ASSAY) NON - R	Unit CE ANTIGEN (HBsAg) UL S/CO	TRA NEGATIVE: < 1.0

Hepatitis B Virus (HBV) is a member of the Hepadna virus family causing infection of the liver with extremely variable clinical features. Hepatitis B is transmitted primarily by body fluids especially serum and also spread effectively sexually and from mother to baby. In most individuals HBV hepatitis is self limiting, but 1-2 % normal adolescent and adults develop Chronic Hepatitis. Frequency of chronic HBV infection is 5-10% in immunocompromised patients and 80 % neonates. The initial serological marker of acute infection is HBsAg which typically appears 2-3 months after infection and disappears 12-20 weeks after onset of symtoms. Persistence of HBsAg for more than 6 months indicates carrier state or Chronic Liver disease.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) V DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

 KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana

 KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana

 0171-2643898, +91 99910 43898
 care@koshealthcare.com

 www.koshealthcare.com
 www.koshealthcare.com



Page 4 of 5





		Chopra gy & Microbiology) Consultant Pathologist	Dr. Yugan MD CEO & Consultant	(Pathology)
IAME	: Mrs. MAMTA			
GE/ GENDER	: 33 YRS/FEMALE	PAT	FIENT ID	: 1549190
COLLECTED BY	:	REG	G. NO./LAB NO.	: 012407150048
REFERRED BY	: LOOMBA HOSPITAL (AMI	BALA CANTT) REG	GISTRATION DATE	: 15/Jul/2024 11:20 AM
ARCODE NO.	:01513194	COL	LECTION DATE	: 15/Jul/2024 11:22AM
LIENT CODE.	: KOS DIAGNOSTIC LAB	RE	PORTING DATE	: 15/Jul/2024 11:38AM
LIENT ADDRESS	: 6349/1, NICHOLSON ROA	AD, AMBALA CANTT		
Test Name		Value	Unit	Biological Reference interval
		VDR	L	
'DRL		NON REACTIV	Έ	NON REACTIVE
DRL by IMMUNOCHROMA	TOGRAPHY	NON REACTIV	Έ	NON REACTIVE
by IMMUNOCHROMAT			Ϋ́Ε	NON REACTIVE
by IMMUNOCHROMA <u>NTERPRETATION:</u> .Does not become p .High titer (>1:16) -	positive until 7 - 10 days after <i>active disease.</i>	appearance ofchancre.		NON REACTIVE
by IMMUNOCHROMA <u>NTERPRETATION:</u> .Does not become p .High titer (>1:16) - b.Low titer (<1:8) - b	positive until 7 - 10 days after active disease. iological falsepositive test in 9	appearance ofchancre. 0% cases or due to late o	r late latent syphillis.	NON REACTIVE
by IMMUNOCHROMA <u>NTERPRETATION</u> : .Does not become p .High titer (>1:16) - b.Low titer (<1:8) - b .Treatment of prim 5.Rising titer (4X) inc	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse,reinfection, or 1	appearance ofchancre. 0% cases or due to late o l e decline tonegative VDF treatment failure and nee	r late latent syphillis. RL within 2 years. ed for retreatment.	NON REACTIVE
by IMMUNOCHROMAT NTERPRETATION: Does not become p Ligh titer (>1:16) - Low titer (<1:8) - b Treatment of prim Strising titer (4X) inco May benonreactive	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse,reinfection, or t e in early primary, late latent,	appearance ofchancre. 0% cases or due to late o e decline tonegative VDF treatment failure and nee and late syphillis (appro	r late latent syphillis. RL within 2 years. ed for retreatment. x. 25% ofcases).	
by IMMUNOCHROMAT <u>NTERPRETATION</u> : .Does not become p .High titer (>1:16) - .Low titer (<1:8) - b .Treatment of prim .Rising titer (4X) inco .May benonreactive .Reactive and weak .HORTTERM FALSE P	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse,reinfection, or t e in early primary, late latent, cly reactive tests should always OSITIVE TEST RESULTS (<6 MOR	appearance ofchancre. 0% cases or due to late o re decline tonegative VDF treatment failure and nee and late syphillis (appro 5 be confirmedwith FTA-A NTHS DURATION) MAY OC	r late latent syphillis. RL within 2 years. ed for retreatment. x. 25% ofcases). BS (fluorescent trepon e	
by IMMUNOCHROMAT <u>NTERPRETATION</u> : .Does not become p .High titer (>1:16) - b.Low titer (<1:8) - b .Treatment of prim .Rising titer (4X) inc .May benonreactiv. .Reactive and weak CHORTTERM FALSE P .Acute viral illnesse	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse, reinfection, or t e in early primary, late latent, cly reactive tests should always OSITIVE TEST RESULTS (<6 MOI es (e.g., hepatitis, measles, info	appearance ofchancre. 0% cases or due to late o re decline tonegative VDF treatment failure and nee and late syphillis (appro 5 be confirmedwith FTA-A NTHS DURATION) MAY OC	r late latent syphillis. RL within 2 years. ed for retreatment. x. 25% ofcases). BS (fluorescent trepon e	
by IMMUNOCHROMAT <u>NTERPRETATION</u> : .Does not become p .High titer (>1:16) - .Low titer (<1:8) - b .Treatment of prim .Rising titer (4X) inc .May benonreactive .Reactive and weak CHORTTERM FALSE P .Acute viral illnesse .M. pneumoniae; C	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse, reinfection, or t e in early primary, late latent, cly reactive tests should always OSITIVE TEST RESULTS (<6 MOI es (e.g., hepatitis, measles, infi- hlamydia; Malaria infection.	appearance ofchancre. 0% cases or due to late o re decline tonegative VDF treatment failure and nee and late syphillis (appro 5 be confirmedwith FTA-A NTHS DURATION) MAY OC	r late latent syphillis. RL within 2 years. ed for retreatment. x. 25% ofcases). BS (fluorescent trepon e	
by IMMUNOCHROMAT <u>NTERPRETATION</u> : .Does not become p .High titer (>1:16) - b.Low titer (<1:8) - b .Treatment of prim .Rising titer (4X) inc .May benonreactiv. .Reactive and weak CHORTTERM FALSE P .Acute viral illnesse	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse, reinfection, or t e in early primary, late latent, cly reactive tests should always OSITIVE TEST RESULTS (<6 MOI es (e.g., hepatitis, measles, infi- hlamydia; Malaria infection.	appearance ofchancre. 0% cases or due to late o re decline tonegative VDF treatment failure and nee and late syphillis (appro 5 be confirmedwith FTA-A NTHS DURATION) MAY OC	r late latent syphillis. RL within 2 years. ed for retreatment. x. 25% ofcases). BS (fluorescent trepon e	
by IMMUNOCHROMAT <u>NTERPRETATION:</u> .Does not become p .High titer (>1:16) - .Low titer (<1:8) - b .Treatment of prim .Rising titer (4X) inc .May benonreactive .May benonreactive .Reactive and weak .HORTTERM FALSE P .Acute viral illnesse .M. pneumoniae; C .Some immunizatio .Pregnancy (rare)	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse,reinfection, or t e in early primary, late latent, cly reactive tests should always OSITIVE TEST RESULTS (<6 MOI es (e.g., hepatitis, measles, infi- hlamydia; Malaria infection.	appearance ofchancre. 0% cases or due to late o re decline tonegative VDF treatment failure and nee and late syphillis (appro 5 be confirmedwith FTA-A NTHS DURATION) MAY OC ectious mononucleosis)	r late latent syphillis. RL within 2 years. ed for retreatment. x. 25% ofcases). BS (fluorescent trepone CCURIN:	
by IMMUNOCHROMAT <u>NTERPRETATION:</u> .Does not become p .High titer (>1:16) - .Low titer (<1:8) - b .Treatment of prim .Rising titer (4X) inc .May benonreactiv. Reactive and weak HORTTERM FALSE P .Acute viral illnesse .M. pneumoniae; C .Some immunizatio .Pregnancy (rare) ONGTERM FALSE PC .Serious underlying	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse, reinfection, or t e in early primary, late latent, cly reactive tests should always OSITIVE TEST RESULTS (<6 MON es (e.g., hepatitis, measles, infi- hlamydia; Malaria infection. ms	appearance ofchancre. 0% cases or due to late o the decline tonegative VDF treatment failure and nee and late syphillis (appro 5 be confirmedwith FTA-A NTHS DURATION) MAY OC ectious mononucleosis)	r late latent syphillis. RL within 2 years. ed for retreatment. x. 25% ofcases). BS (fluorescent trepone CCURIN:	
by IMMUNOCHROMAT <u>NTERPRETATION:</u> .Does not become p .High titer (>1:16) - .Low titer (<1:8) - b .Treatment of prim .Rising titer (4X) inc .May benonreactiv. Reactive and weak CHORTTERM FALSE P .Acute viral illnesse .M. pneumoniae; C .Some immunizatio .Pregnancy (rare) ONGTERM FALSE PC .Serious underlying .Intravenous drug u	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse, reinfection, or t e in early primary, late latent, cly reactive tests should always OSITIVE TEST RESULTS (<6 MON es (e.g., hepatitis, measles, infi- hlamydia; Malaria infection. ms	appearance ofchancre. 0% cases or due to late o the decline tonegative VDF treatment failure and nee and late syphillis (appro 5 be confirmedwith FTA-A NTHS DURATION) MAY OC ectious mononucleosis)	r late latent syphillis. RL within 2 years. ed for retreatment. x. 25% ofcases). BS (fluorescent trepone CCURIN:	
by IMMUNOCHROMAT <u>NTERPRETATION:</u> .Does not become p .High titer (>1:16) - .Low titer (<1:8) - b .Treatment of prim .Rising titer (4X) inc .May benonreactiv. Reactive and weak ChORTTERM FALSE P .Acute viral illnesse .M. pneumoniae; C .Some immunizatio .Pregnancy (rare) ONGTERM FALSE PC .Serious underlying .Intravenous drug u .Rheumatoid arthri	positive until 7 - 10 days after active disease. iological falsepositive test in 9 ary syphillis causes progressiv dicates relapse, reinfection, or t e in early primary, late latent, cly reactive tests should always OSITIVE TEST RESULTS (<6 MON es (e.g., hepatitis, measles, infi- hlamydia; Malaria infection. ms	appearance ofchancre. 0% cases or due to late o the decline tonegative VDF treatment failure and nee and late syphillis (appro 5 be confirmedwith FTA-A NTHS DURATION) MAY OC ectious mononucleosis)	r late latent syphillis. RL within 2 years. ed for retreatment. x. 25% ofcases). BS (fluorescent trepone CCURIN:	

*** End Of Report ***





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com

