

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	: Mr. BALJINDER SINGH	PATIENT ID	: 1556693
AGE/ GENDER	: 44 YRS/MALE	REG. NO./LAB NO.	: 012407220044
COLLECTED BY	:	REGISTRATION DATE	: 22/Jul/2024 12:56 PM
REFERRED BY	:	COLLECTION DATE	: 22/Jul/2024 12:57PM
BARCODE NO.	: 01513623	REPORTING DATE	: 22/Jul/2024 01:53PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

VIRAL MARKERS COMBO PANEL: 2.0

HEPATITIS C VIRUS (HCV) ANTIBODY: TOTAL

HEPATITIS C ANTIBODY (HCV) TOTAL: SERUM 0.12 S/CO NEGATIVE: < 1.00
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY) POSITIVE: > 1.00

HEPATITIS C ANTIBODY (HCV) TOTAL NON - REACTIVE
RESULT

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

INTERPRETATION:-

RESULT (INDEX)	REMARKS
< 1.00	NON - REACTIVE/NOT - DETECTED
> =1.00	REACTIVE/ASYMPTOMATIC/INFECTIVE STATE/CARRIER STATE.

Hepatitis C (HCV) is an RNA virus of Favivirus group transmitted via blood transfusions, transplantation, injection drug abusers, accidental needle punctures in healthcare workers, dialysis patients and rarely from mother to infant. 10 % of new cases show sexual transmission. As compared to HAV & HBV , chronic infection with HCV occurs in 85 % of infected individuals. In high risk population, the predictive value of Anti HCV for HCV infection is > 99% whereas in low risk populations it is only 25 %.

USES:

1. Indicator of past or present infection, but does not differentiate between Acute/ Chronic/Resolved Infection.
2. Routine screening of low and high prevalence population including blood donors.

NOTE:

1. False positive results are seen in Auto-immune disease, Rheumatoid Factor, HYpergammaglobulinemia, Paraproteinemia, Passive antibody transfer, Anti-idiotypes and Anti-superoxide dismutase.
2. False negative results are seen in early Acute infection, Immunosuppression and Immuno— incompetence.
3. HCV-RNA PCR recommended in all reactive results to differentiate between past and present infection.





DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)



DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mr. BALJINDER SINGH	PATIENT ID	: 1556693
AGE/ GENDER	: 44 YRS/MALE	REG. NO./LAB NO.	: 012407220044
COLLECTED BY	:	REGISTRATION DATE	: 22/Jul/2024 12:56 PM
REFERRED BY	:	COLLECTION DATE	: 22/Jul/2024 12:57PM
BARCODE NO.	: 01513623	REPORTING DATE	: 22/Jul/2024 03:05PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

ANTI HUMAN IMMUNODEFICIENCY VIRUS (HIV) DUO ULTRA WITH (P-24 ANTIGEN DETECTION)

HIV 1/2 AND P24 ANTIGEN: SERUM	0.26	S/CO	NEGATIVE: < 1.00 POSITIVE: > 1.00
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			

HIV 1/2 AND P24 ANTIGEN RESULT	NON - REACTIVE
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	

INTERPRETATION:-


RESULT (INDEX)	REMARKS
< 1.00	NON - REACTIVE
> = 1.00	PROVISIONALLY REACTIVE


Non-Reactive result implies that antibodies to HIV 1/ 2 have not been detected in the sample . This means that patient has either not been exposed to HIV 1/ 2 infection or the sample has been tested during the "window phase" i.e. before the development of detectable levels of antibodies. Hence a Non Reactive result does not exclude the possibility of exposure or infection with HIV 1/ 2.

RECOMMENDATIONS:

- Results to be clinically correlated
- Rarely falsenegativity/positivity may occur.




DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mr. BALJINDER SINGH	PATIENT ID	: 1556693
AGE/ GENDER	: 44 YRS/MALE	REG. NO./LAB NO.	: 012407220044
COLLECTED BY	:	REGISTRATION DATE	: 22/Jul/2024 12:56 PM
REFERRED BY	:	COLLECTION DATE	: 22/Jul/2024 12:57PM
BARCODE NO.	: 01513623	REPORTING DATE	: 22/Jul/2024 01:53PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

HEPATITIS B SURFACE ANTIGEN (HBsAg) ULTRA

HEPATITIS B SURFACE ANTIGEN (HBsAg): 0.16 S/CO
 SERUM
 NEGATIVE: < 1.0
 POSITIVE: > 1.0

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

HEPATITIS B SURFACE ANTIGEN (HBsAg) NON REACTIVE
 RESULT


by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)


INTERPRETATION:

RESULT IN INDEX VALUE	REMARKS
< 1.30	NEGATIVE (-ve)
>=1.30	POSITIVE (+ve)

Hepatitis B Virus (HBV) is a member of the Hepadna virus family causing infection of the liver with extremely variable clinical features. Hepatitis B is transmitted primarily by body fluids especially serum and also spread effectively sexually and from mother to baby. In most individuals HBV hepatitis is self limiting, but 1-2 % normal adolescent and adults develop Chronic Hepatitis. Frequency of chronic HBV infection is 5-10% in immunocompromised patients and 80 % neonates. The initial serological marker of acute infection is HBsAg which typically appears 2-3 months after infection and disappears 12-20 weeks after onset of symptoms. Persistence of HBsAg for more than 6 months indicates carrier state or Chronic Liver disease.




DR. VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR. YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mr. BALJINDER SINGH	PATIENT ID	: 1556693
AGE/ GENDER	: 44 YRS/MALE	REG. NO./LAB NO.	: 012407220044
COLLECTED BY	:	REGISTRATION DATE	: 22/Jul/2024 12:56 PM
REFERRED BY	:	COLLECTION DATE	: 22/Jul/2024 12:57 PM
BARCODE NO.	: 01513623	REPORTING DATE	: 22/Jul/2024 01:17 PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

VDRL by IMMUNOCHROMATOGRAPHY	NON REACTIVE		NON REACTIVE
---------------------------------	--------------	--	--------------

INTERPRETATION:

- Does not become positive until 7 - 10 days after appearance of chancre.
- High titer (>1:16) - active disease.**
- Low titer (<1:8) - biological falsepositive test in 90% cases or due to late or late latent syphilis.**
- Treatment of primary syphilis causes progressive decline to negative VDRL within 2 years.
- Rising titer (4X) indicates relapse, reinfection, or treatment failure and need for retreatment.
- May be nonreactive in early primary, late latent, and late syphilis (approx. 25% of cases).
- Reactive and weakly reactive tests should always be confirmed with FTA-ABS (fluorescent treponemal antibody absorption test).**

SHORT TERM FALSE POSITIVE TEST RESULTS (<6 MONTHS DURATION) MAY OCCUR IN:

- Acute viral illnesses (e.g., hepatitis, measles, infectious mononucleosis)
- M. pneumoniae; Chlamydia; Malaria infection.
- Some immunizations
- Pregnancy (rare)

LONG TERM FALSE POSITIVE TEST RESULTS (>6 MONTHS DURATION) MAY OCCUR IN:

- Serious underlying disease e.g., collagen vascular diseases, leprosy, malignancy.
- Intravenous drug users.
- Rheumatoid arthritis, thyroiditis, AIDS, Sjogren's syndrome.
- <10 % of patients older than age 70 years.
- Patients taking some anti-hypertensive drugs.

*** End Of Report ***




 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)

