Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. ISHA SOOD**		
AGE/ GENDER	: 42 YRS/FEMALE	**PATIENT ID**	: 1562181
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012407270018**
REFERRED BY	:	**REGISTRATION DATE**	: 27/Jul/2024 09:33 AM
BARCODE NO.	: 01513897	**COLLECTION DATE**	: 27/Jul/2024 09:36AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Jul/2024 10:54AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

CLINICAL CHEMISTRY/BIOCHEMISTRY

GLUCOSE FASTING (F)

GLUCOSE FASTING (F): PLASMA *by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)*	91.57	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0

INTERPRETATION

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.
2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mrs. ISHA SOOD**		
AGE/ GENDER	: 42 YRS/FEMALE	**PATIENT ID**	: 1562181
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012407270018**
REFERRED BY	:	**REGISTRATION DATE**	: 27/Jul/2024 09:33 AM
BARCODE NO.	: 01513897	**COLLECTION DATE**	: 27/Jul/2024 09:36AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Jul/2024 10:54AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

CALCIUM

CALCIUM: SERUM	9.4	mg/dL	8.50 - 10.60
by ARSENAZO III, SPECTROPHOTOMETRY			

INTERPRETATION:-

1.Serum calcium (total) estimation is used for the diagnosis and monitoring of a wide range of disorders including diseases of bone, kidney, parathyroid gland, or gastrointestinal tract.
2. Calcium levels may also reflect abnormal vitamin D or protein levels.
3.The calcium content of an adult is somewhat over 1 kg (about 2% of the body weight).Of this, 99% is present as calcium hydroxyapatite in bones and <1% is present in the extra-osseous intracellular space or extracellular space (ECS).
4. In serum, calcium is bound to a considerable extent to proteins (approximately 40%), 10% is in the form of inorganic complexes, and 50% is present as free or ionized calcium.
NOTE:-Calcium ions affect the contractility of the heart and the skeletal musculature, and are essential for the function of the nervous system. In addition, calcium ions play an important role in blood clotting and bone mineralization.

HYPOCALCEMIA (LOW CALCIUM LEVELS) CAUSES :-

1.Due to the absence or impaired function of the parathyroid glands or impaired vitamin-D synthesis.
2. Chronic renal failure is also frequently associated with hypocalcemia due to decreased vitamin-D synthesis as well as hyperphosphatemia and skeletal resistance to the action of parathyroid hormone (PTH).
3.***NOTE:-*** A characteristic symptom of hypocalcemia is latent or manifest tetany and osteomalacia.

HYPERCALCEMIA (INCREASE CALCIUM LEVELS) CAUSES:-

1.Increased mobilization of calcium from the skeletal system or increased intestinal absorption.
2.Primary hyperparathyroidism (pHPT)
3.Bone metastasis of carcinoma of the breast, prostate, thyroid gland, or lung.
NOTE:-Severe hypercalcemia may result in cardiac arrhythmia.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. ISHA SOOD**		
AGE/ GENDER	: 42 YRS/FEMALE	**PATIENT ID**	: 1562181
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012407270018**
REFERRED BY	:	**REGISTRATION DATE**	: 27/Jul/2024 09:33 AM
BARCODE NO.	: 01513897	**COLLECTION DATE**	: 27/Jul/2024 09:36AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Jul/2024 10:54AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	0.925	ng/mL	0.35 - 1.93
THYROXINE (T4): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	7.26	µgm/dL	4.87 - 12.60
THYROID STIMULATING HORMONE (TSH): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	1.536	µIU/mL	0.35 - 5.50

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

*TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%.Hence time of the day has influence on the measured serum TSH concentrations.*TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4)and trilodothyronine (T3).Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG),and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (eg: phenytoin , salicylates).
3. Serum T4 levles in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.
4. TSH may be normal in central hypothyroidism , recent rapid correction of hyperthyroidism or hypothroidism , pregnancy , phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	**Refferance Range (ng/mL)**	**Age**	**Refferance Range (µg/dL)**	**Age**	**Reference Range (µIU/mL)**
0 - 7 Days	**0.20 - 2.65**	**0 - 7 Days**	**5.90 – 18.58**	**0 - 7 Days**	**2.43 - 24.3**
7 Days - 3 Months	**0.36 - 2.59**	**7 Days - 3 Months**	**6.39 - 17.66**	**7 Days - 3 Months**	**0.58 - 11.00**
3 - 6 Months	**0.51 - 2.52**	**3 - 6 Months**	**6.75 – 17.04**	**3 Days – 6 Months**	**0.70 - 8.40**

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. ISHA SOOD**		
AGE/ GENDER	: 42 YRS/FEMALE	**PATIENT ID**	: 1562181
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012407270018**
REFERRED BY	:	**REGISTRATION DATE**	: 27/Jul/2024 09:33 AM
BARCODE NO.	: 01513897	**COLLECTION DATE**	: 27/Jul/2024 09:36AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Jul/2024 10:54AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name			Value	Unit	Biological Reference interval
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 – 5.50
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35– 5.50
RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (µIU/mL)					
1st Trimester			0.10 – 2.50		
2nd Trimester			0.20 – 3.00		
3rd Trimester			0.30 – 4.10		

INCREASED TSH LEVELS:
1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
2.Hypothyroid patients receiving insufficient thyroid replacement therapy.
3.Hashimotos thyroiditis
4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.
5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:
1.Toxic multi-nodular goitre & Thyroiditis.
2.Over replacement of thyroid harmone in treatment of hypothyroidism.
3.Autonomously functioning Thyroid adenoma
4.Secondary pituatary or hypothalmic hypothyroidism
5.Acute psychiatric illness
6.Severe dehydration.
7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.
8.Pregnancy: 1st and 2nd Trimester

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)