



	MD (Pathology & M	Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist		n Chopra (Pathology) Pathologist	
NAME	: Mrs. KANIKA				
AGE/ GENDER	: 34 YRS/FEMALE		PATIENT ID	: 1564812	
COLLECTED BY	:		REG. NO./LAB NO.	: 012407290068	
REFERRED BY	:		REGISTRATION DATE	: 29/Jul/2024 07:01 PM	
BARCODE NO.	: 01514092		COLLECTION DATE	: 29/Jul/2024 07:02PM	
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPORTING DATE		: 29/Jul/2024 09:28PM	
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AM	IBALA CANTT			
Test Name		Value	Unit	Biological Reference interval	
		ENDOC	RINOLOGY		
	TH	YROID FUNC	CTION TEST: TOTAL		
TRIIODOTHYRONINE	(T3): SERUM escent microparticle immunoassa	0.769	ng/mL	0.35 - 1.93	
THYROXINE (T4): SEF		10.93	µgm/dL	4.87 - 12.60	
	NG HORMONE (TSH): SERUM escent microparticle immunoassa rasensitive	3.121 (Y)	μIU/mL	0.35 - 5.50	

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.

2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (eg: phenytoin , salicylates).

3. Serum T4 levles in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.

4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)		
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (μIU/mL)	
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3	
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00	
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days – 6 Months	0.70 - 8.40	





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

 KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana

 KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana

 0171-2643898, +91 99910 43898
 care@koshealthcare.com
 www.koshealthcare.com



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT





	Dr. Vinay Chopra MD (Pathology & Microbiolog Chairman & Consultant Patho		(Pathology)
NAME	: Mrs. KANIKA		
AGE/ GENDER	: 34 YRS/FEMALE	PATIENT ID	: 1564812
COLLECTED BY	:	REG. NO./LAB NO.	: 012407290068
REFERRED BY	:	REGISTRATION DATE	: 29/Jul/2024 07:01 PM
BARCODE NO.	: 01514092	COLLECTION DATE	: 29/Jul/2024 07:02PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPORTING DATE	: 29/Jul/2024 09:28PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CA	NTT	

Test Name			Value	Unit	:	Biological Reference interva
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00	
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50	
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35- 5.50	
	RECO	MMENDATIONS OF TSH LI	EVELS DURING PRE	GNANCY (µIU/mL)	•	
1st Trimester			0.10 - 2.50			
2nd Trimester			0.20 - 3.00			
3rd Trimester			0.30 - 4.10			

INCREASED TSH LEVELS:

1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.

2. Hypothyroid patients receiving insufficient thyroid replacement therapy.

3.Hashimotos thyroiditis

4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.

5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

1.Toxic multi-nodular goitre & Thyroiditis.

2. Over replacement of thyroid harmone in treatment of hypothyroidism.

3. Autonomously functioning Thyroid adenoma

4. Secondary pituatary or hypothalmic hypothyroidism

5. Acute psychiatric illness

6.Severe dehydration.

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8.Pregnancy: 1st and 2nd Trimester

*** End Of Report **





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

