

# **KOS Diagnostic Lab**

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 05/Aug/2024 09:32AM

**NAME** : Mrs. KAMLA SURI

**AGE/ GENDER** : 80 YRS/FEMALE **PATIENT ID** : 1570677

**COLLECTED BY** :012408050019 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 05/Aug/2024 08:08 AM BARCODE NO. :01514486 **COLLECTION DATE** : 05/Aug/2024 08:12AM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

## **ENDOCRINOLOGY**

REPORTING DATE

## THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 1.32 ng/mL 0.35 - 1.93by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROXINE (T4): SERUM 4.87 - 12.60 7.56 μgm/dL

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROID STIMULATING HORMONE (TSH): SERUM 1.513 μIU/mL 0.35 - 5.50

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

## INTERPRETATION:

CLIENT CODE.

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and trilodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs
- 3. Serum T4 levies in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (μg/dL)	Age	Reference Range ( μΙυ/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





# **KOS Diagnostic Lab**

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 05/Aug/2024 09:32AM

: Mrs. KAMLA SURI **NAME** 

**AGE/ GENDER** : 80 YRS/FEMALE **PATIENT ID** : 1570677

**COLLECTED BY** REG. NO./LAB NO. :012408050019

REFERRED BY **REGISTRATION DATE** : 05/Aug/2024 08:08 AM BARCODE NO. :01514486 **COLLECTION DATE** : 05/Aug/2024 08:12AM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name		Value L			Biological Reference interva
0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00	
0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50	
0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35- 5.50	
RECON	IMENDATIONS OF TSH LE	VELS DURING PREC	GNANCY ( μIU/mL)		
1st Trimester			0.10 - 2.50		
2nd Trimester			0.20 - 3.00		
3rd Trimester			0.30 - 4.10		
	0.92 - 2.28 0.35 - 1.93 0.35 - 1.93 RECOM 1st Trimester 2nd Trimester	0.92 - 2.28	0.74 - 2.40 6 - 12 Months 7.10 - 16.16 0.92 - 2.28 1 - 10 Years 6.00 - 13.80 0.35 - 1.93 11 - 19 Years 4.87 - 13.20 0.35 - 1.93 > 20 Years (Adults) 4.87 - 12.60 RECOMMENDATIONS OF TSH LEVELS DURING PRECOUNTY PREC	0.74 - 2.40       6 - 12 Months       7.10 - 16.16       6 - 12 Months         0.92 - 2.28       1 - 10 Years       6.00 - 13.80       1 - 10 Years         0.35 - 1.93       11 - 19 Years       4.87 - 13.20       11 - 19 Years         0.35 - 1.93       > 20 Years (Adults)       4.87 - 12.60       > 20 Years (Adults)         RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (μIU/mL)         1st Trimester       0.10 - 2.50         2nd Trimester       0.20 - 3.00	0.74 - 2.40   6 - 12 Months   7.10 - 16.16   6 - 12 Months   0.70 - 7.00     0.92 - 2.28   1 - 10 Years   6.00 - 13.80   1 - 10 Years   0.60 - 5.50     0.35 - 1.93   11 - 19 Years   4.87 - 13.20   11 - 19 Years   0.50 - 5.50     0.35 - 1.93   > 20 Years (Adults)   4.87 - 12.60   > 20 Years (Adults)   0.35 - 5.50     RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (μΙU/mL)     1st Trimester   0.10 - 2.50     2nd Trimester   0.20 - 3.00

REPORTING DATE

### **INCREASED TSH LEVELS:**

CLIENT CODE.

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

### **DECREASED TSH LEVELS:**

- 1.Toxic multi-nodular goitre & Thyroiditis.
- 2. Over replacement of thyroid harmone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituatary or hypothalmic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester

\*\*\* End Of Report \*



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana