

Dr. Vinay Chopra  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

Dr. Yugam Chopra  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mr. SACHIN GOEL	<b>PATIENT ID</b>	: 1579037
<b>AGE/ GENDER</b>	: 40 YRS/MALE	<b>REG. NO./LAB NO.</b>	: 012408130023
<b>COLLECTED BY</b>	: SURJESH	<b>REGISTRATION DATE</b>	: 13/Aug/2024 09:38 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 13/Aug/2024 09:47AM
<b>BARCODE NO.</b>	: 01514992	<b>REPORTING DATE</b>	: 13/Aug/2024 03:19PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

### CLINICAL PATHOLOGY

#### MICROALBUMIN/CREATININE RATIO - RANDOM URINE

<b>MICROALBUMIN: RANDOM URINE</b> by SPECTROPHOTOMETRY	455.27 <sup>H</sup>	mg/L	0 - 25
<b>CREATININE: RANDOM URINE</b> by SPECTROPHOTOMETRY	24.59	mg/dL	20 - 320
<b>MICROALBUMIN/CREATININE RATIO - RANDOM URINE</b> by SPECTROPHOTOMETRY	1851.44 <sup>H</sup>	mg/g	0 - 30

#### INTERPRETATION:-

PHYSIOLOGICALLY NORMAL:	mg/L	0 - 30
MICROALBUMINURIA:	mg/L	30 - 300
GROSS PROTEINURIA:	mg/L	> 300

Long standing un-treated Diabetes and Hypertension can lead to renal dysfunction.

2. Diabetic nephropathy or kidney disease is the most common cause of end stage renal disease(ERSD) or kidney failure.

3. Presence of Microalbuminuria is an early indicator of onset of compromised renal function in these patients.

4. Microalbuminuria is the condition when urinary albumin excretion is between 30-300 mg & above this it is called as macroalbuminuria, the presence of which indicates serious kidney disease.

5. Microalbuminuria is not only associated with kidney disease but of cardiovascular disease in patients with diabetes & hypertension.

6. Microalbuminuria reflects vascular damage & appear to be a marker of early arterial disease & endothelial dysfunction.

**NOTE:-** IF A PATIENT HAS = 1+ PROTEINURIA (30 mg/dl OR 300 mg/L) BY URINE DIPSTICK (URINE ANALYSIS), OVERT PROTEINURIA IS PRESENT AND TESTING FOR MICROALBUMIN IS INAPPROPRIATE. IN SUCH A CASE, URINE PROTEIN:CREATININE RATIO OR 24 HOURS TOTAL URINE MICROPROTEIN IS APPROPRIATE.

\*\*\* End Of Report \*\*\*



  
 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)

