

## **KOS Diagnostic Lab**

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mrs. BHAGYA SHRI

**AGE/ GENDER** : 31 YRS/FEMALE **PATIENT ID** : 1584463

**COLLECTED BY** REG. NO./LAB NO. : 012408190025

REFERRED BY **REGISTRATION DATE** : 19/Aug/2024 11:06 AM : 19/Aug/2024 11:17AM BARCODE NO. :01515308 **COLLECTION DATE** 

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

## IMMUNOPATHOLOGY/SEROLOGY ANTI THYROID PEROXIDASE (TPO/AMA) ANTIBODIES

REPORTING DATE

ANTI TPO/AMA ANTIBODIES: SERUM

by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

< 1.00

IU/mL

0.00 - 10.0

: 19/Aug/2024 01:14PM

DIABETES (II): < 25.0

**INTERPRETATION:** 

CLIENT CODE.

1. Thyroperoxidase (TPO) is an enzyme involved inthyroid hormone synthesis, catalyzing the oxidation of iodide on tyrosine residues in thyroglobulin for the synthesis of triiodothyronine and thyroxine (tetraiodothyronine).
2. TPO is a membrane-associated hemo glycoprotein expressed only in thyrocytes and is one of the most important thyroid gland antigens.
3. Anti-TPO is technically superior and a more specific method for measuring thyroid auto-antibodies, It is especially useful in patients presenting with subclinical hypothyroidism where TSH is elevated but Free T4 levels are normal.

**INCREASED LEVELS (Autoimmune thyroid disease):** 

- Hashimoto thyroiditis.
- Idiopathic myxedema.
   Graves disease
- 4. Post-partum thyroiditis.
- 5. Primary hypothyroidism due to Hashimoto thyroiditis.

1. The highest TPO antibody levels are observed in patients suffering from Hashimoto thyroiditis. In this disease, the prevalence of TPO antibodies is about 90% of cases, confirming the autoimmune origin of the disease.

2. These auto-antibodies also frequently occur (60%-80%) in the course of Graves disease.

3. In patients with subclinical hypothyroidism, the presence of TPO antibodies is associated with an increased risk of developing overt

hypothyroidism.

\*\*\* End Of Report \*\*\*



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

