

KOS Diagnostic Lab





Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. UDAY KUMAR

AGE/ GENDER : 70 YRS/MALE **PATIENT ID** : 1585173

COLLECTED BY :012408200012 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Aug/2024 08:25 AM BARCODE NO. :01515340 **COLLECTION DATE** : 20/Aug/2024 12:24PM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 21/Aug/2024 02:36PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Value Unit **Biological Reference interval** Test Name

CLINICAL CHEMISTRY/BIOCHEMISTRY

GLUCOSE FASTING (F) AND POST PRANDIAL (PP)

GLUCOSE FASTING (F): PLASMA mg/dL NORMAL: < 100.0 110.53^H

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0

DIABETIC: > 0R = 126.0

GLUCOSE POST PRANDIAL (PP): PLASMA 162.63^H mg/dL NORMAL: < 140.00

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 140.0 - 200.0

DIABETIC: > 0R = 200.0

INTERPRETATION:

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

- 1. A fasting plasma glucose below 100 mg/dL and post-prandial plasma glucose level below 140 mg/dl is considered normal.
- 2. A fasting plasma glucose level between 100 125 mg/dl and post-prandial plasma glucose level between 140 200 mg/dL is considered as glucose intolerant or pre diabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
- 3. A fasting plasma glucose level of above 125 mg/dL and post-prandial plasma glucose level above 200 mg/dL is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

* End Of Report ***



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

