

Dr. Vinay Chopra  
MD (Pathology & Microbiology)  
Chairman & Consultant Pathologist

Dr. Yugam Chopra  
MD (Pathology)  
CEO & Consultant Pathologist

NAME : Mrs. PARAMJEET KAUR  
AGE/ GENDER : 48 YRS/FEMALE  
COLLECTED BY :  
REFERRED BY : LOOMBA HOSPITAL (AMBALA CANTT)  
BARCODE NO. : 01515833  
CLIENT CODE. : KOS DIAGNOSTIC LAB  
CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT  
PATIENT ID : 1591427  
REG. NO./LAB NO. : 012408270063  
REGISTRATION DATE : 27/Aug/2024 06:33 PM  
COLLECTION DATE : 27/Aug/2024 06:33PM  
REPORTING DATE : 30/Aug/2024 07:28AM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## MICROBIOLOGY

### CULTURE AEROBIC BACTERIA AND ANTIBIOTIC SENSITIVITY: URINE

#### CULTURE AND SUSCEPTIBILITY: URINE

DATE OF SAMPLE 27-08-2024  
SPECIMEN SOURCE URINE  
INCUBATION PERIOD 48 HOURS  
by AUTOMATED BROTH CULTURE  
CULTURE STERILE  
by AUTOMATED BROTH CULTURE  
ORGANISM NO AEROBIC PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF INCUBATION AT 37°C  
by AUTOMATED BROTH CULTURE

#### AEROBIC SUSCEPTIBILITY: URINE

##### INTERPRETATION:

1. In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection.
2. Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out" catheterization or from patients with indwelling catheters.

##### SUSCEPTIBILITY:

1. A test interpreted as **SENSITIVE** implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent recommended for that type of infection and infecting species, unless otherwise indicated..
2. A test interpreted as **INTERMEDIATE** implies that the "infection due to the isolate may be appropriately treated in body sites where the drugs are physiologically concentrated or when a high dosage of drug can be used".
3. A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

##### CAUTION:

Conditions which can cause a false Negative culture:

1. Patient is on antibiotics. Please repeat culture post therapy.
2. Anaerobic bacterial infection.
3. Fastidious aerobic bacteria which are not able to grow on routine culture media.
4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.
5. Renal tuberculosis to be confirmed by AFB studies.

\*\*\* End Of Report \*\*\*



DR. VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR. YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)

