

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.



KOS Diagnostic Lab (A Unit of KOS Healthcare)

	Dr. Vinay Chopra MD (Pathology & Microbiology		Dr. Yugam Chopra MD (Pathology)	
		onsultant Pathologist		
NAME	: Mr. OM PARKASH			
AGE/ GENDER	: 53 YRS/MALE		PATIENT ID	: 1603765
COLLECTED BY	:		REG. NO./LAB NO.	: 012409060001
REFERRED BY	:		<b>REGISTRATION DATE</b>	: 06/Sep/2024 05:53 AM
BARCODE NO.	: 01516384		COLLECTION DATE	: 06/Sep/2024 05:55AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		REPORTING DATE	: 06/Sep/2024 07:20AM
LIENT ADDRESS	: 6349/1, NICHOLSON ROA	D, AMBALA CANTT		
Test Name		Value	Unit	Biological Reference interval
	CLI	NICAL CHEMIS	TRY/BIOCHEMISTR	Y
		GLUCOSE	FASTING (F)	
GLUCOSE FASTING ( by glucose oxidas	F): PLASMA se - peroxidase (god-pod)	121.55 <sup>H</sup>	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0
est (after consumpti . A fasting plasma g	lucose level below 100 mg/dl i lucose level between 100 - 12! ion of 75 gms of glucose) is rec lucose level of above 125 mg/d ing plasma glucose level in exc	commended for all s dl is highly suggestiv	uch patients. /e of diabetic state. A repe	at post-prandial is strongly recommended for a
est (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
est (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
est (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
est (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
est (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
est (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	at post-prandial is strongly recommended for a
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	at post-prandial is strongly recommended for a
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	at post-prandial is strongly recommended for a
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	at post-prandial is strongly recommended for a
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	at post-prandial is strongly recommended for a
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv cess of 125 mg/dl or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	eat post-prandial is strongly recommended for al
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dI is highly suggestiv cess of 125 mg/dI or	uch patients. /e of diabetic state. A repe ) both occasions is confirm	at post-prandial is strongly recommended for a
test (after consumpti 3. A fasting plasma g	lucose level between 100 - 129 ion of 75 gms of glucose) is rec lucose level of above 125 mg/o	commended for all s dl is highly suggestiv ess of 125 mg/dl or *** End Of Re	uch patients. /e of diabetic state. A repe ) both occasions is confirm	at post-prandial is strongly recommended for a

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com

