



	MD (I	Vinay Chopra Pathology & Microbiology) man & Consultant Pathologist		(Pathology)
NAME	: Mr. ROHIT BHUD	IRAJA		
AGE/ GENDER	: 44 YRS/MALE		PATIENT ID	: 1613847
COLLECTED BY	:		REG. NO./LAB NO.	: 012409150016
REFERRED BY	:		REGISTRATION DATE	: 15/Sep/2024 09:10 AM
BARCODE NO.	:01516992		COLLECTION DATE	: 15/Sep/2024 09:14AM
CLIENT CODE.	: KOS DIAGNOSTIC	LAB	REPORTING DATE	: 15/Sep/2024 10:26AM
CLIENT ADDRESS	: 6349/1, NICHOLS	ON ROAD, AMBALA CANTT		
Test Name		Value	Unit	Biological Reference interval
		CLINICAL CHEMIS	TRY/BIOCHEMISTR	Y
		CHOLESTE	ROL: SERUM	

INTERPRETATION:

NATIONAL LIPID ASSOCIATION RECOMMENDATIONS (NLA-2014)	CHOLESTEROL IN ADULTS (mg/dL)	CHOLESTEROL IN ADULTS (mg/dL)	
DESIRABLE	< 200.0	< 170.0	
BORDERLINE HIGH	200.0 - 239.0	171.0 - 199.0	
HIGH	>= 240.0	>= 200.0	

NOTE:

Measurements in the same patient can show physiological & analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.
 As per National Lipid association - 2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com



Page 1 of 2





NAME : Mr. ROHIT BHUDIRALA AGE/ GENDER : 44 YRS/MALE PATIENT ID : 1613847 COLLECTED BY : REG. NO./LAB NO. : 012409150016 REFERED BY : REGISTRATION DATE : 15/Sep/2024 09:10 AM BARCODE NO. : 01516992 COLLECTION DATE : 15/Sep/2024 09:14 AM CLIENT CODE : KOS DIAGNOSTIC LAB REPORTING DATE : 15/Sep/2024 10:26 AM CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT Test Name Value Unit Biological Reference inte URIC ACID URIC ACID URICASED PEROXIDASE <i>NITEMPERTATION:</i> 1.GOUT OCCURS When high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint. 2. Uric Acid is the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the instituinal tract by microbial degradation. <i>NICREASED:</i> <i>(A)</i> DUE TO INCREASED PEROVIDIASE <i>E</i> Polycythemain etc. <i>(B)</i> DUE TO DECREASED EXCREATION (BY KIDNEYS) 1. Jacobia dispastion. 2. This Acid is the and product of matipasia. 5. Psoriasis. 5. Solitele cell anaemia etc. <i>(B)</i> DUE TO DECREASED EXCREATION (BY KIDNEYS) 1. Jactic acidosis. 4. Aspirin Ingestion (Less than 2 grams per day). 5. Diabetic kellency of Zim, En and Millowing Singerse. 4. Spiring Ingestion. 2. This acid asses exclasses. <i>(B)</i> DUE TO DECREASED EXCREATION (BY KIDNEYS) 1. Jactic acidosis. 4. Aspirin Ingestion (Less than 2 grams per day). 5. Diabetic kellency of Zim, En and molyddenum. 2. Thatale diurentes. 4. Appirin Ingestion (Less than 2 grams per day). 5. Diabetic kellency of Zim, En and Millowing disease. 4. Multiple Screases. <i>(M)</i> DUE TO DIETARY DEFICIENCY 1. Dietary deficiency of Zim, En and Millowing disease. 4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc. <i>(B)</i> DUE TO MERASED EXCREATION 1. Dietary deficiency of Zim, En and Millowing disease. 4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc. <i>(B)</i> DUE TO MERASED EXCREATION 1. Dietary deficiency of Zim, En and Millowing disease. 4. Syndrome of inappropri		MD (Pathology a	Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist		Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist		
CULLECTED BY EXECT NO./LAB NO. : 012409150016 REFERRED BY IS REGISTRATION DATE : 15/Sep/2024 09:10 AM BARCODE NO. : 015160992 COLLECTION DATE : 15/Sep/2024 09:14AM CLIENT CODE : KOS DIAGNOSTIC LAB REPORTING DATE : 15/Sep/2024 09:14AM CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT URIC ACID URIC ACID </th <th>NAME</th> <th>: Mr. ROHIT BHUDIRAJA</th> <th></th> <th></th> <th></th>	NAME	: Mr. ROHIT BHUDIRAJA					
REFERENDENT Image: Structure S	AGE/ GENDER	: 44 YRS/MALE	PA	TIENT ID	: 1613847		
BARCODE NO. : 101516992 COLLECTION DATE : 15/Sep/2024 09:14AM CLIENT CODE : KOS DIAGNOSTIC LAB REPORTING DATE : 15/Sep/2024 09:14AM CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT Test Name Value Unit Biological Reference inte URIC ACID URIC ACID URIC ACID: SERUM 6.51 mg/dL 3.60 - 7.70 by URICASE - OXIDASE PEROXIDASE MIREPRETATION: 1. GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint. 2. Uric Acid is the end product of purime metabolism. Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the netsetinal tract by microbial degradation. NCREASED- (A) DUE TO INCREASED PRODUCTION- 1. Idiopatite jurines (organ meats, legumes, anchovies, etc). 3. Stokle cella naemia etc. 8) DUE TO DECREASED EXCREATION (8Y KIDNEYS) 1. Alcohol ingestion. 2. Thaisde diuretics. 3. Stokle cella naemia etc. 8) DUE TO DECREASED EXCREATION (8Y KIDNEYS) 1. Alcohol ingestion. 2. Thaisde diuretics. 3. Diabetic, ketoasidosis or starvation. 3. Diabetic, ketoasidosis or starvation. 3. Diabetic, ketoasis or starvation. 3. Diabetic, ketoasidosis or starvation. 3. Olabetic, ketoasidosis or starvation. 4. Aputer D DIETARY DEFICIENCY 1. Dietary deficiency of Zinc, Iron and molybdenum. 2. Thaized diverties. 3. Multipe sclerosis. 3.	COLLECTED BY	:	RI	EG. NO./LAB NO.	: 012409150016		
ELIENT CODE :: KOS DIAGNOSTIC LAB	REFERRED BY	:	RI	EGISTRATION DATE	: 15/Sep/2024 09:10 AM		
CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT Test Name Value Unit Biological Reference inte LIRIC ACID URIC ACID by URICASE 0.51 mg/dL 3.60 - 7.70 by URICACID 2.01c Acid is the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation. NCREASED NORCEASED Columnation of the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation. NORCEASED Columnation of the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation. Aprice To increases D PRODUCTION: 2.2xcessive dietary purines (organ meats, legumes, anchovies, etc.). 3.0 to increase at myeloid metaplasia. 3.0 to indicate acidosis. 4. Polycythemania vera & myeloid metaplasia. 3.0 to indicate acidosis. 9. Due to increase to contraction (BY KIDNEYS) 3.1 acit a cidosis. 1. Alopicin	BARCODE NO.	:01516992	CC	DLLECTION DATE	: 15/Sep/2024 09:14AM		
Test Name Value Unit Biological Reference inter URIC ACID	CLIENT CODE.	: KOS DIAGNOSTIC LAB	RI	EPORTING DATE	: 15/Sep/2024 10:26AM		
URIC ACID URIC ACID URICASE PEROXIDASE URICASE PEROXIDASE URICASE OXIDASE PEROXIDASE UNICASE - OXIDASE PEROXIDON: - Lidopathic primary gout. 2. Excessive dietary purines (organ meats.legumes.anchovies, etc). 3. Cytolytic treatment of malignancies especially leukemais & lymphomas. 4. Polycythemal vera & myeloid metaplasia. 3. Psortasis. 5. Sickle cell anaemia etc. (B) DUE TO DECREASED EXCREATION (BY KIDNEYS) 1. Alcohol ingestion, (Bast han 2 grams per day). 5. Diabetic ketoacidosis or starvation. 5. Diabetic ketoacidosis or starvation. 5. Diabetic ketoacidosis or starvation. 5. Pactional cell cell cell anaemia etc. (B) DUE TO DICTEASED EXCREATION (BY KIDNEYS) 1. Diabetic ketoacidosis or starvation. 5. Pactore of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc. (B) DUE TO INCREASED EXCREATION 1. Drugs:-Probenecid , sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estrophysical cell cell cell cell cell cell cell c	CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD					
URIC ACID: SERUM 6.51 mg/dL 3.60 - 7.70 by URICASE - OXIDASE PEROXIDASE INTERRETATION: 1.GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint. 2.Uric Acid is the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation. INCREASED: (A) DUE TO INCREASED PRODUCTION: 1.Idiopathic primary gout. 2.Excessive dietary purines (organ meats,legumes,anchovies, etc). 3.Cytolytic treatment of malignancies especially leukemais & lymphomas. 4.Polycythemai vera & myeloid metaplasia. 5.Sickle cell anaemia etc. (B) DUE TO DECREASED EXCREATION (BY KIDNEYS) 1.Alcohol ingestion. 2.Thiazide diuretics. 3.Lactic acidosis. 4.Aspin ingestion (less than 2 grams per day). 5.Diabetic ketoacidosis or starvation. 5.Diabetic ketoacidosis or starvation. 5.Renal failure due to any cause etc. DECREASED: (A) DUE TO DIETARY DEFICIENCY 1.Dietary deficiency of Zinc, Iron and molybdenum. 2.Fanconi syndrome & Wilsons disease. 3.Multiple sclerosis. 4.Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc. (B) DUE TO DIECREASED EXCREATION 1.Duetro INCREASED EXCREATION 1.Dietary deficiency of Zinc, iron and molybdenum. 2.Fanconi syndrome & Wilsons disease. 3.Multiple sclerosis. 4.Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc. (B) DUE TO INCREASED EXCREATION 1.Durgs:-Probenecid , sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estre	Test Name		Value	Unit	Biological Reference interval		
 JRIC ACID: SERUM by URICASE - OXIDASE PEROXIDASE b) OUT COLORS when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint. 2. Uric Acid is the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation. NOREASED: A) DUE TO INCREASED PRODUCTION:- 1. Idiopathic primary gout. 2. Excessive dietary purines (organ meats,legumes,anchovies, etc). 3. Cytolytic treatment of malignancies especially leukemais & lymphomas. 4. Polycythemai vera & myeloid metaplasia. 5. Sickle cell anaemia etc. B) DUE TO DECREASED EXCREATION (BY KIDNEYS) 1. Alcohol ingestion. 2. Thiazide diuretics. 3. Lactic acidosis. 4. Aspinin ingestion (less than 2 grams per day). 5. Diabetic ketoacidosis or starvation. 5. DECREASED: ANDUE TO DIFLARY DEFICIENCY 1. Dietary deficiency of Zinc, Iron and molybdenum. 2. Fanconi syndrome & Wilsons disease. 4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc. B) DUE TO INCREASED EXCREATION 1. Drugs:-Probenecid , sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estructure of the propertice of sulface of the propertice of a grams per day of the propertice of an ACTH, anti-coagulants and estructure of the propertice of sulface of a sulface of the propertice of the prop			URIC	ACID			
by URICASE - OXIDASE PEROXIDASE INTERPETATION:- 1. GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint. 2. Uric Acid is the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the ntestinal tract by microbial degradation. INCREASED:- (A),DUE TO INCREASED PRODUCTION:- 1.1diopathic primary gout. 2. Excessive dietary purines (organ meats, legumes, anchovies, etc). 3. Cytolytic treatment of malignancies especially leukemais & lymphomas. 4. Polycythemai vera & myeloid metaplasia. 5. Psoriasis. 5. Sickle cell anaemia etc. (B),DUE TO DECREASED EXCREATION (BY KIDNEYS) 1. Alcohol ingestion. 2. Thiazide diuretics. 3. Lactic acidosis. 4. Aspirin ingestion (less than 2 grams per day). 5. Diabetic ketoacidosis or starvation. 5. Renal failure due to any cause etc. DECREASED:- (A),DUE TO DIETRAY DEFICIENCY 1. Dietary deficiency of Zinc, Iron and molybdenum. 2. Fanconi syndrome & Wilsons disease. 3. Multiple scierosis. 4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc. (B).DUE TO INCREASED EXCREATION 1. Drugs:-Probenecid , sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estro	JRIC ACID SERUM				3 60 - 7 70		
(B). DUE TO INCREASED EXCREATION 1. Drugs: - Probenecid, sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estro	NCREASED:- (A).DUE TO INCREASE 1.Idiopathic primary 2.Excessive dietary p 3.Cytolytic treatmen 4.Polycythemai vera 5.Psoriasis. 5.Sickle cell anaemia (B).DUE TO DECREASI 1.Alcohol ingestion. 2.Thiazide diuretics. 3.Lactic acidosis. 4.Aspirin ingestion (I 5.Diabetic ketoacido 6.Renal failure due to DECREASED:- (A).DUE TO DIETARY I 1.Dietary deficiency (2.Fanconi syndrome 3.Multiple sclerosis	D PRODUCTION:- gout. urines (organ meats,legumes,an t of malignancies especially leuk & myeloid metaplasia. etc. ED EXCREATION (BY KIDNEYS) ess than 2 grams per day). sis or starvation. o any cause etc. DEFICIENCY of Zinc, Iron and molybdenum. & Wilsons disease.	emais & lymphomas				
*** End Of Report ***			s (more than 4 grams	per day), corticosterroi	ids and ACTH, anti-coagulants and estrogens e		
			*** End Of Rep	ort ***			





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

V DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

 KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana

 KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana

 0171-2643898, +91 99910 43898
 care@koshealthcare.com
 www.koshealthcare.com



Page 2 of 2