

KOS Diagnostic Lab (A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE PATIENT ID : 1615643

COLLECTED BY : REG. NO./LAB NO. : 012409170006

 REFERRED BY
 : 17/Sep/2024 08:35 AM

 BARCODE NO.
 : 01517107
 COLLECTION DATE
 : 17/Sep/2024 08:37 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 17/Sep/2024 09:13 AM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

SWASTHYA WELLNESS PANEL: GT COMPLETE BLOOD COUNT (CBC)

RED BLOOD CELLS (RBCS) COUNT AND INDICES

| HAEMOGLOBIN (HB) by CALORIMETRIC | 11.4 ^L | gm/dL | 12.0 - 16.0 |
|--|-------------------|--------------|---|
| RED BLOOD CELL (RBC) COUNT by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE | 4.74 | Millions/cmm | 3.50 - 5.50 |
| PACKED CELL VOLUME (PCV) by calculated by automated hematology analyzer | 36.9 | % | 35.0 - 49.0 |
| MEAN CORPUSCULAR VOLUME (MCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER | 77.9 ^L | fL | 80.0 - 100.0 |
| MEAN CORPUSCULAR HAEMOGLOBIN (MCH) by calculated by automated hematology analyzer | 24.1 ^L | pg | 27.0 - 34.0 |
| MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) by calculated by automated hematology analyzer | 30.9 ^L | g/dL | 32.0 - 36.0 |
| RED CELL DISTRIBUTION WIDTH (RDW-CV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER | 14.3 | % | 11.00 - 16.00 |
| RED CELL DISTRIBUTION WIDTH (RDW-SD) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER | 41.7 | fL | 35.0 - 56.0 |
| MENTZERS INDEX by CALCULATED | 16.43 | RATIO | BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0 |
| GREEN & KING INDEX by CALCULATED | 23.55 | RATIO | BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0 |
| WHITE BLOOD CELLS (WBCS) | | | |
| TOTAL LEUCOCYTE COUNT (TLC) by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY | 8110 | /cmm | 4000 - 12000 |
| NUCLEATED RED BLOOD CELLS (NRBCS) by automated 6 part hematology analyzer | NIL | | 0.00 - 20.00 |
| NUCLEATED RED BLOOD CELLS (nRBCS) % by calculated by automated hematology analyzer DIFFERENTIAL LEUCOCYTE COUNT (DLC) | NIL | % | < 10 % |
| NEUTROPHILS by flow cytometry by sf cube & microscopy | 41 ^L | % | 50 - 70 |



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY : REG. NO./LAB NO. : 012409170006

 REFERRED BY
 : 17/Sep/2024 08:35 AM

 BARCODE NO.
 : 01517107
 COLLECTION DATE
 : 17/Sep/2024 08:37 AM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 17/Sep/2024 09:13AM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

| Test Name | Value | Unit | Biological Reference interval |
|--|---------------------|------|-------------------------------|
| LYMPHOCYTES by Flow cytometry by SF cube & microscopy | 52 ^H | % | 20 - 45 |
| EOSINOPHILS by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY | 2 | % | 1 - 6 |
| MONOCYTES by Flow cytometry by SF cube & microscopy | 5 | % | 3 - 12 |
| BASOPHILS by flow cytometry by sf cube & microscopy ABSOLUTE LEUKOCYTES (WBC) COUNT | 0 | % | 0 - 1 |
| ABSOLUTE NEUTROPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY | 3325 | /cmm | 2000 - 7500 |
| ABSOLUTE LYMPHOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY | 4217 | /cmm | 800 - 4900 |
| ABSOLUTE EOSINOPHIL COUNT by flow cytometry by sf cube & microscopy | 162 | /cmm | 40 - 440 |
| ABSOLUTE MONOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY | 406 | /cmm | 80 - 880 |
| ABSOLUTE BASOPHIL COUNT by flow cytometry by sf cube & microscopy PLATELETS AND OTHER PLATELET PREDICTIVE MARKEI | 0 RS . | /cmm | 0 - 110 |
| PLATELET COUNT (PLT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE | 374000 | /cmm | 150000 - 450000 |
| PLATELETCRIT (PCT) by hydro dynamic focusing, electrical impedence | 0.46 ^H | % | 0.10 - 0.36 |
| MEAN PLATELET VOLUME (MPV) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE | 12 ^H | fL | 6.50 - 12.0 |
| PLATELET LARGE CELL COUNT (P-LCC) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE | 154000 ^H | /cmm | 30000 - 90000 |
| PLATELET LARGE CELL RATIO (P-LCR) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE | 41.3 | % | 11.0 - 45.0 |
| PLATELET DISTRIBUTION WIDTH (PDW) by hydro dynamic focusing, electrical impedence NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD | 16.1 | % | 15.0 - 17.0 |



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)









Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE PATIENT ID : 1615643

COLLECTED BY : REG. NO./LAB NO. : 012409170006

 REFERRED BY
 : 17/Sep/2024 08:35 AM

 BARCODE NO.
 : 01517107
 COLLECTION DATE
 : 17/Sep/2024 08:37 AM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 17/Sep/2024 01:40PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

GLYCOSYLATED HAEMOGLOBIN (HBA1C)

GLYCOSYLATED HAEMOGLOBIN (HbA1c): 5.3 % 4.0 - 6.4

WHOLE BLOOD

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

ESTIMATED AVERAGE PLASMA GLUCOSE 105.41 mg/dL 60.00 - 140.00

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

INTERPRETATION:

| AS PER AMERICAN DI | ABETES ASSOCIATION (ADA): | | |
|--|---------------------------|-------------------|--|
| REFERENCE GROUP | GLYCOSYLATED HEMOGL | OGIB (HBAIC) in % | |
| Non diabetic Adults >= 18 years <5.7 | | | |
| At Risk (Prediabetes) | 5.7 – 6.4 | | |
| Diagnosing Diabetes | >= 6.5 | | |
| | Age > 19 Years | | |
| | Goals of Therapy: | < 7.0 | |
| Therapeutic goals for glycemic control | Actions Suggested: >8.0 | | |
| | Age < 19 Y | ears | |
| | Goal of therapy: | <7.5 | |

COMMENTS:

- 1.Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients.

 2.Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high
- concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.

 3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be
- 4.High HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications 5.Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.
- 6.HbA1c results from patients with HbSS,HbSC and HbD must be interpreted with caution, given the pathological processes including anemia,increased red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.
- 7. Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 17/Sep/2024 09:50AM

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY REG. NO./LAB NO. :012409170006

REFERRED BY **REGISTRATION DATE** : 17/Sep/2024 08:35 AM BARCODE NO. :01517107 **COLLECTION DATE** : 17/Sep/2024 08:37AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

ERYTHROCYTE SEDIMENTATION RATE (ESR)

ERYTHROCYTE SEDIMENTATION RATE (ESR)

0 - 20

by MODIFIED WESTERGREN AUTOMATED METHOD

INTERPRETATION:

- 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
- 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
- 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus
 CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

- ESR and C reactive protein (C-RP) are both markers of inflammation.
 Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
 CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
 If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.

- 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
- 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY :012409170006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 17/Sep/2024 08:35 AM BARCODE NO. :01517107 **COLLECTION DATE** : 17/Sep/2024 08:37AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 17/Sep/2024 10:39AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F)

99.47 GLUCOSE FASTING (F): PLASMA mg/dL NORMAL: < 100.0

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0

INTERPRETATION
IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY : REG. NO./LAB NO. : 012409170006

 REFERRED BY
 : 17/Sep/2024 08:35 AM

 BARCODE NO.
 : 01517107
 COLLECTION DATE
 : 17/Sep/2024 08:37 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 17/Sep/2024 10:39 AM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

| Test Name | Value | Unit | Biological Reference interval |
|--|---------------|---------|--|
| | | | |
| | LIPID PROFILE | : BASIC | |
| CHOLESTEROL TOTAL: SERUM by CHOLESTEROL OXIDASE PAP | 131.99 | mg/dL | OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0 |
| TRIGLYCERIDES: SERUM by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC) | 104.72 | mg/dL | OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0 |
| HDL CHOLESTEROL (DIRECT): SERUM by SELECTIVE INHIBITION | 42.31 | mg/dL | LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0 |
| LDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY | 68.74 | mg/dL | OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0 |
| NON HDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY | 89.68 | mg/dL | OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0 |
| VLDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY | 20.94 | mg/dL | 0.00 - 45.00 |
| TOTAL LIPIDS: SERUM by CALCULATED, SPECTROPHOTOMETRY | 368.7 | mg/dL | 350.00 - 700.00 |
| CHOLESTEROL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY | 3.12 | RATIO | LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0 |
| LDL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY | 1.62 | RATIO | LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0 |



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY :012409170006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 17/Sep/2024 08:35 AM BARCODE NO. :01517107 **COLLECTION DATE** : 17/Sep/2024 08:37AM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 17/Sep/2024 10:39AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

TRIGLYCERIDES/HDL RATIO: SERUM **RATIO** 3.00 - 5.002.48^L by CALCULATED, SPECTROPHOTOMETRY

INTERPRETATION:

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co-primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement

CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE PATIENT ID : 1615643

COLLECTED BY : REG. NO./LAB NO. : 012409170006

 REFERRED BY
 : 17/Sep/2024 08:35 AM

 BARCODE NO.
 : 01517107
 COLLECTION DATE
 : 17/Sep/2024 08:37 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 17/Sep/2024 10:39 AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

| LIVER FUNCTION TEST (C | COMPLETE |) |
|------------------------|----------|---|
|------------------------|----------|---|

| BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY | 0.29 | mg/dL | INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20 |
|--|-------------------|-------|---|
| BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY | 0.09 | mg/dL | 0.00 - 0.40 |
| BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY | 0.2 | mg/dL | 0.10 - 1.00 |
| SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE | 17.7 | U/L | 7.00 - 45.00 |
| SGPT/ALT: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE | 19.7 | U/L | 0.00 - 49.00 |
| AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY | 0.9 | RATIO | 0.00 - 46.00 |
| ALKALINE PHOSPHATASE: SERUM by Para nitrophenyl phosphatase by amino meth propanol | 196.3 YL | U/L | 50.00 - 370.00 |
| GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY | 10.12 | U/L | 0.00 - 55.0 |
| TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY | 6.39 | gm/dL | 6.20 - 8.00 |
| ALBUMIN: SERUM by BROMOCRESOL GREEN | 4.58 | gm/dL | 3.50 - 5.50 |
| GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY | 1.81 ^L | gm/dL | 2.30 - 3.50 |
| A : G RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY | 2.53 ^H | RATIO | 1.00 - 2.00 |

INTERPRETATION

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Reference Range.

USE:- Differential diagnosis of diseases of hepatobiliary system and pancreas.

INCREASED:

| DRUG HEPATOTOXICITY | > 2 |
|--|----------------------------|
| ALCOHOLIC HEPATITIS | > 2 (Highly Suggestive) |
| CIRRHOSIS | 1.4 - 2.0 |
| INTRAHEPATIC CHOLESTATIS | > 1.5 |
| HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS | > 1.3 (Slightly Increased) |



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY : REG. NO./LAB NO. : 012409170006

 REFERRED BY
 : 17/Sep/2024 08:35 AM

 BARCODE NO.
 : 01517107
 COLLECTION DATE
 : 17/Sep/2024 08:37 AM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 17/Sep/2024 10:39AM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

DECREASED:

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

PROGNOSTIC SIGNIFICANCE:

| NORMAL | < 0.65 |
|----------------------|-----------|
| GOOD PROGNOSTIC SIGN | 0.3 - 0.6 |
| POOR PROGNOSTIC SIGN | 1.2 - 1.6 |



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUĞAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana
KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana
0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY : REG. NO./LAB NO. : 012409170006

 REFERRED BY
 : 17/Sep/2024 08:35 AM

 BARCODE NO.
 : 01517107
 COLLECTION DATE
 : 17/Sep/2024 08:37 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 17/Sep/2024 10:39 AM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

| Test Name | Value | Unit | Biological Reference interval |
|--|-------------------|----------------|-------------------------------|
| | KIDNEY FUNCTION T | EST (COMPLETE) | |
| UREA: SERUM | 13.41 | mg/dL | 10.00 - 50.00 |
| by UREASE - GLUTAMATE DEHYDROGENASE (GLDH) CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY | 0.59 | mg/dL | 0.40 - 1.20 |
| BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETRY | 6.27 ^L | mg/dL | 7.0 - 25.0 |
| BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM | 10.63 | RATIO | 10.0 - 20.0 |
| by CALCULATED, SPECTROPHOTOMETRY UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY | 22.73 | RATIO | |
| URIC ACID: SERUM by URICASE - OXIDASE PEROXIDASE | 3.36 | mg/dL | 2.50 - 6.80 |
| CALCIUM: SERUM by ARSENAZO III, SPECTROPHOTOMETRY | 9.96 | mg/dL | 8.50 - 10.60 |
| PHOSPHOROUS: SERUM by PHOSPHOMOLYBDATE, SPECTROPHOTOMETRY | 4.33 | mg/dL | 2.30 - 4.70 |

ELECTROLYTES SODIUM: SERUM

 SODIUM: SERUM
 141
 mmol/L
 135.0 - 150.0

 by ISE (ION SELECTIVE ELECTRODE)
 4.4
 mmol/L
 3.50 - 5.00

 by ISE (ION SELECTIVE ELECTRODE)
 105.75
 mmol/L
 90.0 - 110.0

 by ISE (ION SELECTIVE ELECTRODE)
 105.75
 mmol/L
 90.0 - 110.0

ESTIMATED GLOMERULAR FILTERATION RATE

ESTIMATED GLOMERULAR FILTERATION RATE 139.8

(eGFR): SERUM by CALCULATED INTERPRETATION:

To differentiate between pre- and post renal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

- 1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.
- 2. Catabolic states with increased tissue breakdown.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 17/Sep/2024 10:39AM

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY :012409170006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 17/Sep/2024 08:35 AM BARCODE NO. :01517107 **COLLECTION DATE** : 17/Sep/2024 08:37AM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

REPORTING DATE

3. GI haemorrhage.

CLIENT CODE.

- 4. High protein intake.
- 5. Impaired renal function plus
- 6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushing's syndrome, high protein diet, burns, surgery, cachexia, high fever).
- 7. Urine reabsorption (e.g. ureter colostomy)
- 8. Reduced muscle mass (subnormal creatinine production)
- 9. Certain drugs (e.g. tetracycline, glucocorticoids)

INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2. Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN:

- 1. Acute tubular necrosis.
- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.
- 5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
- 8. Pregnancy.

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure.

INAPPROPIATE RATIO:

- 1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio)
- 2. Cephalosporin therapy (interferes with creatinine measurement). ESTIMATED GLOMERULAR FILTERATION RATE:

| ESTIMINATED GEOMERODIAN TIETERATION INVIE: | | | | | |
|--|--|-----------------------|---|--|--|
| CKD STAGE | DESCRIPTION | GFR (mL/min/1.73m2) | ASSOCIATED FINDINGS | | |
| G1 | Normal kidney function | >90 | No proteinuria | | |
| G2 | Kidney damage with normal or high GFR | >90 | Presence of Protein , Albumin or cast in urine | | |
| G3a | Mild decrease in GFR | 60 -89 | | | |
| G3b | Moderate decrease in GFR | 30-59 | | | |
| G4 | Severe decrease in GFR | 15-29 | | | |
| G5 | Kidney failure | <15 | | | |



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY REG. NO./LAB NO. :012409170006

REFERRED BY **REGISTRATION DATE** : 17/Sep/2024 08:35 AM BARCODE NO. :01517107 **COLLECTION DATE** : 17/Sep/2024 08:37AM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 17/Sep/2024 10:39AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

COMMENTS:

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.

2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 17/Sep/2024 10:39AM

: Miss. PIHU **NAME**

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY :012409170006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 17/Sep/2024 08:35 AM BARCODE NO. :01517107 **COLLECTION DATE** : 17/Sep/2024 08:37AM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

ENDOCRINOLOGY

REPORTING DATE

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 1.214 ng/mL 0.35 - 1.93by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROXINE (T4): SERUM 7.47 4.87 - 13.20 μgm/dL

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROID STIMULATING HORMONE (TSH): SERUM 1.903 μIU/mL 0.50 - 5.50

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

CLIENT CODE.

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and trilodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

| CLINICAL CONDITION | Т3 | T4 | TSH |
|------------------------------|-----------------------|-----------------------|---------------------------------|
| Primary Hypothyroidism: | Reduced | Reduced | Increased (Significantly) |
| Subclinical Hypothyroidism: | Normal or Low Normal | Normal or Low Normal | High |
| Primary Hyperthyroidism: | Increased | Increased | Reduced (at times undetectable) |
| Subclinical Hyperthyroidism: | Normal or High Normal | Normal or High Normal | Reduced |

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs
- 3. Serum T4 levies in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

| TRIIODOTHYRONINE (T3) | | THYROXI | NE (T4) | THYROID STIMULATING HORMONE (TS | |
|-----------------------|-----------------------------|-------------------|-----------------------------|---------------------------------|------------------------------|
| Age | Refferance Range (ng/mL) | Age | Refferance Range (μg/dL) | Age | Reference Range (μΙυ/mL) |
| 0 - 7 Days | 0.20 - 2.65 | 0 - 7 Days | 5.90 - 18.58 | 0 - 7 Days | 2.43 - 24.3 |
| 7 Days - 3 Months | 0.36 - 2.59 | 7 Days - 3 Months | 6.39 - 17.66 | 7 Days - 3 Months | 0.58 - 11.00 |
| 3 - 6 Months | 0.51 - 2.52 | 3 - 6 Months | 6.75 – 17.04 | 3 Days – 6 Months | 0.70 - 8.40 |



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Miss. PIHU

AGE/ GENDER : 11 YRS/FEMALE **PATIENT ID** : 1615643

COLLECTED BY : REG. NO./LAB NO. : 012409170006

 REFERRED BY
 : 17/Sep/2024 08:35 AM

 BARCODE NO.
 : 01517107
 COLLECTION DATE
 : 17/Sep/2024 08:37 AM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 17/Sep/2024 10:39AM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

| Test Name | | | Value | Unit | | Biological Reference interval |
|---------------------|-------------|----------------------|------------------|---------------------|-------------|-------------------------------|
| 6 - 12 Months | 0.74 - 2.40 | 6 - 12 Months | 7.10 – 16.16 | 6 – 12 Months | 0.70 - 7.00 | |
| 1 - 10 Years | 0.92 - 2.28 | 1 - 10 Years | 6.00 - 13.80 | 1 – 10 Years | 0.60 - 5.50 | |
| 11- 19 Years | 0.35 - 1.93 | 11 - 19 Years | 4.87- 13.20 | 11 – 19 Years | 0.50 - 5.50 | |
| > 20 years (Adults) | 0.35 - 1.93 | > 20 Years (Adults) | 4.87 - 12.60 | > 20 Years (Adults) | 0.35- 5.50 | |
| | RECOM | IMENDATIONS OF TSH L | EVELS DURING PRE | GNANCY (µIU/mL) | | |
| 1st Trimester | | | 0.10 - 2.50 | | | |
| 2nd Trimester | | | 0.20 - 3.00 | | | |
| 3rd Trimester | | | 0.30 - 4.10 | | | |

INCREASED TSH LEVELS:

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

- 1.Toxic multi-nodular goitre & Thyroiditis.
- $2. Over \ replacement \ of \ thyroid \ harmone \ in \ treatment \ of \ hypothyroid ism.$
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituatary or hypothalmic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester

*** End Of Report **



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana