

**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 03:05PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## HAEMATOLOGY

### HAEMOGLOBIN (HB)

HAEMOGLOBIN (HB)	12.2	gm/dL	12.0 - 16.0
by CALORIMETRIC			

#### INTERPRETATION:-

Hemoglobin is the protein molecule in red blood cells that carries oxygen from the lungs to the body's tissues and returns carbon dioxide from the tissues back to the lungs.

A low hemoglobin level is referred to as ANEMIA or low red blood count.

#### ANEMIA ( DECREASED HAEMOGLOBIN):

- 1) Loss of blood (traumatic injury, surgery, bleeding, colon cancer or stomach ulcer)
- 2) Nutritional deficiency (iron, vitamin B12, folate)
- 3) Bone marrow problems (replacement of bone marrow by cancer)
- 4) Suppression by red blood cell synthesis by chemotherapy drugs
- 5) Kidney failure
- 6) Abnormal hemoglobin structure (sickle cell anemia or thalassemia).

#### POLYCYTHEMIA (INCREASED HAEMOGLOBIN):

- 1) People in higher altitudes (Physiological)
- 2) Smoking (Secondary Polycythemia)
- 3) Dehydration produces a falsely rise in hemoglobin due to increased haemoconcentration
- 4) Advanced lung disease (for example, emphysema)
- 5) Certain tumors
- 6) A disorder of the bone marrow known as polycythemia rubra vera,
- 7) Abuse of the drug erythropoietin (Epogen) by athletes for blood doping purposes (increasing the amount of oxygen available to the body by chemically raising the production of red blood cells).

**NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD**



  
 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 03:08PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		


Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------


### BLOOD GROUP (ABO) AND RH FACTOR TYPING

ABO GROUP  
 by SLIDE AGGLUTINATION  
 RH FACTOR TYPE  
 by SLIDE AGGLUTINATION

B  
 POSITIVE



  
**DR.VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
**DR.YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 05:07PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

<b>BLEEDING TIME (BT)</b>			
BLEEDING TIME (BT) by DUKE METHOD	1 MIN 45 SEC	MINS	1 - 5



  
**DR.VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
**DR.YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra  
MD (Pathology & Microbiology)  
Chairman & Consultant Pathologist

Dr. Yugam Chopra  
MD (Pathology)  
CEO & Consultant Pathologist

**NAME** : Mrs. NISHA  
**AGE/ GENDER** : 34 YRS/FEMALE  
**COLLECTED BY** :  
**REFERRED BY** : LOOMBA HOSPITAL (AMBALA CANTT)  
**BARCODE NO.** : 01517560  
**CLIENT CODE.** : KOS DIAGNOSTIC LAB  
**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**PATIENT ID** : 1622519  
**REG. NO./LAB NO.** : 012409230051  
**REGISTRATION DATE** : 23/Sep/2024 02:34 PM  
**COLLECTION DATE** : 23/Sep/2024 02:57PM  
**REPORTING DATE** : 23/Sep/2024 05:07PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

**CLOTTING TIME (CT)**

CLOTTING TIME (CT) by CAPILLARY TUBE METHOD	5 MIN 20 SEC	MINS	4 - 9
--	--------------	------	-------



DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 04:00PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

### CLINICAL CHEMISTRY/BIOCHEMISTRY

#### GLUCOSE RANDOM (R)


GLUCOSE RANDOM (R): PLASMA	91.77	mg/dL	NORMAL: < 140.00
by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)			PREDIABETIC: 140.0 - 200.0
			DIABETIC: > OR = 200.0


#### INTERPRETATION

##### IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A random plasma glucose level below 140 mg/dl is considered normal.
2. A random glucose level between 140 - 200 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
3. A random glucose level of above 200 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



  
**DR. VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
**DR. YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)





**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 04:00PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## ENDOCRINOLOGY

### THYROID STIMULATING HORMONE (TSH)

THYROID STIMULATING HORMONE (TSH): SERUM 1.768  $\mu$ IU/mL 0.35 - 5.50

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

#### INTERPRETATION:

AGE	REFERENCE RANGE ( $\mu$ IU/mL)
0 – 5 DAYS	0.70 – 15.20
6 Days – 2 Months	0.70 – 11.00
3 – 11 Months	0.70 – 8.40
1 – 5 Years	0.70 – 7.00
6 – 10 Years	0.60 – 5.50
11 - 15	0.50 – 5.50
> 20 Years (Adults)	0.27 – 5.50
<b>PREGNANCY</b>	
1st Trimester	0.10 - 3.00
2nd Trimester	0.20 - 3.00
3rd Trimester	0.30 - 4.10

**NOTE:- TSH levels are subjected to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50 %. Hence time of the day has influence on the measured serum TSH concentration.**

**USE:-** TSH controls biosynthesis and release of thyroid hormones T4 & T3. It is a sensitive measure of thyroid function, especially useful in early or subclinical hypothyroidism, before the patient develops any clinical findings or goitre or any other thyroid function abnormality.

#### INCREASED LEVELS:

- 1.Primary or untreated hypothyroidism, may vary from 3 times to more than 100 times normal depending on degree of hypofunction.
- 2.Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3.Hashimotos thyroiditis.
- 4.DRUGS: Amphetamines, Iodine containing agents and dopamine antagonist.
- 5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge.

#### DECREASED LEVELS:

- 1.Toxic multi-nodular goitre & Thyroiditis.
- 2.Over replacement of thyroid hormone in treatment of hypothyroidism.
- 3.Autonomously functioning Thyroid adenoma
- 4.Secondary pituitary or hypothalamic hypothyroidism
- 5.Acute psychiatric illness
- 6.Severe dehydration.





**DR.VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)



**DR.YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 04:00PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8.Pregnancy: 1st and 2nd Trimester

**LIMITATIONS:**

- 1.TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.
- 2.Autoimmune disorders may produce spurious results.



  
**DR.VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
**DR.YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 05:05PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

### PROLACTIN

#### PROLACTIN: SERUM

129.47<sup>H</sup>

ng/mL

3 - 25

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

#### INTERPRETATION:

1. Prolactin is secreted by the anterior pituitary gland and controlled by the hypothalamus.  
 2. The major chemical controlling prolactin secretion is dopamine, which inhibits prolactin secretion from the pituitary.  
 3. Physiological function of prolactin is the stimulation of milk production. In normal individuals, the prolactin level rises in response to physiologic stimuli such as sleep, exercise, nipple stimulation, sexual intercourse, hypoglycemia, postpartum period, and also is elevated in the newborn infant.

#### INCREASED (HYPERPROLACTEMIA):

1. Prolactin-secreting pituitary adenoma (prolactinoma, which is 5 times more frequent in females than males).  
 2. Functional and organic disease of the hypothalamus.  
 3. Primary hypothyroidism.  
 4. Section compression of the pituitary stalk.  
 5. Chest wall lesions and renal failure.  
 6. Ectopic tumors.  
 7. DRUGS:- Anti-Dopaminergic drugs like antipsychotic drugs, antinausea/antiemetic drugs, Drugs that affect CNS serotonin metabolism, serotonin receptors, or serotonin reuptake (anti-depressants of all classes, ergot derivatives, some illegal drugs such as cannabis), Antihypertensive drugs, Opiates, High doses of estrogen or progesterone, anticonvulsants (valproic acid), anti-tuberculous medications (Isoniazid).

#### SIGNIFICANCE:

1. In loss of libido, galactorrhea, oligomenorrhea or amenorrhea, and infertility in premenopausal females.  
 2. Loss of libido, impotence, infertility, and hypogonadism in males. Postmenopausal and premenopausal women, as well as men, can also suffer from decreased muscle mass and osteoporosis.  
 3. In males, prolactin levels >13 ng/mL are indicative of hyperprolactinemia.  
 4. In women, prolactin levels >27 ng/mL in the absence of pregnancy and postpartum lactation are indicative of hyperprolactinemia.  
 5. Clear symptoms and signs of hyperprolactinemia are often absent in patients with serum prolactin levels <100 ng/mL.  
 4. Mild to moderately increased levels of serum prolactin are not a reliable guide for determining whether a prolactin-producing pituitary adenoma is present, 5. Whereas levels >250 ng/mL are usually associated with a prolactin-secreting tumor.

#### CAUTION:

Prolactin values that exceed the reference values may be due to macroprolactin (prolactin bound to immunoglobulin). Macroprolactin should be evaluated if signs and symptoms of hyperprolactinemia are absent, or pituitary imaging studies are not informative.





**DR. VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)



**DR. YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)





**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 04:00PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

### ANTI MULLERIAN HORMONE (AMH) GEN II

ANTI MULLERIAN HORMONE (AMH) GEN II: SERUM 0.733 ng/mL 0.05 - 11.00  
 by ECLIA (ELECTROCHEMILUMINESCENCE IMMUNOASSAY)

#### INTERPRETATION:-

A Correlation of FERTILITY POTENTIAL and AMH levels are :

OVARIAN FERTILITY POTENTIAL	AMH VALUES IN (ng/mL)
OPTIMAL FERTILITY:	4.00 – 6.80 ng/mL
SATISFACTORY FERTILITY:	2.20 – 4.00 ng/mL
LOW FERTILITY:	0.30 – 2.20 ng/mL
VERY LOW/UNDETECTABLE:	0.00 – 0.30 ng/mL
HIGH LEVEL:	>6.8 ng/mL (PCOD/GRANULOSA CELL TUMOUR)

Anti Mullerian Hormone (AMH) is also known as Mullerian Inhibiting Substance provided by sertoli cells of the testis in males and by ovarian granulosa cells in females upto antral stage in females.

#### IN MALES:

1.It is used to evaluate testicular presence and function in infants with intersex conditions or ambiguous genitalia, and to distinguish between cryptorchidism and anorchia in males

#### IN FEMALES:

- During reproductive age, follicular AMH production begins during the primary stage, peaks in preantral stage & has influence on follicular sensitivity to FSH which is important in selection for follicular dominance. AMH levels thus represents the pool or number of primordial follicles but not the quality of oocytes. AMH does not vary significantly during menstrual cycle & hence can be measured independently of day of cycle.
- Polycystic ovarian syndrome can elevate AMH 2 to 5 fold higher than age specific reference range & predict anovulatory, irregular cycles, ovarian tumours like Granulosa cell tumour are often associated with higher AMH levels.
- Obese women are often associated with diminished ovarian reserve and can have 65% lower mean AMH levels than non-obese women.
- In females, AMH levels do not change significantly throughout the menstrual cycle and decrease with age.
- Assess Ovarian Reserve - correlates with the number of antral follicles in the ovaries.
- Evaluate fertility potential and ovarian response in IVF - Women with low AMH levels are more likely to be poor ovarian responders.
- Assess the condition of Polycystic Ovary and premature ovarian failure.

A combination of Age, Ultrasound markers-Ovarian Volume and Antral Follicle Count, AMH and FSH levels are useful for optimal assessment of ovarian reserve. Studies in various fertility clinics are ongoing to establish optimal AMH concentration for predicting response to invitro fertilization, however, given below is suggested interpretative reference.

AMH levels (ng/mL)	Suggested patient	Anticipated Antral	Anticipated FSH levels	Anticipated Response
--------------------	-------------------	--------------------	------------------------	----------------------



  
 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 04:00PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
	Categorization for fertility based on AMH for age group (20 to 45 yrs)	Follicle counts	(day 3) to IVF/COH cycle
Below 0.3	Very low	Below 4	Above 20 Negligible/Poor
0.3 to 2.19	Low	4 - 10	Usually 16 - 20 Reduced
2.19 to 4.00	Satisfactory	11 - 25	Within reference range or between 11 - 15 Safe/Normal
Above 4.00	Optimal	Upto 30 and Above	Within reference range or between 11 – 15 or Above 15 Possibly Excessive

**INCREASED:**

1. Polycystic ovarian syndrome (most common)
2. Ovarian Tumour: Granulosa cell tumour

**DECREASED:**

1. Anorchia , Abnormal or absence of testis in males
2. Pseudohermaphroditism
3. Post Menopause

**NOTE:**

1. AMH measurement alone is seldom sufficient for diagnosis and results should be interpreted in the light of clinical finding and other relevant test such as ovarian ultrasonography (In fertility applications); abdominal or testicular ultrasound (intersex or testicular function applications); measurement of sex steroids (estradiol, Progesterone, Testosterone), FSH, Inhibin B (For fertility), and Inhibin A and B (for tumour work up).
2. Conversion of AMH from ng/mL to pmol/L can be performed by using equation  $1 \text{ ng/mL} = 7.14 \text{ pmol/L}$



  
**DR. VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
**DR. YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 04:19PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

### IMMUNOPATHOLOGY/SEROLOGY

#### HEPATITIS C VIRUS (HCV) ANTIBODY: TOTAL

HEPATITIS C ANTIBODY (HCV) TOTAL: SERUM	0.12	S/CO	NEGATIVE: < 1.00
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			
HEPATITIS C ANTIBODY (HCV) TOTAL	NON - REACTIVE		POSITIVE: > 1.00
RESULT			
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			

#### INTERPRETATION:-

RESULT (INDEX)	REMARKS
< 1.00	NON - REACTIVE/NOT - DETECTED
> =1.00	REACTIVE/ASYMPTOMATIC/INFECTIVE STATE/CARRIER STATE.

Hepatitis C (HCV) is an RNA virus of Favivirus group transmitted via blood transfusions, transplantation, injection drug abusers, accidental needle punctures in healthcare workers, dialysis patients and rarely from mother to infant. 10 % of new cases show sexual transmission. As compared to HAV & HBV , chronic infection with HCV occurs in 85 % of infected individuals. In high risk population, the predictive value of Anti HCV for HCV infection is > 99% whereas in low risk populations it is only 25 %.

#### USES:

- Indicator of past or present infection, but does not differentiate between Acute/ Chronic/Resolved Infection.
- Routine screening of low and high prevalence population including blood donors.

#### NOTE:

- False positive results are seen in Auto-immune disease, Rheumatoid Factor, HYpergammaglobulinemia, Paraproteinemia, Passive antibody transfer, Anti-idiotypes and Anti-superoxide dismutase.
- False negative results are seen in early Acute infection, Immunosuppression and Immuno— incompetence.
- HCV-RNA PCR recommended in all reactive results to differentiate between past and present infection.



  
**DR.VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
**DR.YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 05:05PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

**ANTI HUMAN IMMUNODEFICIENCY VIRUS (HIV) DUO ULTRA WITH (P-24 ANTIGEN DETECTION)**

HIV 1/2 AND P24 ANTIGEN: SERUM	0.13	S/CO	NEGATIVE: < 1.00 POSITIVE: > 1.00
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			
HIV 1/2 AND P24 ANTIGEN RESULT	NON - REACTIVE		
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			

**INTERPRETATION:-**


RESULT (INDEX)	REMARKS
< 1.00	NON - REACTIVE
> = 1.00	PROVISIONALLY REACTIVE

Non-Reactive result implies that antibodies to HIV 1/ 2 have not been detected in the sample . This means that patient has either not been exposed to HIV 1/ 2 infection or the sample has been tested during the "window phase" i.e. before the development of detectable levels of antibodies. Hence a Non Reactive result does not exclude the possibility of exposure or infection with HIV 1/ 2.

**RECOMMENDATIONS:**

- Results to be clinically correlated
- Rarely falsenegativity/positivity may occur.



  
**DR.VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
**DR.YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)





**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 04:00PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

### HEPATITIS B SURFACE ANTIGEN (HBsAg) ULTRA

HEPATITIS B SURFACE ANTIGEN (HBsAg): 0.03 S/CO  
 SERUM  
 NEGATIVE: < 1.0  
 POSITIVE: > 1.0

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

HEPATITIS B SURFACE ANTIGEN (HBsAg) NON REACTIVE  
 RESULT

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

#### INTERPRETATION:

RESULT IN INDEX VALUE	REMARKS
< 1.30	NEGATIVE (-ve)
>=1.30	POSITIVE (+ve)

Hepatitis B Virus (HBV) is a member of the Hepadna virus family causing infection of the liver with extremely variable clinical features. Hepatitis B is transmitted primarily by body fluids especially serum and also spread effectively sexually and from mother to baby. In most individuals HBV hepatitis is self limiting, but 1-2 % normal adolescent and adults develop Chronic Hepatitis. Frequency of chronic HBV infection is 5-10% in immunocompromised patients and 80 % neonates. The initial serological marker of acute infection is HBsAg which typically appears 2-3 months after infection and disappears 12-20 weeks after onset of symptoms. Persistence of HBsAg for more than 6 months indicates carrier state or Chronic Liver disease.



  
**DR. VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
**DR. YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

Dr. Yugam Chopra  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. NISHA	<b>PATIENT ID</b>	: 1622519
<b>AGE/ GENDER</b>	: 34 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012409230051
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 23/Sep/2024 02:34 PM
<b>REFERRED BY</b>	: LOOMBA HOSPITAL (AMBALA CANTT)	<b>COLLECTION DATE</b>	: 23/Sep/2024 02:57PM
<b>BARCODE NO.</b>	: 01517560	<b>REPORTING DATE</b>	: 23/Sep/2024 03:14PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
VDRL by IMMUNOCHROMATOGRAPHY	NON REACTIVE		NON REACTIVE

**INTERPRETATION:**

- Does not become positive until 7 - 10 days after appearance of chancre.
- High titer (>1:16) - active disease.**
- Low titer (<1:8) - biological falsepositive test in 90% cases or due to late or late latent syphilis.**
- Treatment of primary syphilis causes progressive decline to negative VDRL within 2 years.
- Rising titer (4X) indicates relapse, reinfection, or treatment failure and need for retreatment.
- May be nonreactive in early primary, late latent, and late syphilis (approx. 25% of cases).
- Reactive and weakly reactive tests should always be confirmed with FTA-ABS (fluorescent treponemal antibody absorption test).**

**SHORT TERM FALSE POSITIVE TEST RESULTS (<6 MONTHS DURATION) MAY OCCUR IN:**

- Acute viral illnesses (e.g., hepatitis, measles, infectious mononucleosis)
- M. pneumoniae; Chlamydia; Malaria infection.
- Some immunizations
- Pregnancy (rare)

**LONG TERM FALSE POSITIVE TEST RESULTS (>6 MONTHS DURATION) MAY OCCUR IN:**

- Serious underlying disease e.g., collagen vascular diseases, leprosy, malignancy.
- Intravenous drug users.
- Rheumatoid arthritis, thyroiditis, AIDS, Sjogren's syndrome.
- <10 % of patients older than age 70 years.
- Patients taking some anti-hypertensive drugs.

\*\*\* End Of Report \*\*\*



  
 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)

