

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mrs. SHEELA
AGE/ GENDER : 40 YRS/FEMALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 01517685
CLIENT CODE. : KOS DIAGNOSTIC LAB
CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

PATIENT ID : 1624753
REG. NO./LAB NO. : 012409250037
REGISTRATION DATE : 25/Sep/2024 11:48 AM
COLLECTION DATE : 25/Sep/2024 11:49AM
REPORTING DATE : 25/Sep/2024 01:51PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

TUMOUR MARKER

CARCINO EMBRYONIC ANTIGEN (CEA)

CARCINO EMBRYONIC ANTIGEN (CEA): SERUM by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)	0.74	ng/mL	< 5.0
---	------	-------	-------

INTERPRETATION:

1. Carcinoembryonic antigen (CEA) is a glycoprotein normally found in embryonic endodermal epithelium.
2. Increased levels may be found in patients with primary colorectal cancer or other malignancies including medullary thyroid carcinoma and breast, gastrointestinal tract, liver, lung, ovarian, pancreatic, and prostatic cancers.
3. Serial monitoring of CEA should begin prior to initiation of cancer therapy to verify post therapy decrease in concentration and to establish a baseline for evaluating possible recurrence. Levels generally return to normal within 1 to 4 months after removal of cancerous tissue.

CLINICAL SIGNIFICANCE:

1. Monitoring colorectal cancer and selected other cancers such as medullary thyroid carcinoma
2. May be useful in assessing the effectiveness of chemotherapy or radiation treatment.

NOTE:

1. Carcinoembryonic antigen levels should not be used for screening of the general population for undetected cancers.
2. Grossly elevated carcino-embryonic antigen (CEA) concentrations (>20 ng/mL) in a patient with compatible symptoms are strongly suggestive of the presence of cancer and also suggest metastasis.
3. Most healthy subjects (97%) have values < or =3.0 ng/mL.
4. After removal of a colorectal tumor, the serum CEA concentration should return to normal by 6 weeks, unless there is residual tumor.
5. Increases in test values over time in a patient with a history of cancer suggest tumor recurrence.

*** End Of Report ***



DR. VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR. YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)

