Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. JASVIR KAUR**		
AGE/ GENDER	: 40 YRS/FEMALE	**PATIENT ID**	: 1627654
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012409270066**
REFERRED BY	:	**REGISTRATION DATE**	: 27/Sep/2024 07:16 PM
BARCODE NO.	: 01517842	**COLLECTION DATE**	: 27/Sep/2024 07:17PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Sep/2024 07:26PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

HAEMATOLOGY

COMPLETE BLOOD COUNT (CBC)

Test Name	Value	Unit	Biological Reference interval
RED BLOOD CELLS (RBCS) COUNT AND INDICES			
HAEMOGLOBIN (HB) *by CALORIMETRIC*	**11.3[L]**	**gm/dL**	**12.0 - 16.0**
RED BLOOD CELL (RBC) COUNT *by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE*	4.31	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	**35.5[L]**	**%**	**37.0 - 50.0**
MEAN CORPUSCULAR VOLUME (MCV) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	82.2	fL	80.0 - 100.0
MEAN CORPUSCULAR HAEMOGLOBIN (MCH) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	**26.2[L]**	**pg**	**27.0 - 34.0**
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	**31.8[L]**	**g/dL**	**32.0 - 36.0**
RED CELL DISTRIBUTION WIDTH (RDW-CV) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	14.1	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	43.4	fL	35.0 - 56.0
MENTZERS INDEX *by CALCULATED*	19.07	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX *by CALCULATED*	26.87	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CELLS (WBCS)			
TOTAL LEUCOCYTE COUNT (TLC) *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	7320	/cmm	4000 - 11000
NUCLEATED RED BLOOD CELLS (nRBCS) *by AUTOMATED 6 PART HEMATOLOGY ANALYZER*	NIL		0.00 - 20.00
NUCLEATED RED BLOOD CELLS (nRBCS) % *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	NIL	%	< 10 %
DIFFERENTIAL LEUCOCYTE COUNT (DLC)			
NEUTROPHILS *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	**72[H]**	**%**	**50 - 70**

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. JASVIR KAUR**		
AGE/ GENDER	: 40 YRS/FEMALE	**PATIENT ID**	: 1627654
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012409270066**
REFERRED BY	:	**REGISTRATION DATE**	: 27/Sep/2024 07:16 PM
BARCODE NO.	: 01517842	**COLLECTION DATE**	: 27/Sep/2024 07:17PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Sep/2024 07:26PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
LYMPHOCYTES ***by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY***	**15^{L}**	**%**	**20 - 40**
EOSINOPHILS ***by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY***	**8^{H}**	**%**	**1 - 6**
MONOCYTES *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	5	%	2 - 12
BASOPHILS *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	0	%	0 - 1
ABSOLUTE LEUKOCYTES (WBC) COUNT			
ABSOLUTE NEUTROPHIL COUNT *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	5270	/cmm	2000 - 7500
ABSOLUTE LYMPHOCYTE COUNT *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	1098	/cmm	800 - 4900
ABSOLUTE EOSINOPHIL COUNT ***by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY***	**586^{H}**	**/cmm**	**40 - 440**
ABSOLUTE MONOCYTE COUNT *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	366	/cmm	80 - 880
PLATELETS AND OTHER PLATELET PREDICTIVE MARKERS.			
PLATELET COUNT (PLT) *by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE*	175000	/cmm	150000 - 450000
PLATELETCRIT (PCT) *by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE*	0.2	%	0.10 - 0.36
MEAN PLATELET VOLUME (MPV) *by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE*	11	fL	6.50 - 12.0
PLATELET LARGE CELL COUNT (P-LCC) *by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE*	62000	/cmm	30000 - 90000
PLATELET LARGE CELL RATIO (P-LCR) *by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE*	35.2	%	11.0 - 45.0
PLATELET DISTRIBUTION WIDTH (PDW) *by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE*	16.4	%	15.0 - 17.0

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. JASVIR KAUR**		
AGE/ GENDER	: 40 YRS/FEMALE	**PATIENT ID**	: 1627654
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012409270066**
REFERRED BY	:	**REGISTRATION DATE**	: 27/Sep/2024 07:16 PM
BARCODE NO.	: 01517842	**COLLECTION DATE**	: 27/Sep/2024 07:17PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Sep/2024 07:53PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

IMMUNOPATHOLOGY/SEROLOGY

TYPHOID COMBO SCREEN (TYPHOID ANTIGEN, IgG AND IgM): SERUM

TYPHOID ANTIGEN - SERUM *by ICT (IMMUNOCHROMATOGRAPHY)*	NEGATIVE (-ve)		NEGATIVE (-ve)
TYPHI DOT ANTIBODY IgG *by ICT (IMMUNOCHROMATOGRAPHY)*	NEGATIVE (-ve)		NEGATIVE (-ve)
TYPHI DOT ANTIBODY IgM *by ICT (IMMUNOCHROMATOGRAPHY)*	NEGATIVE (-ve)		NEGATIVE (-ve)

INTE*RPRETATION:*

Typhoid fever is a life threatening illness caused by the bacterium Salmonella typhus.The infection is acquired typically by ingestion. On reaching the gut, the bacilli attach themselves to the epithelial cells of the intestinal villi and penetrate the lamina and submucosa. They are then phagocytosed there by polymorphs and mesenteric lymph nodes, where they multiply and, via the thoracic duct, enter the blood stream. A transient bacteremia follows, during which the bacilli are seeded in the liver, gall bladder, spleen, bone marrow, lymph nodes, and kidneys, where further multiplication takes place. Towards the end of the incubation period, there occurs a massive bacteremia from these sites, heralding the onset of the clinical symptoms.

The diagnosis of typhoid consists of isolation of the bacilli and the demonstration of antibodies. The isolation of the bacilli is very time consuming and antibody detection is not very specific. Other tests include the Widal reaction. The advantage of this test is that it takes only 10-20 minutes and requires only a small amount of stool/serum/plasma to perform. It is the easiest and most specific method for detecting S. typhi infection.

RELATIVE SENSTIVITY OF TYPHOID ANTIGEN DETECTION: 98.7%
RELATIVE SPECIFICITY OF TYPHOID ANTIGEN DETECTION: 97.4%

DETECTABLE IgM RESPONSE:

ONSET OF FEVER	PERCENT POSITIVE
4 - 6 DAYS	43.5
6 - 9 DAYS	92.9
> 9 DAYS	99.5

1.This is a solid phase, immunochromatographic ELISA assay that detects specific IgM and IgG Antibodies against the OUTER MEMBRAN PROTEIN(OMP) of theSalmonella species. IgM antibodies appear in the serum 2-3 days post infection and are indicative of a recent infection while the IgG antibodies appear later and are useful for presumptive diagnosis of Enteric fever if the patient presents more than a week after onset of symptoms.

2.This is a useful screening assay for the early detection of Enteric fever and has a high sensitivity. However the test has moderate specificity and false positive results may be obtained in the following situations:

- Antibodies against Salmonella may cross react with other antibodies.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mrs. JASVIR KAUR**		
AGE/ GENDER	: 40 YRS/FEMALE	**PATIENT ID**	: 1627654
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012409270066**
REFERRED BY	:	**REGISTRATION DATE**	: 27/Sep/2024 07:16 PM
BARCODE NO.	: 01517842	**COLLECTION DATE**	: 27/Sep/2024 07:17PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Sep/2024 07:53PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

- Unrelated infections may lead to production of specific Salmonella antibodies if the patient has previously been exposed to Salmonella infection **(ANAMNESTIC RESPONSE)**.

NOTE:-Rapid blood culture performed during 1st week of infection is highly recommended for confirmation of all IgM positive results. In case the patient has presented after the first week of infection, a thorough clinical correlation and confirmatory Widal test must be performed to establish the diagnosis.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. JASVIR KAUR**		
AGE/ GENDER	: 40 YRS/FEMALE	**PATIENT ID**	: 1627654
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012409270066**
REFERRED BY	:	**REGISTRATION DATE**	: 27/Sep/2024 07:16 PM
BARCODE NO.	: 01517842	**COLLECTION DATE**	: 27/Sep/2024 07:17PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Sep/2024 08:51PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

DENGUE FEVER ANTIGEN NS1 - ELISA (QUANTITATIVE)

DENGUE NS1 ANTIGEN QUANTITATIVE *by ELISA (ENZYME LINKED IMMUNOSORBENT ASSAY)*	0.19	INDEX	NEGATIVE: < 0.90 BORDERLINE: 0.90 - 1.10 POSITIVE: >=1.10
DENGUE NS1 ANTIGEN RESULT *by ELISA (ENZYME LINKED IMMUNOSORBENT ASSAY)*	NEGATIVE (-ve)		NEGATIVE (-ve)

INTERPRETATION

DENGUE ANTIGEN NS1		
VALUE	**UNIT**	**RESULT**
< 0.90	INDEX	NEGATIVE (-ve)
0.90 - 1.10	INDEX	BORDERLINE
>=1.10	INDEX	POSITIVE (+ve)

1.The test becomes positive within 0-9 days of exposure to the virus (positive results are obtained within 24 hours of exposure in the overwhelming majority of patients) and generally remains positive till 15 days after exposure. The Dengue NS-1 antigen test is extremely useful in the early diagnosis of the disease thus helping in proper follow up and monitoring of the patients.
2.The IgM antibodies on the other hand take a minimum of 5-10 days in primary infection and 4-5 days in secondary infections to test positive and hence are suitable for the diagnosis of dengue fever only when the fever is approximately one week old.

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)