



	Dr. Vinay Chopr MD (Pathology & Mic Chairman & Consulta	robiology)	Dr. Yugam MD CEO & Consultant	(Pathology)	
NAME	: Mr. NARINDER KUMAR				
AGE/ GENDER	: 65 YRS/MALE	PA	TIENT ID	: 1436763	
COLLECTED BY	:	RI	EG. NO./LAB NO.	: 012409290009	
REFERRED BY	:	RI	EGISTRATION DATE	: 29/Sep/2024 08:09 AM	
BARCODE NO.	:01517921	CO	LLECTION DATE	: 29/Sep/2024 08:11AM	
CLIENT CODE.	: KOS DIAGNOSTIC LAB	RH	EPORTING DATE	: 01/Oct/2024 09:48AM	
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMB	ALA CANTT			
Test Name		Value	Unit	Biological Reference interval	
MICROBIOLOGY CULTURE AEROBIC BACTERIA AND ANTIBIOTIC SENSITIVITY: URINE					
CULTURE AND SUSCEPTIBILITY: URINE					
		29-09-2024			
SPECIMEN SOURCE		URINE			
INCUBATION PERIOD by AUTOMATED BROTH CULTURE		48 HOURS			
CULTURE		STERILE			
		NO AEROBIC 37*C	NO AEROBIC PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF INCUBATION AT 37*C		
AEROBIC SUSCEPTIE	BILITY: URINE				
INTERPRETATION: 1. In urine culture an significant. However	d sensitivity, presence of more than in symptomatic patients , a smaller n	100,000 organis umber of bacter	m per mL in midstream ria (100 to 10000/mL) m	sample of urine is considered clinically any signify infection.	

2. Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out" catheterization or from patients with indwelling catheters.

SUSCEPTIBILITY:

1. A test interpreted as SENSTITIVE implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent

recommended for that type of infection and infecting species, unless otherwise indicated.. 2. A test interpreted as **INTERMEDIATE** implies that the Infection due to the isolate may be appropriately treated in body sites where the drugs are physiologically concentrated or when a high dosage of drug can be used". 3.A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal

dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

CAUTION:

Conditions which can cause a false Negative culture: 1. Patient is on antibiotics. Please repeat culture post therapy.

2. Anaerobic bacterial infection.

- 3. Fastidious aerobic bacteria which are not able to grow on routine culture media.
- 4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.

5. Renal tuberculosis to be confirmed by AFB studies.

*** End Of Report ***



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 care@koshealthcare.com www.koshealthcare.com

