





Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

PATIENT ID AGE/ GENDER : 52 YRS/FEMALE : 1628908

COLLECTED BY : 012409290030 : SURJESH REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 11:13AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

SWASTHYA WELLNESS PANEL: 1.5 COMPLETE BLOOD COUNT (CBC)

RED BLOOD CELLS (RBCS) COUNT AND INDICES

HAEMOGLOBIN (HB) by CALORIMETRIC	12.6	gm/dL	12.0 - 16.0
RED BLOOD CELL (RBC) COUNT by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	4.57	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV)	41.3	%	37.0 - 50.0
by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER MEAN CORPUSCULAR VOLUME (MCV)	90.5	fL	80.0 - 100.0
by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER MEAN CORPUSCULAR HAEMOGLOBIN (MCH) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	27.5	pg	27.0 - 34.0
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	30.4 ^L	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION WIDTH (RDW-CV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	15.4	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	52.1	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED	19.8	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX by CALCULATED	30.42	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CELLS (WBCS)			
TOTAL LEUCOCYTE COUNT (TLC) by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	6820	/cmm	4000 - 11000
NUCLEATED RED BLOOD CELLS (nRBCS) by AUTOMATED 6 PART HEMATOLOGY ANALYZER	NIL		0.00 - 20.00
NUCLEATED RED BLOOD CELLS (nRBCS) % by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	NIL	%	< 10 %
DIFFERENTIAL LEUCOCYTE COUNT (DLC)			
NEUTROPHILS by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	67	%	50 - 70



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY: SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM **BARCODE NO.** : 01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 29/Sep/2024 11:13AM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
YMPHOCYTES by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	27	%	20 - 40
OSINOPHILS by flow cytometry by sf cube & microscopy	3	%	1 - 6
IONOCYTES by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	3	%	2 - 12
ASOPHILS by flow cytometry by sf cube & microscopy BSOLUTE LEUKOCYTES (WBC) COUNT	0	%	0 - 1
BSOLUTE NEUTROPHIL COUNT by flow cytometry by SF cube & microscopy	4569	/cmm	2000 - 7500
BSOLUTE LYMPHOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	1841	/cmm	800 - 4900
BSOLUTE EOSINOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	205	/cmm	40 - 440
BSOLUTE MONOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	205	/cmm	80 - 880
BSOLUTE BASOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY LATELETS AND OTHER PLATELET PREDICTIVE MARKER	0	/cmm	0 - 110
LATELETS AND OTHER PLATELET PREDICTIVE IMARKER LATELET COUNT (PLT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	314000	/cmm	150000 - 450000
LATELETCRIT (PCT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	0.3	%	0.10 - 0.36
IEAN PLATELET VOLUME (MPV) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	10	fL	6.50 - 12.0
LATELET LARGE CELL COUNT (P-LCC) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	76000	/cmm	30000 - 90000
LATELET LARGE CELL RATIO (P-LCR) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	24.1	%	11.0 - 45.0
LATELET DISTRIBUTION WIDTH (PDW) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE OTE: TEST CONDUCTED ON EDTA WHOLE BLOOD	16.1	%	15.0 - 17.0



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: Ilnd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY :012409290030 : SURJESH REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 02:06PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

GLYCOSYLATED HAEMOGLOBIN (HBA1C)

7.5^H **GLYCOSYLATED HAEMOGLOBIN (HbA1c):** 4.0 - 6.4

WHOLE BLOOD

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

ESTIMATED AVERAGE PLASMA GLUCOSE mg/dL 60.00 - 140.00 168.55^H

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

INTERPRETATION:

AS PER AMERICAN DIABETES ASSOCIATION (ADA):				
REFERENCE GROUP GLYCOSYLATED HEMOGLOGIB (HBAIC) in %		OGIB (HBAIC) in %		
Non diabetic Adults >= 18 years	<5.7			
At Risk (Prediabetes)	5.7 – 6.	4		
Diagnosing Diabetes	>= 6.5			
	Age > 19 Years			
	Goals of Therapy:	< 7.0		
Therapeutic goals for glycemic control	Actions Suggested:	>8.0		
	Age < 19 Years			
	Goal of therapy:	<7.5		

COMMENTS:

- 1.Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients. 2. Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.
- 3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be
- 4. High HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications 5. Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.
- $6. Hb A1 c \ results \ from \ patients \ with \ Hb SS, Hb SC \ and \ Hb D \ must \ be \ interpreted \ with \ caution \ , given \ the \ pathological \ processes \ including \ anemia, increased$ red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.
- 7. Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. **COLLECTION DATE** : 29/Sep/2024 04:29PM :01517942

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Value Unit **Biological Reference interval** Test Name

ERYTHROCYTE SEDIMENTATION RATE (ESR)

REPORTING DATE

ERYTHROCYTE SEDIMENTATION RATE (ESR)

mm/1st hr

0 - 20

: 29/Sep/2024 11:34AM

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

INTERPRETATION:

CLIENT CODE.

- 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-
- immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.

 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
- 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

NOTE:

- 1. ESR and C reactive protein (C-RP) are both markers of inflammation.
 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while services and quiping may decrease it. aspirin, cortisone, and quinine may decrease it



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH :012409290030 REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 12:04PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

CLINICAL CHEMISTRY/BIOCHEMISTRY **GLUCOSE FASTING (F)**

GLUCOSE FASTING (F): PLASMA 137.61^H mg/dL NORMAL: < 100.0

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0

INTERPRETATION
IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY: SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) REGISTRATION DATE : 29/Sep/2024 10:34 AM BARCODE NO. : 01517942 COLLECTION DATE : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 29/Sep/2024 12:04PM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
	LIPID PROFILE	BASIC	
CHOLESTEROL TOTAL: SERUM by CHOLESTEROL OXIDASE PAP	178.19	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SERUM by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)	190.85 ^H	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (DIRECT): SERUM by SELECTIVE INHIBITION	52.9	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	87.12	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	125.29	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	38.17	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SERUM by CALCULATED, SPECTROPHOTOMETRY	547.23	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	3.37	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0
LDL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	1.65	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH :012409290030 REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 12:04PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
TRIGLYCERIDES/HDL RATIO: SERUM	3.61	RATIO	3.00 - 5.00
by CALCULATED, SPECTROPHOTOMETRY			

INTERPRETATION:

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement

CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana

0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) REGISTRATION DATE : 29/Sep/2024 10:34 AM

BARCODE NO. : 01517942 COLLECTION DATE : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 12:17PM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

LIVER FUNCTION TEST (COMPLETE)

BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY	0.63	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY	0.18	mg/dL	0.00 - 0.40
BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY	0.45	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	34.7	U/L	7.00 - 45.00
SGPT/ALT: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	16.5	U/L	0.00 - 49.00
AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.1	RATIO	0.00 - 46.00
ALKALINE PHOSPHATASE: SERUM by Para Nitrophenyl Phosphatase by Amino Metheropanol	100.53 HYL	U/L	40.0 - 130.0
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY	25.91	U/L	0.00 - 55.0
TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY	6.53	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL GREEN	3.64	gm/dL	3.50 - 5.50
GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.89	gm/dL	2.30 - 3.50
A : G RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	1.26	RATIO	1.00 - 2.00

INTERPRETATION

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range.

USE:- Differential diagnosis of diseases of hepatobiliary system and pancreas.

INCREASED:

DRUG HEPATOTOXICITY_	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM **BARCODE NO.** : 01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 29/Sep/2024 12:17PM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS		> 1.3 (Slightly Increased)	

DECREASED:

- 1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)
- 2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

PROGNOSTIC SIGNIFICANCE:

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUĞAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana
KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana
0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com

CLIENT CODE.



KOS Diagnostic Lab

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 29/Sep/2024 03:05PM

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH :012409290030 REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval

REPORTING DATE

KIDNEY FUNCTION	IF21	(COIVIE	LETE)
41 12		r	na/dl

UREA: SERUM	41.12	mg/dL	10.00 - 50.00
by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)			
CREATININE: SERUM	1.49 ^H	mg/dL	0.40 - 1.20
by ENZYMATIC, SPECTROPHOTOMETERY			
BLOOD UREA NITROGEN (BUN): SERUM	19.21	mg/dL	7.0 - 25.0
by CALCULATED, SPECTROPHOTOMETRY			
BLOOD UREA NITROGEN (BUN)/CREATININE	12.89	RATIO	10.0 - 20.0
RATIO: SERUM			
by CALCULATED, SPECTROPHOTOMETRY			
UREA/CREATININE RATIO: SERUM	27.6	RATIO	
by CALCULATED, SPECTROPHOTOMETRY			
URIC ACID: SERUM	5.49	mg/dL	2.50 - 6.80
by URICASE - OXIDASE PEROXIDASE			
CALCIUM: SERUM	9.79	mg/dL	8.50 - 10.60
by ARSENAZO III, SPECTROPHOTOMETRY			
PHOSPHOROUS: SERUM	2.61	mg/dL	2.30 - 4.70
by PHOSPHOMOLYBDATE, SPECTROPHOTOMETRY			
<u>ELECTROLYTES</u>			
SODIUM: SERUM	140.2	mmol/L	135.0 - 150.0
by ISE (ION SELECTIVE ELECTRODE)			
POTASSIUM: SERUM	5.04 ^H	mmol/L	3.50 - 5.00
by ISE (ION SELECTIVE ELECTRODE)	0.01		
CHLORIDE: SERUM	105.15	mmol/L	90.0 - 110.0
by ISE (ION SELECTIVE ELECTRODE)			

ESTIMATED GLOMERULAR FILTERATION RATE

ESTIMATED GLOMERULAR FILTERATION RATE 42

(eGFR): SERUM by CALCULATED

INTERPRETATION:

To differentiate between pre- and post renal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

- 1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.
- 2. Catabolic states with increased tissue breakdown.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 29/Sep/2024 04:29PM

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH :012409290030 REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 03:05PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

:01517942

Test Name Value Unit **Biological Reference interval**

COLLECTION DATE

3. GI haemorrhage.

BARCODE NO.

- 4. High protein intake.
- 5. Impaired renal function plus
- 6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushing's syndrome, high protein diet, burns, surgery, cachexia, high fever).
- 7. Urine reabsorption (e.g. ureter colostomy)
- 8. Reduced muscle mass (subnormal creatinine production)
- 9. Certain drugs (e.g. tetracycline, glucocorticoids)

INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2. Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN:

- 1. Acute tubular necrosis.
- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.
- 5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
- 8. Pregnancy.

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure.

INAPPROPIATE RATIO:

- 1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio)
- 2. Cephalosporin therapy (interferes with creatinine measurement). ESTIMATED GLOMERULAR FILTERATION RATE:

CKD STAGE	DESCRIPTION	GFR (mL/min/1.73m2)	ASSOCIATED FINDINGS
CKD STAGE	DESCRIPTION	GFK (IIIL/ IIIIII/ 1./ 3IIIZ)	ASSOCIATED FINDINGS
G1	Normal kidney function	>90	No proteinuria
G2	Kidney damage with	>90	Presence of Protein,
	normal or high GFR		Albumin or cast in urine
G3a	Mild decrease in GFR	60 -89	
G3b	Moderate decrease in GFR	30-59	
G4	Severe decrease in GFR	15-29	
G5	Kidney failure	<15	



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 03:05PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

COMMENTS:

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.

2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH :012409290030 REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 12:17PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
	IRON PRO	FILE	
IRON: SERUM by FERROZINE, SPECTROPHOTOMETRY	52.7	μg/dL	37.0 - 145.0
UNSATURATED IRON BINDING CAPACITY (UIBC) :SERUM by FERROZINE, SPECTROPHOTOMETERY	257.5	μg/dL	150.0 - 336.0
TOTAL IRON BINDING CAPACITY (TIBC) :SERUM by SPECTROPHOTOMETERY	310.2	μg/dL	230 - 430
%TRANSFERRIN SATURATION: SERUM by CALCULATED, SPECTROPHOTOMETERY (FERENE)	16.99	%	15.0 - 50.0
TRANSFERRIN: SERUM by SPECTROPHOTOMETERY (FERENE)	220.24	mg/dL	200.0 - 350.0

INTERPRETATION:-

VARIABLES	ANEMIA OF CHRONIC DISEASE	IRON DEFICIENCY ANEMIA	THALASSEMIA α/β TRAIT
SERUM IRON:	Normal to Reduced	Reduced	Normal
TOTAL IRON BINDING CAPACITY:	Decreased	Increased	Normal
% TRANSFERRIN SATURATION:	FERRIN SATURATION: Decreased		Normal
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increased

IRON:

- 1. Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia.i.e iron deficiency anemia, zinc deficiency anemia, anemia of chronic disease and thalassemia syndromes.
- 2. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.

 TOTAL IRON BINDING CAPACITY (TIBC):

 1. It is essential to isolate iron deficiency anemia, is severely contra-indicated in Thalassemia.

 TOTAL IRON BINDING CAPACITY (TIBC):

 1. It is essential to isolate iron deficiency anemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.

 TOTAL IRON BINDING CAPACITY (TIBC):

 1. It is essential to isolate iron deficiency anemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.

 TOTAL IRON BINDING CAPACITY (TIBC):

 1. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.

 TOTAL IRON BINDING CAPACITY (TIBC):

 1. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.

1.Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) REGISTRATION DATE : 29/Sep/2024 10:34 AM BARCODE NO. : 01517942 COLLECTION DATE : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 29/Sep/2024 12:22PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 1.048 ng/mL 0.35 - 1.93

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROXINE (T4): SERUM 6.06 μgm/dL 4.87 - 12.60

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROID STIMULATING HORMONE (TSH): SERUM 6.164^{H} µIU/mL 0.35 - 5.50

by CMIA (CHEMILUMINESCENT MICROPARTICLE

IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

<u>INTERPRETATION</u>:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and trilodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction (hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (eg: phenytoin , salicylates).
- 3. Serum T4 levles in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

TRIIODOTHY	TRIIODOTHYRONINE (T3) THYROXINE (T4)		NE (T4)	THYROID STIMULATING HORMONE (TSH)		
Age	Refferance Range (ng/mL)	Age	Refferance Range (μg/dL)	Age	Reference Range (μΙυ/mL)	
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3	
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00	
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40	



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUĞAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 29/Sep/2024 04:29PM

COLLECTION DATE

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE PATIENT ID : 1628908

COLLECTED BY : SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) REGISTRATION DATE : 29/Sep/2024 10:34 AM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 12:22PM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

:01517942

Test Name			Value	Unit		Biolog	ical Reference interval
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00		
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50		
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50		
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35- 5.50		
	RECC	MMENDATIONS OF TSH LI	EVELS DURING PREGNANCY (μΙU/mL)				
	1st Trimester		0.10 - 2.50				
	2nd Trimester		0.20 - 3.00			•	
	3rd Trimester		0.30 - 4.10				

INCREASED TSH LEVELS:

BARCODE NO.

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

- 1.Toxic multi-nodular goitre & Thyroiditis.
- $2. Over \ replacement \ of \ thyroid \ harmone \ in \ treatment \ of \ hypothyroid ism.$
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituatary or hypothalmic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



KOS Diagnostic Lab (A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE PATIENT ID : 1628908

COLLECTED BY : SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) REGISTRATION DATE : 29/Sep/2024 10:34 AM BARCODE NO. : 01517942 COLLECTION DATE : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 30/Sep/2024 06:50AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

IMMUNOPATHOLOGY/SEROLOGY ANTI NUCLEAR ANTIBODY/FACTOR (ANA/ANF)

ANTI NUCLEUR ANTIBODIES (ANA): SERUM 0.22 INDEX VALUE NEGATIVE: < 1.0

by ELISA (ENZYME LINKED IMMUNOASSAY)

BORDERLINE: 1.0 - 1.20

POSITIVE: > 1.20

INTERPRETATION:-

1. For diagnostic purposes, ANA value should be used as an adjuvant to other clinical and laboratory data available.

2.Measurement of antinuclear antibodies (ANAs) in serum is the most commonly performed screening test for patients suspected of having a systemic rheumatic disease, also referred to as connective tissue disease.

3.ANAs occur in patients with a variety of autoimmune diseases, both systemic and organ-specific. They are particularly common in the systemic rheumatic diseases, which include lupus erythematosus (LE), discoid LE, drug-induced LE, mixed connective tissue disease, Sjogren syndrome scleroderma (systemic sclerosis), CREST (calcinosis, Raynaud's phenomenon, esophageal dysmotility, sclerodactyly, telangiectasia) syndrome, polymyositis/dermatomyositis, and rheumatoid arthritis.

NOTE:

1. The diagnosis of a systemic rheumatic disease is based primarily on the presence of compatible clinical signs and symptoms. The results of tests for autoantibodies including ANA and specific autoantibodies are ancillary. Additional diagnostic criteria include consistent histopathology or specific radiographic findings. Although individual systemic rheumatic diseases are relatively uncommon, a great many patients present with clinical findings that are compatible with a systemic rheumatic disease ANA screening may be useful for ruling out the disease

2.Secondary, disease specific auto antibodies maybe ordered for patients who are screen positive as ancillary aids for the diagnosis of specific auto-immune disorders.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 29/Sep/2024 12:04PM

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY :012409290030 : SURJESH REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

RHEUMATOID FACTOR (RA): QUANTITATIVE - SERUM

REPORTING DATE

RHEUMATOID (RA) FACTOR QUANTITATIVE: NEGATIVE: < 18.0 **SERUM** BORDERLINE: 18.0 - 25.0

by NEPHLOMETRY POSITIVE: > 25.0

INTERPRETATION:

CLIENT CODE.

RHEUMATOID FACTOR (RA):

1. Rheumatoid factors (RF) are antibodies that are directed against the Fc fragment of IgG altered in its tertiary structure

2. Over 75% of patients with rheumatoid arthritis (RA) have an IgM antibody to IgG immunoglobulin. This autoantibody (RF) is diagnostically useful although it may not be etiologically related to RA.

Inflammatory Markers such as ESR & C-Reactive protein (CRP) are normal in about 60 % of patients with positive RA.

Inflammatory Markers poorly with disease activity, but those patients with high titers tend to have more severe disease course.

The test is useful for diagnosis and prognosis of rheumatoid arthritis.

RHEUMATOID ARTHIRITIS:

The patients with related to RA.

The patients with positive RA.

The patien

- 1. Rheumatoid Arthiritis is a systemic autoimmune disease that is multi-functional in origin and is characterized by chronic inflammation of the

membrane lining (synovium) joints which ledas to progressive joint destruction and in most cases to disability and reduction of quality life.

2. The disease spredas from small to large joints, with greatest damage in early phase.

3. The diagnosis of RA is primarily based on clinical, radiological & immunological features. The most frequent serological test is the measurement of RA factor.

CAUTION (FALSE POSTIVE):
1. PA factor is not specific for Phosometrial arthirities as it is often present in healthy individuals with other autoimmuno diseases and chronic infect.

- 1. RA factor is not specific for Rheumatoid arthiritis, as it is often present in healthy individuals with other autoimmune diseases and chronic infections. 2. Non rheumatoid arthritis (RA) populations are not clearly separate with regard to the presence of rheumatoid factor (RF) (15% of RA patients have a nonreactive titer and 8% of nonrheumatoid patients have a positive titer).

 3. Patients with various nonrheumatoid diseases, characterized by chronic inflammation may have positive tests for RF. These diseases include systemic lupus erythematosus, polymyositis, tuberculosis, syphilis, viral hepatitis, infectious mononucleosis, and influenza.

 4. Anti-CCP have been discovered in joints of patients with RA, but not in other form of joint disease. Anti-CCP2 is HIGHLY SENSITIVE (71%) & more
- specific (98%) than RA factor

5. Upto 30 % of patients with Seronegative Rheumatoid arthiritis also show Anti-CCP antibodies.

6. The positive predictive value of Anti-CCP antibodies for Rheumatoid Arthiritis is far greater than Rheumatoid factor.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY :012409290030 : SURJESH REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 12:22PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

VITAMINS

VITAMIN D/25 HYDROXY VITAMIN D3

VITAMIN D (25-HYDROXY VITAMIN D3): SERUM 41.858 ng/mL DEFICIENCY: < 20.0

by CLIA (CHEMILUMINESCENCE IMMUNOASSAY) INSUFFICIENCY: 20.0 - 30.0

SUFFICIENCY: 30.0 - 100.0 TOXICITY: > 100.0

INTERPRETATION:

DEFICIENT:	< 20	ng/mL
INSUFFICIENT:	21 - 29	ng/mL
PREFFERED RANGE:	30 - 100	ng/mL
INTOXICATION:	> 100	ng/ml

1. Vitamin D compounds are derived from dietary ergocalciferol (from plants, Vitamin D2), or cholecalciferol (from animals, Vitamin D3), or by conversion of 7- dihydrocholecalciferol to Vitamin D3 in the skin upon Ultraviolet exposure.

2.25-OH--Vitamin D represents the main body resevoir and transport form of Vitamin D and transport form of Vitamin D, being stored in adipose tissue and tightly bound by a transport protein while in circulation.

3. Vitamin D plays a primary role in the maintenance of calcium homeostatis. It promotes calcium absorption, renal calcium absorption and phosphate reabsorption, skeletal calcium deposition, calcium mobilization, mainly regulated by parathyroid harmone (PTH).

4. Severe deficiency may lead to failure to mineralize newly formed osteoid in bone, resulting in rickets in children and osteomalacia in adults.

DECREASED:

- 1.Lack of sunshine exposure.
- 2.Inadequate intake, malabsorption (celiac disease)
 3.Depressed Hepatic Vitamin D 25- hydroxylase activity
- 4. Secondary to advanced Liver disease
- 5. Osteoporosis and Secondary Hyperparathroidism (Mild to Moderate deficiency)
- 6.Enzyme Inducing drugs: anti-epileptic drugs like phenytoin, phenobarbital and carbamazepine, that increases Vitamin D metabolism.

1. Hypervitaminosis D is Rare, and is seen only after prolonged exposure to extremely high doses of Vitamin D. When it occurs, it can result in severe hypercalcemia and hyperphophatemia.

CAUTION: Replacement therapy in deficient individuals must be monitored by periodic assessment of Vitamin D levels in order to prevent hypervitaminosis D

NOTE:-Dark coloured individuals as compare to whites, is at higher risk of developing Vitamin D deficiency due to excess of melanin pigment which interefere with Vitamin D absorption.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 29/Sep/2024 01:03PM

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH :012409290030 REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

VITAMIN B12/COBALAMIN

REPORTING DATE

VITAMIN B12/COBALAMIN: SERUM 313.13 190.0 - 830 pg/mL

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

INTERPRETATION:-

CLIENT CODE.

INCREASED VITAMIN B12	DECREASED VITAMIN B12
1.Ingestion of Vitamin C	1.Pregnancy
2.Ingestion of Estrogen	2.DRUGS:Aspirin, Anti-convulsants, Colchicine
3.Ingestion of Vitamin A	3.Ethanol Igestion
4.Hepatocellular injury	4. Contraceptive Harmones
5.Myeloproliferative disorder	5.Haemodialysis
6.Uremia	6. Multiple Myeloma

- 1. Vitamin B12 (cobalamin) is necessary for hematopoiesis and normal neuronal function.
- 2.In humans, it is obtained only from animal proteins and requires intrinsic factor (IF) for absorption.
- 3. The body uses its vitamin B12 stores very economically, reabsorbing vitamin B12 from the ileum and returning it to the liver; very little is excreted
- 4.Vitamin B12 deficiency may be due to lack of IF secretion by gastric mucosa (eg, gastrectomy, gastric atrophy) or intestinal malabsorption (eg, ileal resection, small intestinal diseases).
- 5. Vitamin B12 deficiency frequently causes macrocytic anemia, glossitis, peripheral neuropathy, weakness, hyperreflexia, ataxia, loss of proprioception, poor coordination, and affective behavioral changes. These manifestations may occur in any combination; many patients have the neurologic defects without macrocytic anemia.
- 6.Serum methylmalonic acid and homocysteine levels are also elevated in vitamin B12 deficiency states.
- 7. Follow-up testing for antibodies to intrinsic factor (IF) is recommended to identify this potential cause of vitamin B12 malabsorption. NOTE:A normal serum concentration of vitamin B12 does not rule out tissue deficiency of vitamin B12. The most sensitive test for vitamin B12 deficiency at the cellular level is the assay for MMA. If clinical symptoms suggest deficiency, measurement of MMA and homocysteine should be considered, even if serum vitamin B12 concentrations are normal.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY :012409290030 : SURJESH REG. NO./LAB NO.

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 01:45PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

VITAMIN B9/FOLIC ACID/FOLATE

VITAMIN B9/FOLIC ACID/FOLATE: SERUM by CLIA (CHEMILUMINESCENCE IMMUNOASSAY) 9.98 DEFICIENT: < 3.37

INTERMEDIATE: 3.37 - 5.38

NORMAL: > 5.38

INTERPRETATION

RESULT IN ng/mL	REMARKS
0.35 – 3.37	DEFICIENT
3.38 - 5.38	INTERMEDIATE
5.39 - 100.00	NORMAL

NOTE:

- 1. Drugs like Methotrexate & Leucovorin interfere with folate measurement
- 2. To differentiate vitamin B12 & folate deficiency, measurement of Methyl malonic acid in urine & serum Homocysteine level is suggested 3. Risk of toxicity from folic acid is low as it is a water soluble vitamin regularly excreted in urine
- COMMENTS:

- 1. Folate plays an important role in the synthesis of purine & pyrimidines in the body and is important for the maturation of erythrocytes.

 2. It is widely available from plants and to a lesser extent organ meats, but more than half the folate content of food is lost during cooking.

 3. Folate deficiency is commonly prevalent in alcoholic liver disease, pregnancy and the elderly. It may result from poor intestinal absorption, nutrition deficiency, excessive demand as in pregnancy or in malignancy and in response to certain drugs like Methotrexate & anticonvulsants.

 4. Decreased Levels Megaloblastic anemia, Infantile hyperthyroidism, Alcoholism, Malnutrition, Scurvy, Liver disease, B12 deficiency, dietary amino acid excess, adult Celiac disease, Tropical Sprue, Crohn's disease, Hemolytic anemias, Carcinomas, Myelofibrosis, vitamin B6 deficiency, pregnancy, Whippley's disease, extensive intesting and syvers exfeliative dermatitis. pregnancy, Whipple's disease, extensive intestinal resection and severe exfoliative dermatitis



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY : SURJESH REG. NO./LAB NO. :012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM BARCODE NO. :01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 29/Sep/2024 04:53PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval**

CLINICAL PATHOLOGY

URINE ROUTINE & MICROSCOPIC EXAMINATION

PHYSICAL EXAMINATION

QUANTITY RECIEVED	10	ml
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY		

AMBER YELLOW PALE YELLOW

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

TRANSPARANCY CLEAR CLEAR

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY 1.002 - 1.030 SPECIFIC GRAVITY 1.01

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

CHEMICAL EXAMINATION

ACIDIC

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY **PROTEIN NEGATIVE (-ve)** Trace

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

NEGATIVE (-ve) SUGAR Negative by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

рΗ 5.0 - 7.56.5

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY **NEGATIVE (-ve) BILIRUBIN** Negative

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

NITRITE Negative **NEGATIVE** (-ve)

EU/dL UROBILINOGEN Normal 0.2 - 1.0

KETONE BODIES **NEGATIVE (-ve)** Negative

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

BLOOD Negative **NEGATIVE (-ve)** by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

NEGATIVE (-ve) **NEGATIVE (-ve)** ASCORBIC ACID by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

MICROSCOPIC EXAMINATION

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY.

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mrs. SEEMA GARG

AGE/ GENDER : 52 YRS/FEMALE **PATIENT ID** : 1628908

COLLECTED BY: SURJESH REG. NO./LAB NO. : 012409290030

REFERRED BY : CENTRAL PHOENIX CLUB (AMBALA CANTT) **REGISTRATION DATE** : 29/Sep/2024 10:34 AM **BARCODE NO.** : 01517942 **COLLECTION DATE** : 29/Sep/2024 04:29PM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 29/Sep/2024 04:53PM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

T 131			D: 1 : 1D (
Test Name	Value	Unit	Biological Reference interval
RED BLOOD CELLS (RBCs) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)	/HPF	0 - 3
PUS CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	1-3	/HPF	0 - 5
EPITHELIAL CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	2-4	/HPF	ABSENT
CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
BACTERIA by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
OTHERS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
TRICHOMONAS VAGINALIS (PROTOZOA) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	ABSENT		ABSENT

*** End Of Report **



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana
KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana
0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com