

KOS Diagnostic Lab (A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. VIKAS

AGE/ GENDER : 53 YRS/MALE **PATIENT ID** : 1498256

COLLECTED BY :012409300007 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:13 AM BARCODE NO. :01517980 **COLLECTION DATE** : 30/Sep/2024 12:32PM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Value Unit **Biological Reference interval** Test Name

CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F) AND POST PRANDIAL (PP)

REPORTING DATE

GLUCOSE FASTING (F): PLASMA mg/dL NORMAL: < 100.0 123.71^H

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0

DIABETIC: > 0R = 126.0

: 30/Sep/2024 02:46PM

GLUCOSE POST PRANDIAL (PP): PLASMA 132.94 mg/dL NORMAL: < 140.00 by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)

PREDIABETIC: 140.0 - 200.0 DIABETIC: > OR = 200.0

INTERPRETATION:

CLIENT CODE.

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

- 1. A fasting plasma glucose below 100 mg/dL and post-prandial plasma glucose level below 140 mg/dl is considered normal.
- 2. A fasting plasma glucose level between 100 125 mg/dl and post-prandial plasma glucose level between 140 200 mg/dL is considered as glucose intolerant or pre diabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
- 3. A fasting plasma glucose level of above 125 mg/dL and post-prandial plasma glucose level above 200 mg/dL is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

* End Of Report ***



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

