

# KOS Diagnostic Lab (A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mr. GULSHAN AHUJA

AGE/ GENDER : 41 YRS/MALE PATIENT ID : 1629326

COLLECTED BY : REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 30/Sep/2024 09:16 AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

# SWASTHYA WELLNESS PANEL: 1.5 COMPLETE BLOOD COUNT (CBC)

#### **RED BLOOD CELLS (RBCS) COUNT AND INDICES**

HAEMOGLOBIN (HB) by CALORIMETRIC	15.2	gm/dL	12.0 - 17.0
RED BLOOD CELL (RBC) COUNT by Hydro dynamic focusing, electrical impedence	5.47 <sup>H</sup>	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	45.6	%	40.0 - 54.0
MEAN CORPUSCULAR VOLUME (MCV) by Calculated by automated hematology analyzer	83.2	fL	80.0 - 100.0
MEAN CORPUSCULAR HAEMOGLOBIN (MCH) by Calculated by automated hematology analyzer	27.8	pg	27.0 - 34.0
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	33.4	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION WIDTH (RDW-CV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	12.9	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	40.2	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED	15.21	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX by CALCULATED	19.63	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CELLS (WBCS)			
TOTAL LEUCOCYTE COUNT (TLC) by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	5030	/cmm	4000 - 11000
NUCLEATED RED BLOOD CELLS (nRBCS) by automated 6 part hematology analyzer	NIL		0.00 - 20.00
NUCLEATED RED BLOOD CELLS (nRBCS) % by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	NIL	%	< 10 %
DIFFERENTIAL LEUCOCYTE COUNT (DLC)			
NEUTROPHILS by Flow cytometry by SF cube & microscopy	43 <sup>L</sup>	%	50 - 70



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





CLIENT CODE.

### **KOS Diagnostic Lab**

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 30/Sep/2024 09:16AM

REPORTING DATE

**NAME** : Mr. GULSHAN AHUJA

**AGE/ GENDER PATIENT ID** : 1629326 : 41 YRS/MALE

**COLLECTED BY** REG. NO./LAB NO. : 012409300009

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM **COLLECTION DATE** BARCODE NO. :01517982 : 30/Sep/2024 08:29AM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
LYMPHOCYTES by Flow cytometry by SF cube & microscopy	44 <sup>H</sup>	%	20 - 40
OSINOPHILS by flow cytometry by sf cube & microscopy	3	%	1 - 6
MONOCYTES by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	10	%	2 - 12
BASOPHILS by flow cytometry by sf cube & microscopy ABSOLUTE LEUKOCYTES (WBC) COUNT	0	%	0 - 1
ABSOLUTE NEUTROPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	2163	/cmm	2000 - 7500
ABSOLUTE LYMPHOCYTE COUNT by flow cytometry by sf cube & microscopy	2213	/cmm	800 - 4900
BSOLUTE EOSINOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	151	/cmm	40 - 440
BSOLUTE MONOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	503	/cmm	80 - 880
NBSOLUTE BASOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY PLATELETS AND OTHER PLATELET PREDICTIVE MARKE	0 <b>RS</b>	/cmm	0 - 110
PLATELET COUNT (PLT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	191000	/cmm	150000 - 450000
LATELETCRIT (PCT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	0.22	%	0.10 - 0.36
MEAN PLATELET VOLUME (MPV) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	12	fL	6.50 - 12.0
LATELET LARGE CELL COUNT (P-LCC) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	69000	/cmm	30000 - 90000
LATELET LARGE CELL RATIO (P-LCR) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	36.2	%	11.0 - 45.0
LATELET DISTRIBUTION WIDTH (PDW) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD	16.6	%	15.0 - 17.0



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 30/Sep/2024 02:05PM

NAME : Mr. GULSHAN AHUJA

AGE/ GENDER : 41 YRS/MALE PATIENT ID : 1629326

COLLECTED BY : REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE

 CLIENT ADDRESS
 : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

#### **GLYCOSYLATED HAEMOGLOBIN (HBA1C)**

GLYCOSYLATED HAEMOGLOBIN (HbA1c): 5.6 % 4.0 - 6.4

WHOLE BLOOD

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

ESTIMATED AVERAGE PLASMA GLUCOSE 114.02 mg/dL 60.00 - 140.00

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

#### **INTERPRETATION:**

AS PER AMERICAN DI	ABETES ASSOCIATION (ADA):	
REFERENCE GROUP	GLYCOSYLATED HEMOGL	OGIB (HBAIC) in %
Non diabetic Adults >= 18 years	<5.7	
At Risk (Prediabetes)	5.7 – 6.	4
Diagnosing Diabetes	>= 6.5	
	Age > 19 Years	
	Goals of Therapy:	< 7.0
Therapeutic goals for glycemic control	Actions Suggested:	>8.0
	Age < 19 Y	ears
	Goal of therapy:	<7.5

#### COMMENTS:

- 1.Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients. 2. Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high
- concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.

  3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive competitions, targetting a goal of < 7.0% may not be
- patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be appropriate.

  4. High HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications
- 5. Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.
  6. HbA1c results from patients with HbSS HbSC and HbD must be interpreted with caution, given the pathological processes including anemia increase
- 6.HbA1c results from patients with HbSS,HbSC and HbD must be interpreted with caution, given the pathological processes including anemia,increased red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.
- 7. Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

:012409300009 **COLLECTED BY** REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 30/Sep/2024 09:25AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

### ERYTHROCYTE SEDIMENTATION RATE (ESR)

**ERYTHROCYTE SEDIMENTATION RATE (ESR)** 

0 - 20

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

#### INTERPRETATION:

- 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
- 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
- 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

- ESR and C reactive protein (C-RP) are both markers of inflammation.
   Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
   CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
   If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
- 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
- 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: Mr. GULSHAN AHUJA **NAME** 

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** :012409300009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE :30/Sep/2024 10:18AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

### CLINICAL CHEMISTRY/BIOCHEMISTRY **GLUCOSE FASTING (F)**

**GLUCOSE FASTING (F): PLASMA** 102.19H mg/dL NORMAL: < 100.0

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0

INTERPRETATION
IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

COLLECTED BY : REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 30/Sep/2024 10:18 AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
	LIPID PROFILE	: BASIC	
CHOLESTEROL TOTAL: SERUM by CHOLESTEROL OXIDASE PAP	139.95	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SERUM by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)	116.71	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (DIRECT): SERUM by SELECTIVE INHIBITION	28.6 <sup>L</sup>	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	88.01	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	111.35	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	23.34	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SERUM by CALCULATED, SPECTROPHOTOMETRY	396.61	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	4.89 <sup>H</sup>	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0
LDL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	3.08 <sup>H</sup>	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

:30/Sep/2024 10:18AM

: Mr. GULSHAN AHUJA **NAME** 

AGE/ GENDER : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** :012409300009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
TRIGLYCERIDES/HDL RATIO: SERUM	4.08	RATIO	3.00 - 5.00
by CALCULATED, SPECTROPHOTOMETRY			

REPORTING DATE

#### INTERPRETATION:

CLIENT CODE.

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. GULSHAN AHUJA

AGE/ GENDER : 41 YRS/MALE PATIENT ID : 1629326

COLLECTED BY : REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 30/Sep/2024 10:18 AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

LIVER FUNCTION TEST (COMPLE	IL)
-----------------------------	-----

BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY	0.62	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY	0.19	mg/dL	0.00 - 0.40
BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY	0.43	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	15.1	U/L	7.00 - 45.00
SGPT/ALT: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	25.4	U/L	0.00 - 49.00
AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	0.59	RATIO	0.00 - 46.00
ALKALINE PHOSPHATASE: SERUM by Para nitrophenyl phosphatase by amino methyl propanol	93.38	U/L	40.0 - 130.0
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY	65.79 <sup>H</sup>	U/L	0.00 - 55.0
TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY	6.54	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL GREEN	3.79	gm/dL	3.50 - 5.50
GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.75	gm/dL	2.30 - 3.50
A : G RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	1.38	RATIO	1.00 - 2.00

#### INTERPRETATION

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range.

**USE**:- Differential diagnosis of diseases of hepatobiliary system and pancreas.

#### INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

COLLECTED BY: REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

**CLIENT CODE.** : KOS DIAGNOSTIC LAB **REPORTING DATE** : 30/Sep/2024 10:18AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

#### **DECREASED:**

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

#### PROGNOSTIC SIGNIFICANCE:

1 KO CITOCTTO CICITII TOTILICE	
NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUĞAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana
KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana
0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** REG. NO./LAB NO. :012409300009

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 30/Sep/2024 10:18AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
k	IDNEY FUNCTION TE	ST (COMPLETE)	
UREA: SERUM  by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)	27.06	mg/dL	10.00 - 50.00
CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY	0.95	mg/dL	0.40 - 1.40
BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETRY	12.64	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	13.31	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	28.48	RATIO	
URIC ACID: SERUM by URICASE - OXIDASE PEROXIDASE	7.24	mg/dL	3.60 - 7.70
CALCIUM: SERUM by ARSENAZO III, SPECTROPHOTOMETRY	9.18	mg/dL	8.50 - 10.60
PHOSPHOROUS: SERUM by PHOSPHOMOLYBDATE, SPECTROPHOTOMETRY	3.17	mg/dL	2.30 - 4.70
ELECTROLYTES			
SODIUM: SERUM by ISE (ION SELECTIVE ELECTRODE)	140.3	mmol/L	135.0 - 150.0
POTASSIUM: SERUM by ISE (ION SELECTIVE ELECTRODE)	4.96	mmol/L	3.50 - 5.00
CHLORIDE: SERUM by ISE (ION SELECTIVE ELECTRODE)	105.23	mmol/L	90.0 - 110.0
FSTIMATED GLOMERI II AR FII TERATION RATE			

#### ESTIMATED GLOMERULAR FILTERATION RATE

ESTIMATED GLOMERULAR FILTERATION RATE 103.1

(eGFR): SERUM by CALCULATED

#### **INTERPRETATION:**

To differentiate between pre- and post renal azotemia.

#### INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

- 1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.
- 2. Catabolic states with increased tissue breakdown.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. GULSHAN AHUJA

AGE/ GENDER : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** :012409300009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE :30/Sep/2024 10:18AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

- 3. GI haemorrhage.
- 4. High protein intake.
- 5. Impaired renal function plus
- 6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushing's syndrome, high protein diet, burns, surgery, cachexia, high fever).
- 7. Urine reabsorption (e.g. ureter colostomy)
- 8. Reduced muscle mass (subnormal creatinine production)
- 9. Certain drugs (e.g. tetracycline, glucocorticoids)

#### INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2. Prerenal azotemia superimposed on renal disease.

#### DECREASED RATIO (<10:1) WITH DECREASED BUN:

- 1. Acute tubular necrosis.
- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.
- 5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
- 8. Pregnancy.

#### **DECREASED RATIO (<10:1) WITH INCREASED CREATININE:**

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure.

#### **INAPPROPIATE RATIO:**

- 1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).
- 2. Cephalosporin therapy (interferes with creatinine measurement). ESTIMATED GLOMERULAR FILTERATION RATE:

ESTIMATED GEOIMERGEART	ILI LIU TITOTI TUTTE.		
CKD STAGE	DESCRIPTION	GFR ( mL/min/1.73m2 )	ASSOCIATED FINDINGS
G1	Normal kidney function	>90	No proteinuria
G2	Kidney damage with	>90	Presence of Protein,
	normal or high GFR		Albumin or cast in urine
G3a	Mild decrease in GFR	60 -89	
G3b	Moderate decrease in GFR	30-59	
G4	Severe decrease in GFR	15-29	
G5	Kidney failure	<15	



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 30/Sep/2024 10:18AM

**NAME** : Mr. GULSHAN AHUJA

AGE/ GENDER : 41 YRS/MALE **PATIENT ID** : 1629326

:012409300009 COLLECTED BY REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

COMMENTS:

CLIENT CODE.

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.

2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure

REPORTING DATE

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** :012409300009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE :30/Sep/2024 10:18AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference in
--

	IRON PROFILE		
IRON: SERUM by FERROZINE, SPECTROPHOTOMETRY	49.8 <sup>L</sup>	μg/dL	59.0 - 158.0
UNSATURATED IRON BINDING CAPACITY (UIBC) :SERUM by FERROZINE, SPECTROPHOTOMETERY	220.04	μg/dL	150.0 - 336.0
TOTAL IRON BINDING CAPACITY (TIBC) :SERUM by SPECTROPHOTOMETERY	269.84	μg/dL	230 - 430
%TRANSFERRIN SATURATION: SERUM by CALCULATED, SPECTROPHOTOMETERY (FERENE)	18.46	%	15.0 - 50.0
TRANSFERRIN: SERUM by SPECTROPHOTOMETERY (FERENE)	191.59 <sup>L</sup>	mg/dL	200.0 - 350.0

#### INTERPRETATION:-

VARIABLES	ANEMIA OF CHRONIC DISEASE	IRON DEFICIENCY ANEMIA	THALASSEMIA α/β TRAIT
SERUM IRON:	Normal to Reduced	Reduced	Normal
TOTAL IRON BINDING CAPACITY:	Decreased	Increased	Normal
% TRANSFERRIN SATURATION:	Decreased	Decreased < 12-15 %	Normal
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increased

#### **IRON:**

- 1. Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia. i.e iron deficiency anemia, zinc deficiency anemia, anemia of chronic disease and thalassemia syndromes.
- 2. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.

TOTAL IRON BINDING CAPACITY (TIBC):

1.It is a direct measure of protein transferrin which transports iron from the gut to storage sites in the bone marrow.

% TRANSFERRIN SATURATION:

1.Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: Mr. GULSHAN AHUJA **NAME** 

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** :012409300009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM

CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 30/Sep/2024 10:42AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

### **ENDOCRINOLOGY**

#### THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM	0.754	ng/mL	0.35 - 1.93
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			
THYROXINE (T4): SERUM	7.34	μgm/dL	4.87 - 12.60
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			
THADOID CLIVINI WALING HODINONE (ACH). CEDINA	2.70	ull I /ml	0.25 5.50

THYROID STIMULATING HORMONE (TSH): SERUM

μIU/mL 0.35 - 5.50 by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

#### INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and trilodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	Т3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs
- 3. Serum T4 levies in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMUL	ATING HORMONE (TSH)
Age	Refferance Range (ng/mL)	Age	Refferance Range (μg/dL)	Age	Reference Range ( μΙυ/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 30/Sep/2024 10:42AM

: Mr. GULSHAN AHUJA **NAME** 

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** REG. NO./LAB NO. :012409300009

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name			Value	Unit		Biological Reference interval
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00	
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50	
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50	
	RECOM	MENDATIONS OF TSH LI	VELS DURING PRE	GNANCY ( µIU/mL)		
	1st Trimester			0.10 - 2.50		
	2nd Trimester		0.20 - 3.00			
	3rd Trimester			0.30 - 4.10	•	

REPORTING DATE

#### **INCREASED TSH LEVELS:**

CLIENT CODE.

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

#### **DECREASED TSH LEVELS:**

- 1.Toxic multi-nodular goitre & Thyroiditis.
- 2. Over replacement of thyroid harmone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituatary or hypothalmic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. GULSHAN AHUJA

AGE/ GENDER : 41 YRS/MALE PATIENT ID : 1629326

COLLECTED BY: REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 30/Sep/2024 11:26 AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

#### PROCALCITONIN (PCT)

PROCALCITONIN (PCT): SERUM 0.1 ng/mL < 0.50

by ELFA (ENZYME LINKED FLOUROSCENCE ASSAY)

#### **INTERPRETATION:**

Procalcitonin, the prohormone of calcitonin is below limit of detection 500 pg/ml in healthy individuals. It rises in response to an inflammatory stimulus especially of bacterial origin. It does not rise significantly with viral or non inflammations.

PROCALCITONIN (VALUE IN ng/mL)	INFERENCE
< 0.50 ng/mL	Minor local bacterial infection is possible. Severe systemic infection (Sepsis) is not likely
0.50- < 2.0 ng/mL	Systemic infection is possible, but various conditions are known to induce PCT as well (see below). Suggest repeat after 6-24 hours for a definitive diagnosis
2.0 - < 10.0 ng/mL	Systemic infection (Sepsis) is likely, unless other causes are known
>=10.0 ng/mL	Important systemic inflammatory response, almost exclusively due to severe bacterial sepsis or septic shock

#### PCT levels can be elevated in non infectious causes like:

- 1. The first days after a major trauma, major surgical intervention, burns, treatment with OKT3 antibodies and other drugs stimulating the release of pro-inflammatory cytokines, small cell lung cancer, medullary C-cell carcinoma of thyroid.
- 2. Patients with prolonged or severe cardiogenic shock, prolonged severe organ perfusion anomalies.
- 3. Neonates < 48 hrs of life.
- 4.Patients with PCT values 2000 pg/mL should be closely monitored both clinically and by reassessing PCT within 6-24 hrs.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





# KOS Diagnostic Lab (A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. GULSHAN AHUJA

AGE/ GENDER : 41 YRS/MALE PATIENT ID : 1629326

COLLECTED BY : REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 30/Sep/2024 11:26 AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

# IMMUNOPATHOLOGY/SEROLOGY CHIKUNGUNYA ANTIBODY IgM SCREENING

CHIKUNGUNYA ANTIBODY IgM QUANTITATIVE 0.14 INDEX VALUE < 1.00

by ELISA (ENZYME LINKED IMMUNOASSAY)

#### INTERPRETATION

- 1.Chikungunya is an insect borne viral disease belonging to genus ALPHAVIRIDAE transmitted to humans by infected Aedes mosquitoe
- 2.lt causes fever and severe joint pain. Other symptoms include muscle pain, headache, nausea, fatigue and rash.
- 3. Joint pain is often debilitating and can vary in duration.
- 4.The disease shares some clinical signs with dengue, and can be misdiagnosed in areas where dengue is common.
- $5. There is no cure for the disease. \ Treatment is focused on relieving the symptoms.\\$

#### **ROUTE OF TRANSMISSION:**

- 1. The virus is transmitted from human to human by the bites of infected female mosquitoes.
- 2.Most commonly, the mosquitoes involved are Aedes aegypti andAedes albopictus, two species which can also transmit other mosquito-borne viruses, including dengue.
- 3. These mosquitoes can be found biting throughout daylight hours, though there may be peaks of activity in the early morning and late afternoon.
- 4. Both species are found biting outdoors, but Ae. aegypti will also readily feed indoors.
- 5. After the bite of an infected mosquito, onset of illness occurs usually between 4 and 8 days but can range from 2 to 12 days.

#### NOTE:

This is a solid phase immunochromatographic assay for the detection of the Chikungunya specific IgM antibodies in the human serum. The test has a sensitivity of 97.5 % and a specificity of 99.1 %.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** :012409300009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE :30/Sep/2024 10:18AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

#### **C-REACTIVE PROTEIN (CRP)**

C-REACTIVE PROTEIN (CRP) QUANTITATIVE:  $7.5^{H}$ mg/L 0.0 - 6.0

**SERUM** 

by NEPHLOMETRY **INTERPRETATION:** 

1. C-reactive protein (CRP) is one of the most sensitive acute-phase reactants for inflammation.

2. CRP levels can increase dramatically (100-fold or more) after severe trauma, bacterial infection, inflammation, surgery, or neoplastic proliferation.

3. CRP levels (Quantitative) has been used to assess activity of inflammatory disease, to detect infections after surgery, to detect transplant

rejection, and to monitor these inflammatory processes.

4. As compared to ESR, CRP shows an earlier rise in inflammatory disorders which begins in 4-6 hrs, the intensity of the rise being higher than ESR and the recovery being earlier than ESR. Unlike ESR, CRP levels are not influenced by hematologic conditions like Anemia, Polycythemia etc., 5. Elevated values are consistent with an acute inflammatory process.

NOTE:

1. Elevated C-reactive protein (CRP) values are nonspecific and should not be interpreted without a complete clinical history.

2. Oral contraceptives may increase CRP levels.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra

MD (Pathology)

CEO & Consultant Pathologist

NAME : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

COLLECTED BY : REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 30/Sep/2024 09:46 AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

#### DENGUE FEVER COMBO SCREENING - (NS1 ANTIGEN, IgG AND IgM)

DENGUE NS1 ANTIGEN - SCREENING
by ICT (IMMUNOCHROMATOGRAPHY)

DENGUE ANTIBODY IgG - SCREENING
by ICT (IMMUNOCHROMATOGRAPHY)

DENGUE ANTIBODY IGM - SCREENING

NEGATIVE (-ve)

NEGATIVE (-ve)

NEGATIVE (-ve)

DENGUE ANTIBODY IgM - SCREENING NEGATIVE (-ve)

by ICT (IMMUNOCHROMATOGRAPHY)

NEGATIVE (-ve)

#### **INTERPRETATION:**-

- 1. This is a solid phase immunochromatographic ELISA test for the qualitative detection of the specific IgG and IgM antibodies against the Dengue virus.
- 2.The IgM antibodies take a minimum of 5-10 days in primary infection and 4-5 days in secondary infections to test positive and hence are suitable for the diagnosis of dengue fever only when the fever is approximately one week old.
- 3.The IgG antibodies develop at least two weeks after exposure to primary infection and subsequently remain positive for the rest of the life. A positive result is incapable of differentiating a current infection from a past infection.
- 4.The Dengue NS-1 antigen test is most suited for early diagnosis (within the first week of exposure).



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. GULSHAN AHUJA

AGE/ GENDER : 41 YRS/MALE PATIENT ID : 1629326

COLLECTED BY : REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 30/Sep/2024 09:25 AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

#### WIDAL SLIDE AGGLUTINATION TEST

SALMONELLA TYPHI O	1:20	TITRE	1:80
by SLIDE AGGLUTINATION			
SALMONELLA TYPHI H	NIL	TITRE	1:160
by SLIDE AGGLUTINATION			
SALMONELLA PARATYPHI AH	NIL	TITRE	1:160
by SLIDE AGGLUTINATION			
SALMONELLA PARATYPHI BH	NIL	TITRE	1:160
by SLIDE AGGLUTINATION			

#### **INTERPRETATION:**

- 1.Titres of 1:80 or more for "O" agglutinin is considered significant.
- 2. Titres of 1:160 or more for "H" agglutinin is considered significant.

#### LIMITATIONS

- 1. Agglutinins usually appear by 5th to 6th day of illness of enteric fever, hence a negative result in early stage is inconclusive. The titre then rises till 3rd or 4th week, after which it declines gradually.
- 2.Lower titres may be found in normal individuals.
- 3.A single positive result has less significance than the rising agglutination titre, since demonstration of rising titre four or more in 1st and 3rd week is considered as a definite evidence of infection.
- 4.A simultaneous rise in H agglutinins is suggestive of paratyphoid infection.

#### NOTE:

- 1.Individuals with prior infection or immunization with TAB vaccine may develop an ANAMNESTIC RESPONSE (False-Positive) during an unrelated fever i.e High titres of antibodies to various antigens. This may be differentiated by repitition of the test after a week.
- 2. The anamnestic response shows only a transient rise, while in enteric fever rise is sustained.
- 3.H agglutinins tend to persist for many months after vaccination but O agglutinins tend to disappear sooner i.e within 6 months. Therefore rise in Oagglutinins indicate recent infection.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** :012409300009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE :30/Sep/2024 10:48AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

#### **VITAMINS**

#### VITAMIN D/25 HYDROXY VITAMIN D3

VITAMIN D (25-HYDROXY VITAMIN D3): SERUM by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

20.35<sup>L</sup>

ng/mL

**DEFICIENCY:** < 20.0

INSUFFICIENCY: 20.0 - 30.0 SUFFICIENCY: 30.0 - 100.0

**TOXICITY: > 100.0** 

**INTERPRETATION:** 

DEFICIENT:	< 20	ng/mL
INSUFFICIENT:	21 - 29	ng/mL
PREFFERED RANGE:	30 - 100	ng/mL
INTOXICATION:	> 100	ng/mL

- 1. Vitamin D compounds are derived from dietary ergocalciferol (from plants, Vitamin D2), or cholecalciferol (from animals, Vitamin D3), or by conversion of 7- dihydrocholecalciferol to Vitamin D3 in the skin upon Ultraviolet exposure.
- 2.25-OH--Vitamin D represents the main body resevoir and transport form of Vitamin D and transport form of Vitamin D, being stored in adipose tissue and tightly bound by a transport protein while in circulation.
- 3. Vitamin D plays a primary role in the maintenance of calcium homeostatis. It promotes calcium absorption, renal calcium absorption and phosphate reabsorption, skeletal calcium deposition, calcium mobilization, mainly regulated by parathyroid harmone (PTH).
- 4. Severe deficiency may lead to failure to mineralize newly formed osteoid in bone, resulting in rickets in children and osteomalacia in adults. **DECREASED:**
- 1.Lack of sunshine exposure.
- 2.Inadequate intake, malabsorption (celiac disease)
  3.Depressed Hepatic Vitamin D 25- hydroxylase activity
- 4. Secondary to advanced Liver disease
- 5. Osteoporosis and Secondary Hyperparathroidism (Mild to Moderate deficiency)
- 6.Enzyme Inducing drugs: anti-epileptic drugs like phenytoin, phenobarbital and carbamazepine, that increases Vitamin D metabolism.
- 1. Hypervitaminosis D is Rare, and is seen only after prolonged exposure to extremely high doses of Vitamin D. When it occurs, it can result in severe hypercalcemia and hyperphophatemia.

CAUTION: Replacement therapy in deficient individuals must be monitored by periodic assessment of Vitamin D levels in order to prevent hypervitaminosis D

NOTE:-Dark coloured individuals as compare to whites, is at higher risk of developing Vitamin D deficiency due to excess of melanin pigment which interefere with Vitamin D absorption.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

pg/mL

190.0 - 830

**NAME** : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

**COLLECTED BY** :012409300009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM BARCODE NO. :01517982 **COLLECTION DATE** : 30/Sep/2024 08:29AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE :30/Sep/2024 10:59AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit **Biological Reference interval** 

#### **VITAMIN B12/COBALAMIN**

VITAMIN B12/COBALAMIN: SERUM

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

INTERPRETATION:-

MATERIA RETATION:	
INCREASED VITAMIN B12	DECREASED VITAMIN B12
1.Ingestion of Vitamin C	1.Pregnancy
2.Ingestion of Estrogen	2.DRUGS:Aspirin, Anti-convulsants, Colchicine
3.Ingestion of Vitamin A	3.Ethanol Igestion
4.Hepatocellular injury	4. Contraceptive Harmones
5.Myeloproliferative disorder	5.Haemodialysis
6.Uremia	6. Multiple Myeloma

186.04<sup>L</sup>

- 1. Vitamin B12 (cobalamin) is necessary for hematopoiesis and normal neuronal function.
- 2.In humans, it is obtained only from animal proteins and requires intrinsic factor (IF) for absorption.
- 3. The body uses its vitamin B12 stores very economically, reabsorbing vitamin B12 from the ileum and returning it to the liver; very little is
- 4. Vitamin B12 deficiency may be due to lack of IF secretion by gastric mucosa (eg, gastrectomy, gastric atrophy) or intestinal malabsorption (eg, ileal resection, small intestinal diseases).
- 5.Vitamin B12 deficiency frequently causes macrocytic anemia, glossitis, peripheral neuropathy, weakness, hyperreflexia, ataxia, loss of proprioception, poor coordination, and affective behavioral changes. These manifestations may occur in any combination; many patients have the neurologic defects without macrocytic anemia.
- 6. Serum methylmalonic acid and homocysteine levels are also elevated in vitamin B12 deficiency states.
- 7. Follow-up testing for antibodies to intrinsic factor (IF) is recommended to identify this potential cause of vitamin B12 malabsorption. NOTE: A normal serum concentration of vitamin B12 does not rule out tissue deficiency of vitamin B12. The most sensitive test for vitamin B12 deficiency at the cellular level is the assay for MMA. If clinical symptoms suggest deficiency, measurement of MMA and homocysteine should be considered, even if serum vitamin B12 concentrations are normal.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mr. GULSHAN AHUJA

**AGE/ GENDER** : 41 YRS/MALE **PATIENT ID** : 1629326

COLLECTED BY : REG. NO./LAB NO. : 012409300009

 REFERRED BY
 : 30/Sep/2024 08:27 AM

 BARCODE NO.
 : 01517982
 COLLECTION DATE
 : 30/Sep/2024 08:29 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 30/Sep/2024 09:02 AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

### CLINICAL PATHOLOGY

### **URINE ROUTINE & MICROSCOPIC EXAMINATION**

#### **PHYSICAL EXAMINATION**

QUANTITY RECIEVED	10	ml
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY		

COLOUR PALE YELLOW PALE YELLOW

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

TRANSPARANCY CLEAR CLEAR

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

SPECIFIC GRAVITY >=1.030 1.002 - 1.030

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

#### **CHEMICAL EXAMINATION**

REACTION ACIDIC

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

PROTEIN Trace NEGATIVE (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

SUGAR Negative NEGATIVE (-ve)
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

pH <=5.0 5.0 - 7.5

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

BILIRUBIN Negative NEGATIVE (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

NITRITE Negative NEGATIVE (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY.

UROBILINOGEN

Normal

EU/dL

0.2 - 1.0

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

KETONE BODIES

Negative

NEGATIVE (-ve)

ETONE BODIES Negative NEGATIVE (-ve) by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

BLOOD Negative NEGATIVE (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

ASCORBIC ACID NEGATIVE (-ve) NEGATIVE (-ve) by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

MICROSCOPIC EXAMINATION



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





CLIENT CODE.

### **KOS Diagnostic Lab**

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

: 30/Sep/2024 09:02AM

**NAME** : Mr. GULSHAN AHUJA

**AGE/ GENDER PATIENT ID** : 1629326 : 41 YRS/MALE

**COLLECTED BY** REG. NO./LAB NO. : 012409300009

REFERRED BY **REGISTRATION DATE** : 30/Sep/2024 08:27 AM **COLLECTION DATE** BARCODE NO. :01517982 : 30/Sep/2024 08:29AM

: KOS DIAGNOSTIC LAB **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Value	Unit	Biological Reference interval
NEGATIVE (-ve)	/HPF	0 - 3
2-4	/HPF	0 - 5
0-3	/HPF	ABSENT
NEGATIVE (-ve)		NEGATIVE (-ve)
ABSENT		ABSENT
	NEGATIVE (-ve)  2-4  0-3  NEGATIVE (-ve)  NEGATIVE (-ve)  NEGATIVE (-ve)  NEGATIVE (-ve)	NEGATIVE (-ve) /HPF  2-4 /HPF  0-3 /HPF  NEGATIVE (-ve)  NEGATIVE (-ve)  NEGATIVE (-ve)  NEGATIVE (-ve)

REPORTING DATE

End Of Report



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com