

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Miss. ANIKA
AGE/ GENDER : 22 YRS/FEMALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 01518671
CLIENT CODE. : KOS DIAGNOSTIC LAB
CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

PATIENT ID : 1139591
REG. NO./LAB NO. : 012410100057
REGISTRATION DATE : 10/Oct/2024 10:56 PM
COLLECTION DATE : 10/Oct/2024 10:58PM
REPORTING DATE : 10/Oct/2024 11:21PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

HAEMATOLOGY

PLATELET COUNT (P/C)

PLATELET COUNT (PLT)	430000	/cmm	150000 - 450000
----------------------	--------	------	-----------------

by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE & MICROSCOPY

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Miss. ANIKA	PATIENT ID	: 1139591
AGE/ GENDER	: 22 YRS/FEMALE	REG. NO./LAB NO.	: 012410100057
COLLECTED BY	:	REGISTRATION DATE	: 10/Oct/2024 10:56 PM
REFERRED BY	:	COLLECTION DATE	: 10/Oct/2024 10:58PM
BARCODE NO.	: 01518671	REPORTING DATE	: 10/Oct/2024 11:21PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

IMMUNOPATHOLOGY/SEROLOGY

TYPHOID COMBO SCREEN (TYPHOID ANTIGEN, IgG AND IgM): SERUM

TYPHOID ANTIGEN - SERUM <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)	NEGATIVE (-ve)
TYPHI DOT ANTIBODY IgG <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)	NEGATIVE (-ve)
TYPHI DOT ANTIBODY IgM <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)	NEGATIVE (-ve)

INTERPRETATION:

Typhoid fever is a life threatening illness caused by the bacterium *Salmonella typhus*. The infection is acquired typically by ingestion. On reaching the gut, the bacilli attach themselves to the epithelial cells of the intestinal villi and penetrate the lamina and submucosa. They are then phagocytosed there by polymorphs and mesenteric lymph nodes, where they multiply and, via the thoracic duct, enter the blood stream. A transient bacteremia follows, during which the bacilli are seeded in the liver, gall bladder, spleen, bone marrow, lymph nodes, and kidneys, where further multiplication takes place. Towards the end of the incubation period, there occurs a massive bacteremia from these sites, heralding the onset of the clinical symptoms.

The diagnosis of typhoid consists of isolation of the bacilli and the demonstration of antibodies. The isolation of the bacilli is very time consuming and antibody detection is not very specific. Other tests include the Widal reaction. The advantage of this test is that it takes only 10-20 minutes and requires only a small amount of stool/serum/plasma to perform. It is the easiest and most specific method for detecting *S. typhi* infection.

RELATIVE SENSITIVITY OF TYPHOID ANTIGEN DETECTION: 98.7%

RELATIVE SPECIFICITY OF TYPHOID ANTIGEN DETECTION: 97.4%

DETECTABLE IgM RESPONSE:

ONSET OF FEVER	PERCENT POSITIVE
4 - 6 DAYS	43.5
6 - 9 DAYS	92.9
> 9 DAYS	99.5

1. This is a solid phase, immunochromatographic ELISA assay that detects specific IgM and IgG Antibodies against the OUTER MEMBRANE PROTEIN (OMP) of the *Salmonella* species. IgM antibodies appear in the serum 2-3 days post infection and are indicative of a recent infection while the IgG antibodies appear later and are useful for presumptive diagnosis of Enteric fever if the patient presents more than a week after onset of symptoms.

2. This is a useful screening assay for the early detection of Enteric fever and has a high sensitivity. However the test has moderate specificity and false positive results may be obtained in the following situations:

Antibodies against *Salmonella* may cross react with other antibodies.




DR. VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR. YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Miss. ANIKA	PATIENT ID	: 1139591
AGE/ GENDER	: 22 YRS/FEMALE	REG. NO./LAB NO.	: 012410100057
COLLECTED BY	:	REGISTRATION DATE	: 10/Oct/2024 10:56 PM
REFERRED BY	:	COLLECTION DATE	: 10/Oct/2024 10:58PM
BARCODE NO.	: 01518671	REPORTING DATE	: 10/Oct/2024 11:21PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

. Unrelated infections may lead to production of specific Salmonella antibodies if the patient has previously been exposed to Salmonella infection (**ANAMNESTIC RESPONSE**).

NOTE:-Rapid blood culture performed during 1st week of infection is highly recommended for confirmation of all IgM positive results. In case the patient has presented after the first week of infection, a thorough clinical correlation and confirmatory Widal test must be performed to establish the diagnosis.




DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Miss. ANIKA	PATIENT ID	: 1139591
AGE/ GENDER	: 22 YRS/FEMALE	REG. NO./LAB NO.	: 012410100057
COLLECTED BY	:	REGISTRATION DATE	: 10/Oct/2024 10:56 PM
REFERRED BY	:	COLLECTION DATE	: 10/Oct/2024 10:58PM
BARCODE NO.	: 01518671	REPORTING DATE	: 10/Oct/2024 11:21PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
DENGUE FEVER COMBO SCREENING - (NS1 ANTIGEN, IgG AND IgM)			
DENGUE NS1 ANTIGEN - SCREENING <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)		NEGATIVE (-ve)
DENGUE ANTIBODY IgG - SCREENING <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)		NEGATIVE (-ve)
DENGUE ANTIBODY IgM - SCREENING <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)		NEGATIVE (-ve)

INTERPRETATION:-

- 1.This is a solid phase immunochromatographic ELISA test for the qualitative detection of the specific IgG and IgM antibodies against the Dengue virus.
- 2.The IgM antibodies take a minimum of 5-10 days in primary infection and 4-5 days in secondary infections to test positive and hence are suitable for the diagnosis of dengue fever only when the fever is approximately one week old.
- 3.The IgG antibodies develop at least two weeks after exposure to primary infection and subsequently remain positive for the rest of the life. A positive result is incapable of differentiating a current infection from a past infection.
- 4.The Dengue NS-1 antigen test is most suited for early diagnosis (within the first week of exposure).

*** End Of Report ***





DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)



DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS , MD (PATHOLOGY)

