

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	: Mr. BABLU	PATIENT ID	: 1646083
AGE/ GENDER	: 29 YRS/MALE	REG. NO./LAB NO.	: 012410170057
COLLECTED BY	:	REGISTRATION DATE	: 17/Oct/2024 02:33 PM
REFERRED BY	:	COLLECTION DATE	: 17/Oct/2024 02:34PM
BARCODE NO.	: 01519076	REPORTING DATE	: 17/Oct/2024 03:43PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

IMMUNOPATHOLOGY/SEROLOGY

DENGUE FEVER COMBO SCREENING - (NS1 ANTIGEN, IgG AND IgM)

DENGUE NS1 ANTIGEN - SCREENING <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)		NEGATIVE (-ve)
DENGUE ANTIBODY IgG - SCREENING <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)		NEGATIVE (-ve)
DENGUE ANTIBODY IgM - SCREENING <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)		NEGATIVE (-ve)

INTERPRETATION:-

- This is a solid phase immunochromatographic ELISA test for the qualitative detection of the specific IgG and IgM antibodies against the Dengue virus.
- The IgM antibodies take a minimum of 5-10 days in primary infection and 4-5 days in secondary infections to test positive and hence are suitable for the diagnosis of dengue fever only when the fever is approximately one week old.
- The IgG antibodies develop at least two weeks after exposure to primary infection and subsequently remain positive for the rest of the life. A positive result is incapable of differentiating a current infection from a past infection.
- The Dengue NS-1 antigen test is most suited for early diagnosis (within the first week of exposure).

*** End Of Report ***



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.