

KOS Diagnostic Lab

(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SUKHWINDER

AGE/ GENDER : 45 YRS/MALE **PATIENT ID** : 1458831

COLLECTED BY : REG. NO./LAB NO. : 012410190001

 REFERRED BY
 : 19/Oct/2024 05:38 AM

 BARCODE NO.
 : 01519146
 COLLECTION DATE
 : 19/Oct/2024 08:42AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 19/Oct/2024 10:25AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 0.869 ng/mL 0.35 - 1.93 by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROXINE (T4): SERUM 7.27 μgm/dL 4.87 - 12.60

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROID STIMULATING HORMONE (TSH): SERUM 5.225 μIU/mL 0.35 - 5.50 by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triiodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction (hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	Т3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.; phenytoin, salicylates).
- 3. Serum T4 levels in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (μg/dL)	Age	Reference Range (µIU/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 – 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





KOS Diagnostic Lab

(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SUKHWINDER

AGE/ GENDER : 45 YRS/MALE PATIENT ID : 1458831

COLLECTED BY : REG. NO./LAB NO. : 012410190001

 REFERRED BY
 : 19/Oct/2024 05:38 AM

 BARCODE NO.
 : 01519146
 COLLECTION DATE
 : 19/Oct/2024 08:42AM

CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** : 19/Oct/2024 10:25AM

CLIENT ADDRESS: 6349/1, NICHOLSON ROAD, AMBALA CANTT

6 - 12 Months 0.74 - 2.40 6 - 12 Months 7.10 - 16.16 6 - 12 Months 0.70 - 7.00 1 - 10 Years 0.92 - 2.28 1 - 10 Years 6.00 - 13.80 1 - 10 Years 0.60 - 5.50 11- 19 Years 0.35 - 1.93 11 - 19 Years 4.87 - 13.20 11 - 19 Years 0.50 - 5.50 > 20 years (Adults) 0.35 - 1.93 > 20 Years (Adults) 4.87 - 12.60 > 20 Years (Adults) 0.35 - 5.50 RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (μΙU/mL) 1st Trimester 0.10 - 2.50 2nd Trimester 0.20 - 3.00							
1 - 10 Years 0.92 - 2.28 1 - 10 Years 6.00 - 13.80 1 - 10 Years 0.60 - 5.50 11 - 19 Years 0.35 - 1.93 11 - 19 Years 4.87 - 13.20 11 - 19 Years 0.50 - 5.50 > 20 years (Adults) 0.35 - 1.93 > 20 Years (Adults) 4.87 - 12.60 > 20 Years (Adults) 0.35 - 5.50 RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (μU/mL) 1st Trimester 0.10 - 2.50 2nd Trimester 0.20 - 3.00	Test Name			Value	Unit		Biological Reference interva
11- 19 Years	6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00	
> 20 years (Adults)	1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (µIU/mL) 1st Trimester 0.10 – 2.50 2nd Trimester 0.20 – 3.00	11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50	
1st Trimester 0.10 - 2.50 2nd Trimester 0.20 - 3.00	> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35- 5.50	
2nd Trimester 0.20 – 3.00		RECOM	MENDATIONS OF TSH L	EVELS DURING PRE	GNANCY (µIU/mL)		
3.27 3.00	1st Trimester			0.10 - 2.50			
3rd Trimester 0.30 – 4.10	2nd Trimester			0.20 - 3.00			
0.00	3rd Trimester			0.30 - 4.10			

INCREASED TSH LEVELS:

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

- 1.Toxic multi-nodular goiter & Thyroiditis.
- $2. Over \ replacement \ of \ thyroid \ hormone \ in \ treatment \ of \ hypothyroid ism.$
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituitary or hypothalamic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester

*** End Of Report **



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana