



	Dr. Vinay ChopraDr. Yugam ChopraMD (Pathology & Microbiology)MD (Pathology)Chairman & Consultant PathologistCEO & Consultant Pathologist			
NAME	: Mr. HARBANS LAL			
AGE/ GENDER	: 70 YRS/MALE		ENT ID	: 1647599
COLLECTED BY	:		NO./LAB NO.	: 012410190017
REFERRED BY	BY :		TRATION DATE : 19/Oct/2024 08:47 AM	
BARCODE NO.	: 01519162	COLLECTION DATE		: 19/Oct/2024 08:48AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		RTING DATE	: 19/Oct/2024 12:39PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, A	AMBALA CANTT		
Test Name		Value	Unit	Biological Reference interval
		CAL CHEMISTRY/ E FASTING (F) AND		
GLUCOSE FASTING (F): PLASMA by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)		137.83 ^H	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0
GLUCOSE POST PRANDIAL (PP): PLASMA by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)		220.81 ^H	mg/dL	NORMAL: < 140.00 PREDIABETIC: 140.0 - 200.0 DIABETIC: > 0R = 200.0
INTERPRETATION:				

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose below 100 mg/dL and post-prandial plasma glucose level below 140 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl and post-prandial plasma glucose level between 140 – 200 mg/dL is considered as glucose intolerant or pre diabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dL and post-prandial plasma glucose level above 200 mg/dL is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

*** End Of Report ***





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

