

## **KOS Diagnostic Lab**

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mrs. SHWETA KUMARI

AGE/ GENDER : 25 YRS/FEMALE **PATIENT ID** : 1649576

**COLLECTED BY** : SURJESH REG. NO./LAB NO. :012410210078

REFERRED BY **REGISTRATION DATE** : 21/Oct/2024 05:43 PM BARCODE NO. :01519324 **COLLECTION DATE** : 21/Oct/2024 05:52PM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 24/Oct/2024 05:38AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

### **MICROBIOLOGY**

#### CULTURE AEROBIC BACTERIA AND ANTIBIOTIC SENSITIVITY: URINE

#### **CULTURE AND SUSCEPTIBILITY: URINE**

DATE OF SAMPLE 22-10-2024 SPECIMEN SOURCE URINE INCUBATION PERIOD 48 HOURS by AUTOMATED BROTH CULTURE

**STERILE CULTURE** 

by AUTOMATED BROTH CULTURE

NO AEROBIC PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF **ORGANISM** by AUTOMATED BROTH CULTURE

**INCUBATION AT 37\*C** 

#### **AEROBIC SUSCEPTIBILITY: URINE**

#### INTERPRETATION:

1. In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection.

2. Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out"

catheterization or from patients with indwelling catheters. **SUSCEPTIBILITY:** 

1. A test interpreted as SENSTITIVE implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent

recommended for that type of infection and infecting species, unless otherwise indicated..

2. A test interpreted as **INTERMEDIATE** implies that the" Infection due to the isolate may be appropriately treated in body sites where the drugs are

physiologically concentrated or when a high dosage of drug can be used".

3.A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distributions are not inhibited by the usually achievable concentration of the agents with normal distribution and the properties of the agents with a second c dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

#### **CAUTION:**

Conditions which can cause a false Negative culture:

- 1. Patient is on antibiotics. Please repeat culture post therapy.
- 2. Anaerobic bacterial infection.
- 3. Fastidious aerobic bacteria which are not able to grow on routine culture media.4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.
- 5. Renal tuberculosis to be confirmed by AFB studies.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



# KOS Diagnostic Lab (A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mrs. SHWETA KUMARI

**AGE/ GENDER PATIENT ID** : 25 YRS/FEMALE : 1649576

**COLLECTED BY** : SURJESH : 012410210078 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 21/Oct/2024 05:43 PM **COLLECTION DATE** BARCODE NO. :01519324 : 21/Oct/2024 05:52PM REPORTING DATE CLIENT CODE. : KOS DIAGNOSTIC LAB : 24/Oct/2024 05:38AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Test Name Value** Unit **Biological Reference interval** 

End Of Report



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com