



	Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist	Dr. Yugam MD CEO & Consultant	(Pathology)
NAME	: Mr. HARKESH BHARDWAJ		
AGE/ GENDER	: 63 YRS/MALE	PATIENT ID	: 1649971
COLLECTED BY	: SURJESH	REG. NO./LAB NO.	: 012410220022
REFERRED BY	: CENTRAL PHOENIX CLUB (AMBALA CANTT)	REGISTRATION DATE	: 22/Oct/2024 09:35 AM
BARCODE NO.	: 01519348	COLLECTION DATE	: 22/Oct/2024 10:44AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPORTING DATE	: 24/Oct/2024 01:03PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		
Test Name	Value	Unit	Biological Reference interval
	MICROI CULTURE AEROBIC BACTERIA ANI	BIOLOGY D ANTIBIOTIC SENS	SITIVITY: URINE
CULTURE AND SU	SCEPTIBILITY: URINE		
DATE OF SAMPLE	22-10-202	24	
SPECIMEN SOURCE			

SPECIMEN SOURCE	URINE
INCUBATION PERIOD by Automated broth culture	48 HOURS
CULTURE by AUTOMATED BROTH CULTURE	STERILE
ORGANISM by AUTOMATED BROTH CULTURE	NO AEROBIC PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF INCUBATION AT 37*C
AEROBIC SUSCEPTIBILITY: URINE	

INTERPRETATION:

 In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection.
Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out" activity is a frame activity in the indicate infection. catheterization or from patients with indwelling catheters.

SUSCEPTIBILITY:

1. A test interpreted as **SENSTITIVE** implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent recommended for that type of infection and infecting species, unless otherwise indicated.. 2. A test interpreted as **INTERMEDIATE** implies that the" Infection due to the isolate may be appropriately treated in body sites where the drugs are

physiologically concentrated or when a high dosage of drug can be used". 3.A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal

dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

CAUTION:

Conditions which can cause a false Negative culture:

1. Patient is on antibiotics. Please repeat culture post therapy.

2. Anaerobic bacterial infection.

3. Fastidious aerobic bacteria which are not able to grow on routine culture media.

4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.

5. Renal tuberculosis to be confirmed by AFB studies.

*** End Of Report ***





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana 0171-2643898, +91 99910 43898 care@koshealthcare.com www.koshealthcare.com



TEST PERFORMED AT KOS DIAGNOSTIC LAB. AMBALA CANTT