

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.



		hopra & Microbiology) onsultant Pathologist	Dr. Yugam MD CEO & Consultant	(Pathology)
JAME	: Mr. ANIL KUMAR			
AGE/ GENDER	: 49 YRS/MALE	PATI	ENT ID	: 1652882
COLLECTED BY	:	REG.	NO./LAB NO.	: 012410250007
REFERRED BY	:	REGI	STRATION DATE	: 25/Oct/2024 08:09 AM
BARCODE NO.	: 01519508	COLL	ECTION DATE	: 25/Oct/2024 08:21AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPO	RTING DATE	: 25/Oct/2024 10:58AM
LIENT ADDRESS	: 6349/1, NICHOLSON ROAD), AMBALA CANTT		
Fest Name		Value	Unit	Biological Reference interval
		GLUCOSE FAST	TING (F)	
	CLINI	CAL CHEMISTRY. GLUCOSE FAST		'nY
NTERPRETATION	E - PEROXIDASE (GOD-POD)	116.34 ^H	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0
N ACCORDANCE WIT	H AMERICAN DIABETES ASSOCIA	TION GUIDELINES:		
N ACCORDANCE WIT . A fasting plasma g . A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa	tients.	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for al atory for diabetic state.
N ACCORDANCE WIT . A fasting plasma g . A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa	tients. iabetic state. A repe occasions is confirm	
N ACCORDANCE WIT . A fasting plasma g 2. A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa l is highly suggestive of d ess of 125 mg/dl on both	tients. iabetic state. A repe occasions is confirm	
N ACCORDANCE WIT . A fasting plasma g . A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa l is highly suggestive of d ess of 125 mg/dl on both	tients. iabetic state. A repe occasions is confirm	
N ACCORDANCE WIT . A fasting plasma g 2. A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa l is highly suggestive of d ess of 125 mg/dl on both	tients. iabetic state. A repe occasions is confirm	
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa l is highly suggestive of d ess of 125 mg/dl on both	tients. iabetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for al atory for diabetic state.
N ACCORDANCE WIT . A fasting plasma g 2. A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa l is highly suggestive of d ess of 125 mg/dl on both	tients. iabetic state. A repe occasions is confirm	
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa l is highly suggestive of d ess of 125 mg/dl on both	tients. iabetic state. A repe occasions is confirm	
N ACCORDANCE WIT . A fasting plasma g . A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa l is highly suggestive of d ess of 125 mg/dl on both	tients. iabetic state. A repe occasions is confirm	
N ACCORDANCE WIT . A fasting plasma g . A fasting plasma g est (after consumpt	plucose level below 100 mg/dl is plucose level between 100 - 125 jon of 75 gms of glucose) is recc	s considered normal. mg/dl is considered as g ommended for all such pa l is highly suggestive of d ess of 125 mg/dl on both	tients. iabetic state. A repe occasions is confirm	

KOS Diagnostic Lab (A Unit of KOS Healthcare)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

 KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana

 KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana

 0171-2643898, +91 99910 43898
 care@koshealthcare.com
 www.koshealthcare.com



Page 1 of 1