

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.



		<b>Chopra</b> ogy & Microbiology) Consultant Pathologis		(Pathology)
AME	: Mr. LAXMAN SAINI			
GE/ GENDER	: 50 YRS/MALE		PATIENT ID	: 1644654
OLLECTED BY	:		REG. NO./LAB NO.	: 012410280002
EFERRED BY	:		<b>REGISTRATION DATE</b>	: 28/Oct/2024 06:00 AM
ARCODE NO.	:01519665		COLLECTION DATE	: 28/Oct/2024 06:02AM
LIENT CODE.	: KOS DIAGNOSTIC LAB		REPORTING DATE	: 28/Oct/2024 11:23AM
LIENT ADDRESS	: 6349/1, NICHOLSON RC	AD, AMBALA CANTI		
'est Name		Value	Unit	<b>Biological Reference interval</b>
by GLUCOSE OXIDAS I <u>TERPRETATION</u> ACCORDANCE WIT	E - PEROXIDASE (GOD-POD) H AMERICAN DIABETES ASSO	124.72 <sup>H</sup> CIATION GUIDELINES Il is considered norm	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0
by GLUCOSE OXIDAS <u>ITERPRETATION</u> I ACCORDANCE WIT A fasting plasma g st (after consumpt	E - PEROXIDASE (GOD-POD) H AMERICAN DIABETES ASSO lucose level below 100 mg/c lucose level between 100 - 1 ion of 75 gms of glucose) is r	CIATION GUIDELINES Il is considered norm 25 mg/dl is consider ecommended for all	al. ed as glucose intolerant or such patients	PREDIABETIC: 100.0 - 125.0
VTERPRETATION NACCORDANCE WIT A fasting plasma g A fasting plasma g	E - PEROXIDASE (GOD-POD) H AMERICAN DIABETES ASSO lucose level below 100 mg/c lucose level between 100 - 1 ion of 75 gms of glucose) is r	CIATION GUIDELINES dl is considered norm 25 mg/dl is consider ecommended for all g/dl is highly suggest xcess of 125 mg/dl o	al. ed as glucose intolerant or such patients. ve of diabetic state. A repe n both occasions is confirm	PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0 prediabetic. A fasting and post-prandial blood
by GLUCOSE OXIDAS <u>ITERPRETATION</u> I ACCORDANCE WIT A fasting plasma g st (after consumpt	E - PEROXIDASE (GOD-POD) H AMERICAN DIABETES ASSO lucose level below 100 mg/c lucose level between 100 - 1 ion of 75 gms of glucose) is r	CIATION GUIDELINES Il is considered norm 25 mg/dl is consider ecommended for all	al. ed as glucose intolerant or such patients. ve of diabetic state. A repe n both occasions is confirm	PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0 prediabetic. A fasting and post-prandial blood
by GLUCOSE OXIDAS ITERPRETATION I ACCORDANCE WIT A fasting plasma g st (after consumpt	E - PEROXIDASE (GOD-POD) H AMERICAN DIABETES ASSO lucose level below 100 mg/c lucose level between 100 - 1 ion of 75 gms of glucose) is r	CIATION GUIDELINES dl is considered norm 25 mg/dl is consider ecommended for all g/dl is highly suggest xcess of 125 mg/dl o	al. ed as glucose intolerant or such patients. ve of diabetic state. A repe n both occasions is confirm	PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0 prediabetic. A fasting and post-prandial blood
by GLUCOSE OXIDAS ITERPRETATION I ACCORDANCE WIT A fasting plasma g st (after consumpt	E - PEROXIDASE (GOD-POD) H AMERICAN DIABETES ASSO lucose level below 100 mg/c lucose level between 100 - 1 ion of 75 gms of glucose) is r	CIATION GUIDELINES dl is considered norm 25 mg/dl is consider ecommended for all g/dl is highly suggest xcess of 125 mg/dl o	al. ed as glucose intolerant or such patients. ve of diabetic state. A repe n both occasions is confirm	PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0 prediabetic. A fasting and post-prandial blood
by GLUCOSE OXIDAS <u>ITERPRETATION</u> I ACCORDANCE WIT A fasting plasma g st (after consumpt	E - PEROXIDASE (GOD-POD) H AMERICAN DIABETES ASSO lucose level below 100 mg/c lucose level between 100 - 1 ion of 75 gms of glucose) is r	CIATION GUIDELINES dl is considered norm 25 mg/dl is consider ecommended for all g/dl is highly suggest xcess of 125 mg/dl o	al. ed as glucose intolerant or such patients. ve of diabetic state. A repe n both occasions is confirm	PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0 prediabetic. A fasting and post-prandial blood

KOS Diagnostic Lab (A Unit of KOS Healthcare)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

V DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

