NAME	**: Mr. DEEPAK RISHI**		
AGE/ GENDER	: 64 YRS/MALE	**PATIENT ID**	: 1659887
COLLECTED BY	: SURJESH	**REG. NO./LAB NO.**	**: 012411040032**
REFERRED BY	: CENTRAL PHOENIX CLUB (AMBALA CANTT)	**REGISTRATION DATE**	: 04/Nov/2024 10:19 AM
BARCODE NO.	: 01520037	**COLLECTION DATE**	: 04/Nov/2024 10:21AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 04/Nov/2024 11:29AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

CLINICAL CHEMISTRY/BIOCHEMISTRY

GLUCOSE FASTING (F)

GLUCOSE FASTING (F): PLASMA *by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)*	**190.5^{H}**	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0

INTERPRETATION

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.
2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mr. DEEPAK RISHI**		
AGE/ GENDER	: 64 YRS/MALE	**PATIENT ID**	: 1659887
COLLECTED BY	: SURJESH	**REG. NO./LAB NO.**	**: 012411040032**
REFERRED BY	: CENTRAL PHOENIX CLUB (AMBALA CANTT)	**REGISTRATION DATE**	: 04/Nov/2024 10:19 AM
BARCODE NO.	: 01520037	**COLLECTION DATE**	: 04/Nov/2024 10:21AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 04/Nov/2024 11:19AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
CREATININE			
CREATININE: SERUM *by ENZYMATIC, SPECTROPHOTOMETRY*	**2.01^{H}**	mg/dL	0.40 - 1.40

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mr. DEEPAK RISHI**		
AGE/ GENDER	: 64 YRS/MALE	**PATIENT ID**	: 1659887
COLLECTED BY	: SURJESH	**REG. NO./LAB NO.**	**: 012411040032**
REFERRED BY	: CENTRAL PHOENIX CLUB (AMBALA CANTT)	**REGISTRATION DATE**	: 04/Nov/2024 10:19 AM
BARCODE NO.	: 01520037	**COLLECTION DATE**	: 04/Nov/2024 10:21AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 04/Nov/2024 11:19AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

GLOMERULAR FILTERATION RATE (GFR) - ESTIMATED

ESTIMATED GLOMERULAR FILTERATION RATE (eGFR): SERUM	**33.6[L]**	mL/min/1.73m2	KIDNEY FAILURE: < 15.0

by SPECTROPHOTOMETRY-ENZYMATIC, MDRD CALCULATION

INTERPRETATION:

CKD STAGE	DESCRIPTION	GFR (mL/min/1.73m2)	ASSOCIATED FINDINGS
G1	Normal kidney function	>90	No proteinuria
G2	Kidney damage with normal or high GFR	>90	Presence of Protein , Albumin or cast in urine
G3a	Mild decrease in GFR	60 -89	
G3b	Moderate decrease in GFR	30-59	
G4	Severe decrease in GFR	15-29	
G5	Kidney failure	<15	

COMMENTS:
1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.
2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012
3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD
4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

ADVICE:
KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)