

Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mr. JITENDER	PATIENT ID	: 1660598
AGE/ GENDER	: 58 YRS/MALE	REG. NO./LAB NO.	: 012411040079
COLLECTED BY	:	REGISTRATION DATE	: 04/Nov/2024 03:47 PM
REFERRED BY	:	COLLECTION DATE	: 04/Nov/2024 04:17PM
BARCODE NO.	: 01520084	REPORTING DATE	: 04/Nov/2024 05:46PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

CLINICAL CHEMISTRY/BIOCHEMISTRY
CREATININE PHOSPHOKINASE (CPK-NAC) TOTAL

CREATININE PHOSPHO KINASE (CPK-NAC)	128.5	IU/L	24 - 190
by SPECTROPHOTOMETRY			

Interpretation:-

- 1.Serum creatine kinase (CK) activity is greatly elevated, at some time during the course of the disease, in all types of muscular dystrophy, and especially so in Duchenne type, in which levels up to 50 times the upper limit of normal may be encountered.
- 2.In progressive muscular dystrophy, enzyme activity in serum is highest in infancy and childhood (7-10 years of age) and may be elevated long before the disease is clinically apparent.
- 3.Quite high values of CK are noted in viral myositis, polymyositis, and similar muscle diseases.
- 4.However, in neurogenic Parkinsonism, serum enzyme activity is normal. Very high activity is also encountered in malignant hyperthermia.

Significance:-

- 1.An early rise in CK is also seen after an acute MI, with values peaking at 12 to 24 hours and falling back to normal in 3 to 4 days.
- 2.Although total CK activity has been used as a diagnostic test for MI, it has been replaced by the troponin T and I immunoassays, and is no longer the laboratory test choice for diagnosing and monitoring acute infarctions.
- 3.Serum CK activity may increase in patients with acute cerebrovascular disease or neurosurgical intervention and with cerebral ischemia.
- 4.Serum CK activity also demonstrates an inverse relationship with thyroid activity. About 60% of hypothyroid subjects show an average elevation of CK activity 5-fold over the upper reference limit; elevation of as high as 50-fold may also be found.

Note: Exercise and muscle trauma (contact sports, traffic accidents, intramuscular injections, surgery, convulsions, wasp or bee stings, and burns) can elevate serum creatine kinase values.

*** End Of Report ***




DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)

