

Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Miss. DIYA	PATIENT ID	: 1555271
AGE/ GENDER	: 11 YRS/FEMALE	REG. NO./LAB NO.	: 012411050089
COLLECTED BY	:	REGISTRATION DATE	: 05/Nov/2024 06:52 PM
REFERRED BY	:	COLLECTION DATE	: 05/Nov/2024 06:56PM
BARCODE NO.	: 01520185	REPORTING DATE	: 05/Nov/2024 07:26PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

HAEMATOLOGY

HAEMOGLOBIN (HB)

HAEMOGLOBIN (HB) by CALORIMETRIC	11.1 ^L	gm / dL	12.0 - 16.0
-------------------------------------	-------------------	---------	-------------

INTERPRETATION:-

Hemoglobin is the protein molecule in red blood cells that carries oxygen from the lungs to the bodys tissues and returns carbon dioxide from the tissues back to the lungs.

A low hemoglobin level is referred to as ANEMIA or low red blood count.

ANEMIA (DECREASED HAEMOGLOBIN):

- 1) Loss of blood (traumatic injury, surgery, bleeding, colon cancer or stomach ulcer)
- 2) Nutritional deficiency (iron, vitamin B12, folate)
- 3) Bone marrow problems (replacement of bone marrow by cancer)
- 4) Suppression by red blood cell synthesis by chemotherapy drugs
- 5) Kidney failure
- 6) Abnormal hemoglobin structure (sickle cell anemia or thalassemia).

POLYCYTHEMIA (INCREASED HAEMOGLOBIN):

- 1) People in higher altitudes (Physiological)
- 2) Smoking (Secondary Polycythemia)
- 3) Dehydration produces a falsely rise in hemoglobin due to increased haemoconcentration
- 4) Advanced lung disease (for example, emphysema)
- 5) Certain tumors
- 6) A disorder of the bone marrow known as polycythemia rubra vera,
- 7) Abuse of the drug erythropoetin (Epogen) by athletes for blood doping purposes (increasing the amount of oxygen available to the body by chemically raising the production of red blood cells).

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD




 DR.VINAY CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST
 MBBS , MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Miss. DIYA	PATIENT ID	: 1555271
AGE/ GENDER	: 11 YRS/FEMALE	REG. NO./LAB NO.	: 012411050089
COLLECTED BY	:	REGISTRATION DATE	: 05/Nov/2024 06:52 PM
REFERRED BY	:	COLLECTION DATE	: 05/Nov/2024 06:56PM
BARCODE NO.	: 01520185	REPORTING DATE	: 06/Nov/2024 10:54AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

IMMUNOPATHOLOGY/SEROLOGY

ANTI TISSUE TRANSGLUTAMINASE (tTG) ANTIBODY IgA

ANTI TISSUE TRANSGLUTAMINASE ANTIBODY IgA	11.05	IU/mL	NEGATIVE: < 20.0 POSITIVE: > 20.0
---	-------	-------	--------------------------------------

by ELISA (ENZYME LINKED IMMUNOASSAY)

INTERPRETATION:

1. Anti-transglutaminase antibodies (ATA) are autoantibodies against the transglutaminase protein.
2. Antibodies to tissue transglutaminase are found in patients with several conditions, including coeliac disease, juvenile diabetes, inflammatory bowel disease, and various forms of arthritis.
3. In coeliac disease, ATA are involved in the destruction of the villous extracellular matrix and target the destruction of intestinal villous epithelial cells by killer cells.
4. Deposits of anti-tTG in the intestinal epithelium predict coeliac disease.
5. Celiac disease (gluten-sensitive enteropathy, celiac sprue) results from an immune-mediated inflammatory process following ingestion of wheat, rye, or barley proteins that occurs in genetically susceptible individuals. The inflammation in celiac disease occurs primarily in the mucosa of the small intestine, which leads to villous atrophy.

CLINICAL MANIFESTATIONS RELATED TO GASTROINTESTINAL TRACT:

1. Abdominal pain
2. Malabsorption
3. Diarrhea and Constipation.

CLINICAL MANIFESTATION OF CELIAC DISEASE NOT RESTRICTED TO GIT:

1. Failure to grow (delayed puberty and short stature)
2. Iron deficiency anemia
3. Recurrent fetal loss
4. Osteoporosis and chronic fatigue
5. Recurrent aphthous stomatitis (canker sores)
6. Dental enamel hypoplasia, and dermatitis herpetiformis.
7. Patients with celiac disease may also present with neuropsychiatric manifestations including ataxia and peripheral neuropathy, and are at increased risk for development of non-Hodgkin lymphoma.
8. The disease is also associated with other clinical disorders including thyroiditis, type I diabetes mellitus, Down syndrome, and IgA deficiency.

NOTE:

1. The finding of tissue transglutaminase (tTG)-IgA antibodies is specific for celiac disease and possibly for dermatitis herpetiformis. For individuals with moderately to strongly positive results, a diagnosis of celiac disease is likely and the patient should undergo biopsy to confirm the diagnosis.
2. If patients strictly adhere to a gluten-free diet, the unit value of IgA-anti-tTG should begin to decrease within 6 to 12 months of onset of dietary therapy.

CAUTION:

1. This test should not be solely relied upon to establish a diagnosis of celiac disease. It should be used to identify patients who have an increased probability of having celiac disease and in whom a small intestinal biopsy is recommended.




 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Miss. DIYA	PATIENT ID	: 1555271
AGE/ GENDER	: 11 YRS/FEMALE	REG. NO./LAB NO.	: 012411050089
COLLECTED BY	:	REGISTRATION DATE	: 05/Nov/2024 06:52 PM
REFERRED BY	:	COLLECTION DATE	: 05/Nov/2024 06:56PM
BARCODE NO.	: 01520185	REPORTING DATE	: 06/Nov/2024 10:54AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

2.Affected individuals who have been on a gluten-free diet prior to testing may have a negative result.
 3.For individuals who test negative, IgA deficiency should be considered. If total IgA is normal and tissue transglutaminase (tTG)-IgA is negative there is a low probability of the patient having celiac disease and a biopsy may not be necessary.
 4.If serology is negative or there is substantial clinical doubt remaining, then further investigation should be performed with endoscopy and bowel biopsy. This is especially important in patients with frank malabsorptive symptoms since many syndromes can mimic celiac disease. For the patient with frank malabsorptive symptoms, bowel biopsy should be performed regardless of serologic test results.
 5.The antibody pattern in dermatitis herpetiformis may be more variable than in celiac disease; therefore, both endomysial and tTG antibody determinations are recommended to maximize the sensitivity of the serologic tests.

*** End Of Report ***




DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)

