

Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. JASHANDEEP KAUR	PATIENT ID	: 1669712
AGE/ GENDER	: 34 YRS/FEMALE	REG. NO./LAB NO.	: 012411120055
COLLECTED BY	:	REGISTRATION DATE	: 12/Nov/2024 02:18 PM
REFERRED BY	:	COLLECTION DATE	: 12/Nov/2024 02:19PM
BARCODE NO.	: 01520674	REPORTING DATE	: 12/Nov/2024 04:53PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	0.788	ng/mL	0.35 - 1.93
THYROXINE (T4): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	8.91	µg/dL	4.87 - 12.60
THYROID STIMULATING HORMONE (TSH): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	3.712	µIU/mL	0.35 - 5.50

3rd GENERATION, ULTRA SENSITIVE

INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triiodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction (hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced


LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
2. Normal levels of T4 can also be seen in Hyperthyroid patients with : T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin, salicylates).
3. Serum T4 levels in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	Reference Range (ng/mL)	Age	Reference Range (µg/dL)	Age	Reference Range (µIU/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days - 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 - 12 Months	0.70 - 7.00




 DR. VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR. YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. JASHANDEEP KAUR	PATIENT ID	: 1669712
AGE/ GENDER	: 34 YRS/FEMALE	REG. NO./LAB NO.	: 012411120055
COLLECTED BY	:	REGISTRATION DATE	: 12/Nov/2024 02:18 PM
REFERRED BY	:	COLLECTION DATE	: 12/Nov/2024 02:19PM
BARCODE NO.	: 01520674	REPORTING DATE	: 12/Nov/2024 04:53PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60
RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (μ U/mL)			
1st Trimester			0.10 - 2.50
2nd Trimester			0.20 - 3.00
3rd Trimester			0.30 - 4.10

INCREASED TSH LEVELS:

- 1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2.Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3.Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.
- 5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

- 1.Toxic multi-nodular goiter & Thyroiditis.
- 2.Over replacement of thyroid hormone in treatment of hypothyroidism.
- 3.Autonomously functioning Thyroid adenoma
- 4.Secondary pituitary or hypothalamic hypothyroidism
- 5.Acute psychiatric illness
- 6.Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.
- 8.Pregnancy: 1st and 2nd Trimester




DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. JASHANDEEP KAUR	PATIENT ID	: 1669712
AGE/ GENDER	: 34 YRS/FEMALE	REG. NO./LAB NO.	: 012411120055
COLLECTED BY	:	REGISTRATION DATE	: 12/Nov/2024 02:18 PM
REFERRED BY	:	COLLECTION DATE	: 12/Nov/2024 02:19PM
BARCODE NO.	: 01520674	REPORTING DATE	: 12/Nov/2024 04:53PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

LUTEINISING HORMONE (LH)

LUTEINISING HORMONE (LH): SERUM	4.44	mIU/mL	MALES: 0.57 - 12.07 FOLLICULAR PHASE: 1.80 - 11.78 MID-CYCLE PEAK: 7.59 - 89.08 LUTEAL PHASE: 0.56 - 14.0 POST MENOPAUSAL WITHOUT HRT: 5.16 - 61.99
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			

INTERPRETATION:

1. Luteinizing hormone (LH) is a glycoprotein hormone consisting of 2 non covalently bound subunits (alpha and beta). Gonadotropin-releasing hormone from the hypothalamus controls the secretion of the gonadotropins, FSH and LH, from the anterior pituitary.
2. In both males and females, LH is essential for reproduction. In females, the menstrual cycle is divided by a mid cycle surge of both LH and FSH into a follicular phase and a luteal phase.
3. This "LH surge" triggers ovulation thereby not only releasing the egg, but also initiating the conversion of the residual follicle into a corpus luteum that, in turn, produces progesterone to prepare the endometrium for a possible implantation.
4. LH supports thecal cells in the ovary that provide androgens and hormonal precursors for estradiol production. LH in males acts on testicular interstitial cells of Leydig to cause increased synthesis of testosterone.

The test is useful in the following situations:

1. An adjunct in the evaluation of menstrual irregularities.
2. Evaluating patients with suspected hypogonadism
3. Predicting ovulation & Evaluating infertility
4. Diagnosing pituitary disorders
5. In both males and females, primary hypogonadism results in an elevation of basal follicle-stimulating hormone and luteinizing hormone levels.

FSH AND LH ELEVATED IN:

1. Primary gonadal failure
2. Complete testicular feminization syndrome
3. Precocious puberty (either idiopathic or secondary to a central nervous system lesion)
4. Menopause
5. Primary ovarian hypo dysfunction in females
6. Polycystic ovary disease in females
7. Primary hypogonadism in males

LH IS DECREASED IN:

1. Primary ovarian hyper function in females
2. Primary hypergonadism in males

NOTE

1. FSH and LH are both decreased in failure of the pituitary or hypothalamus.




 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. JASHANDEEP KAUR	PATIENT ID	: 1669712
AGE/ GENDER	: 34 YRS/FEMALE	REG. NO./LAB NO.	: 012411120055
COLLECTED BY	:	REGISTRATION DATE	: 12/Nov/2024 02:18 PM
REFERRED BY	:	COLLECTION DATE	: 12/Nov/2024 02:19PM
BARCODE NO.	: 01520674	REPORTING DATE	: 12/Nov/2024 05:26PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

FOLLICLE STIMULATING HORMONE (FSH)

FOLLICLE STIMULATING HORMONE (FSH): SERUM	2.68	mIU/mL	FEMALE FOLLICULAR PHASE: 2.5 - 11.4
by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)			FEMALE MID-CYCLE PEAK: 3.3 - 21.7
			FEAMLE LUTEAL PHASE: 1.2 - 7.0
			FEMALE POST-MENOPAUSAL: 18.8 - 132
			MALE: 1.0 - 12.1

INTERPRETATION:

1. Gonadotropin-releasing hormone from the hypothalamus controls the secretion of the gonadotropins, follicle-stimulating hormone (FSH) and luteinizing hormone (LH) from the anterior pituitary.
2. The menstrual cycle is divided by a midcycle surge of both FSH and LH into a follicular phase and a luteal phase.
3. FSH appears to control gametogenesis in both males and females.

The test is useful in the following settings:

1. An adjunct in the evaluation of menstrual irregularities.
2. Evaluating patients with suspected hypogonadism.
3. Predicting ovulation
4. Evaluating infertility
5. Diagnosing pituitary disorders
6. In both males and females, primary hypogonadism results in an elevation of basal follicle-stimulating hormone (FSH) and luteinizing hormone (LH) levels.

FSH and LH LEVELS ELEVATED IN:

1. Primary gonadal failure
2. Complete testicular feminization syndrome.
3. Precocious puberty (either idiopathic or secondary to a central nervous system lesion)
4. Menopause (postmenopausal FSH levels are generally >40 IU/L)
5. Primary ovarian hypofunction in females
6. Primary hypogonadism in males

NOTE:

1. Normal or decreased FSH is seen in polycystic ovarian disease in females
2. FSH and LH are both decreased in failure of the pituitary or hypothalamus.




 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. JASHANDEEP KAUR	PATIENT ID	: 1669712
AGE/ GENDER	: 34 YRS/FEMALE	REG. NO./LAB NO.	: 012411120055
COLLECTED BY	:	REGISTRATION DATE	: 12/Nov/2024 02:18 PM
REFERRED BY	:	COLLECTION DATE	: 12/Nov/2024 02:19PM
BARCODE NO.	: 01520674	REPORTING DATE	: 12/Nov/2024 04:53PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

PROLACTIN

PROLACTIN: SERUM	24.97	ng/mL	3 - 25
------------------	-------	-------	--------

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

INTERPRETATION:

1. Prolactin is secreted by the anterior pituitary gland and controlled by the hypothalamus.
 2. The major chemical controlling prolactin secretion is dopamine, which inhibits prolactin secretion from the pituitary.
 3. Physiological function of prolactin is the stimulation of milk production. In normal individuals, the prolactin level rises in response to physiologic stimuli such as sleep, exercise, nipple stimulation, sexual intercourse, hypoglycemia, postpartum period, and also is elevated in the newborn infant.

INCREASED (HYPERPROLACTEMIA):

1. Prolactin-secreting pituitary adenoma (prolactinoma, which is 5 times more frequent in females than males).
 2. Functional and organic disease of the hypothalamus.
 3. Primary hypothyroidism.
 4. Section compression of the pituitary stalk.
 5. Chest wall lesions and renal failure.
 6. Ectopic tumors.
 7. DRUGS:- Anti-Dopaminergic drugs like antipsychotic drugs, anti-nausea/antiemetic drugs, Drugs that affect CNS serotonin metabolism, serotonin receptors, or serotonin reuptake (anti-depressants of all classes, ergot derivatives, some illegal drugs such as cannabis), Antihypertensive drugs, Opiates, High doses of estrogen or progesterone, anticonvulsants (valproic acid), anti-tuberculous medications (Isoniazid).

SIGNIFICANCE:

1. In loss of libido, galactorrhea, oligomenorrhea or amenorrhea, and infertility in premenopausal females.
 2. Loss of libido, impotence, infertility, and hypogonadism in males. Postmenopausal and premenopausal women, as well as men, can also suffer from decreased muscle mass and osteoporosis.
 3. In males, prolactin levels >13 ng/mL are indicative of hyperprolactinemia.
 4. In women, prolactin levels >27 ng/mL in the absence of pregnancy and postpartum lactation are indicative of hyperprolactinemia.
 5. Clear symptoms and signs of hyperprolactinemia are often absent in patients with serum prolactin levels <100 ng/mL.
 4. Mild to moderately increased levels of serum prolactin are not a reliable guide for determining whether a prolactin-producing pituitary adenoma is present, 5. Whereas levels >250 ng/mL are usually associated with a prolactin-secreting tumor.

CAUTION:

Prolactin values that exceed the reference values may be due to macroprolactin (prolactin bound to immunoglobulin). Macroprolactin should be evaluated if signs and symptoms of hyperprolactinemia are absent, or pituitary imaging studies are not informative.

*** End Of Report ***




DR. VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR. YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)

