

Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Dog COCO	PATIENT ID	: 1670237
AGE/ GENDER	: 4 YRS/Female	REG. NO./LAB NO.	: 012411120071
COLLECTED BY	:	REGISTRATION DATE	: 12/Nov/2024 06:37 PM
REFERRED BY	:	COLLECTION DATE	: 12/Nov/2024 06:40PM
BARCODE NO.	: 01520690	REPORTING DATE	: 12/Nov/2024 08:40PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

ENDOCRINOLOGY
PROGESTERONE

PROGESTERONE: SERUM <i>by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)</i>	0.47	ng/mL	FEMALE FOLLICULAR PHASE: 0.10 - 1.50 FEMALE OVULATORY PHASE: 0.40 - 3.00 FEMALE LUTEAL PHASE: 1.20 - 18.80 POST MENOPAUSAL: < 1.40 MALES: < 2.80
---	------	-------	---

INTERPRETATION:

EXPECTED VALUES OF PROGESTERONE DURING PREGNANCY	
	UNITS (ng/mL)
First trimester (0 - 12 Wweeks)	15.8 - 46.0
Second trimester (13 - 28 Wweeks)	15.6 - 74.0
Third trimester (29 - 40 Wweeks)	45.0 - 143.0
Post Menopausal	< 1.40

1. Progesterone is produced by the adrenal glands, corpus luteum, and placenta.
2. After ovulation, there is a significant rise in serum Progesterone levels as the corpus luteum begins To produce progesterone in increasing amounts. This causes changes in the uterus, preparing it for implantation of a fertilized egg. If implantation occurs, the trophoblast begins to secrete human chorionic gonadotropin, which maintains the corpus luteum and its secretion of progesterone. If there is no implantation, the corpus luteum degenerates and circulating progesterone levels decrease rapidly, reaching follicular phase levels about 4 days before the next menstrual period.

The test is indicated for:

1. Ascertaining whether ovulation occurred in a menstrual cycle
2. Evaluation of placental function in pregnancy
3. Workup of some patients with adrenal or testicular tumors

NOTE:

In patients receiving therapy with high biotin doses (ie, >5 mg/day), no specimen should be drawn until at least 8 hours after the last biotin administration.

*** End Of Report ***




 DR.VINAY CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)

