

Dr. Vinay Chopra  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

Dr. Yugam Chopra  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. AARTI	<b>PATIENT ID</b>	: 1671300
<b>AGE/ GENDER</b>	: 35 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012411130059
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 13/Nov/2024 05:08 PM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 13/Nov/2024 05:10PM
<b>BARCODE NO.</b>	: 01520751	<b>REPORTING DATE</b>	: 13/Nov/2024 05:29PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## HAEMATOTOLOGY

### ERYTHROCYTE SEDIMENTATION RATE (ESR)

ERYTHROCYTE SEDIMENTATION RATE (ESR)	<b>103<sup>H</sup></b>	mm/1st hr	0 - 20
--------------------------------------	------------------------	-----------	--------

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

#### INTERPRETATION:

1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

#### CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

#### NOTE:

1. ESR and C - reactive protein (C-RP) are both markers of inflammation.
2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
3. **CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.**
4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



  
 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. AARTI	<b>PATIENT ID</b>	: 1671300
<b>AGE/ GENDER</b>	: 35 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012411130059
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 13/Nov/2024 05:08 PM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 13/Nov/2024 05:10PM
<b>BARCODE NO.</b>	: 01520751	<b>REPORTING DATE</b>	: 13/Nov/2024 05:45PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## IMMUNOPATHOLOGY/SEROLOGY

### TYPHOID COMBO SCREEN (TYPHOID ANTIGEN, IgG AND IgM): SERUM

TYPHOID ANTIGEN - SERUM <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)	NEGATIVE (-ve)
TYPHI DOT ANTIBODY IgG <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)	NEGATIVE (-ve)
TYPHI DOT ANTIBODY IgM <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	WEAKLY POSITIVE (+ve)	NEGATIVE (-ve)

#### INTERPRETATION:

Typhoid fever is a life threatening illness caused by the bacterium *Salmonella typhi*. The infection is acquired typically by ingestion. On reaching the gut, the bacilli attach themselves to the epithelial cells of the intestinal villi and penetrate the lamina and submucosa. They are then phagocytosed there by polymorphs and mesenteric lymph nodes, where they multiply and, via the thoracic duct, enter the blood stream. A transient bacteremia follows, during which the bacilli are seeded in the liver, gall bladder, spleen, bone marrow, lymph nodes, and kidneys, where further multiplication takes place. Towards the end of the incubation period, there occurs a massive bacteremia from these sites, heralding the onset of the clinical symptoms.

The diagnosis of typhoid consists of isolation of the bacilli and the demonstration of antibodies. The isolation of the bacilli is very time consuming and antibody detection is not very specific. Other tests include the Widal reaction. The advantage of this test is that it takes only 10-20 minutes and requires only a small amount of stool/serum/plasma to perform. It is the easiest and most specific method for detecting *S. typhi* infection.

**RELATIVE SENSITIVITY OF TYPHOID ANTIGEN DETECTION: 98.7%**

**RELATIVE SPECIFICITY OF TYPHOID ANTIGEN DETECTION: 97.4%**

#### DETECTABLE IgM RESPONSE:

ONSET OF FEVER	PERCENT POSITIVE
4 - 6 DAYS	43.5
6 - 9 DAYS	92.9
> 9 DAYS	99.5

1. This is a solid phase, immunochromatographic ELISA assay that detects specific IgM and IgG Antibodies against the OUTER MEMBRAN PROTEIN(OMP) of the *Salmonella* species. IgM antibodies appear in the serum 2-3 days post infection and are indicative of a recent infection while the IgG antibodies appear later and are useful for presumptive diagnosis of Enteric fever if the patient presents more than a week after onset of symptoms.

2. This is a useful screening assay for the early detection of Enteric fever and has a high sensitivity. However the test has moderate specificity and false positive results may be obtained in the following situations:

- Antibodies against *Salmonella* may cross react with other antibodies.
- Unrelated infections may lead to production of specific *Salmonella* antibodies if the patient has previously been exposed to



  
**DR. VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
**DR. YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. AARTI	<b>PATIENT ID</b>	: 1671300
<b>AGE/ GENDER</b>	: 35 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012411130059
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 13/Nov/2024 05:08 PM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 13/Nov/2024 05:10PM
<b>BARCODE NO.</b>	: 01520751	<b>REPORTING DATE</b>	: 13/Nov/2024 05:45PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

Salmonella infection (ANAMNESTIC RESPONSE).

NOTE:-Rapid blood culture performed during 1<sup>st</sup> week of infection is highly recommended for confirmation of all IgM positive results. In case the patient has presented after the first week of infection, a thorough clinical correlation and confirmatory Widal test must be performed to establish the diagnosis.



  
 DR.VINAY CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR.YUGAM CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. AARTI	<b>PATIENT ID</b>	: 1671300
<b>AGE/ GENDER</b>	: 35 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012411130059
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 13/Nov/2024 05:08 PM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 13/Nov/2024 05:10PM
<b>BARCODE NO.</b>	: 01520751	<b>REPORTING DATE</b>	: 13/Nov/2024 06:21PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

### CHIKUNGUNYA ANTIBODY IgM SCREENING

CHIKUNGUNYA ANTIBODY IgM QUANTITATIVE SERUM by ELISA (ENZYME LINKED IMMUNOASSAY)	0.36	INDEX VALUE	< 1.00
--	------	-------------	--------

#### INTERPRETATION:

- 1.Chikungunya is an insect borne viral disease belonging to genus ALPHAVIRIDAE transmitted to humans by infected Aedes mosquito.
- 2.It causes fever and severe joint pain. Other symptoms include muscle pain, headache, nausea, fatigue and rash.
- 3.Joint pain is often debilitating and can vary in duration.
- 4.The disease shares some clinical signs with dengue, and can be misdiagnosed in areas where dengue is common.
- 5.There is no cure for the disease. Treatment is focused on relieving the symptoms.

#### ROUTE OF TRANSMISSION:

- 1.The virus is transmitted from human to human by the bites of infected female mosquitoes.
- 2.Most commonly, the mosquitoes involved are Aedes aegypti and Aedes albopictus, two species which can also transmit other mosquito-borne viruses, including dengue.
- 3.These mosquitoes can be found biting throughout daylight hours, though there may be peaks of activity in the early morning and late afternoon.
- 4.Both species are found biting outdoors, but Ae. aegypti will also readily feed indoors.
- 5.After the bite of an infected mosquito, onset of illness occurs usually between 4 and 8 days but can range from 2 to 12 days.

#### NOTE:

This is a solid phase immunochromatographic assay for the detection of the Chikungunya specific IgM antibodies in the human serum. The test has a sensitivity of 97.5 % and a specificity of 99.1 %.



  
 DR.VINAY CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR.YUGAM CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)





Dr. Vinay Chopra  
MD (Pathology & Microbiology)  
Chairman & Consultant Pathologist

Dr. Yugam Chopra  
MD (Pathology)  
CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. AARTI	<b>PATIENT ID</b>	: 1671300
<b>AGE/ GENDER</b>	: 35 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012411130059
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 13/Nov/2024 05:08 PM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 13/Nov/2024 05:10PM
<b>BARCODE NO.</b>	: 01520751	<b>REPORTING DATE</b>	: 13/Nov/2024 05:45PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

**DENGUE FEVER COMBO SCREENING - (NS1 ANTIGEN, IgG AND IgM)**

DENGUE NS1 ANTIGEN - SCREENING <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)		NEGATIVE (-ve)
DENGUE ANTIBODY IgG - SCREENING <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)		NEGATIVE (-ve)
DENGUE ANTIBODY IgM - SCREENING <i>by ICT (IMMUNOCHROMATOGRAPHY)</i>	NEGATIVE (-ve)		NEGATIVE (-ve)

**INTERPRETATION:-**

- 1.This is a solid phase immunochromatographic ELISA test for the qualitative detection of the specific IgG and IgM antibodies against the Dengue virus.
- 2.The IgM antibodies take a minimum of 5-10 days in primary infection and 4-5 days in secondary infections to test positive and hence are suitable for the diagnosis of dengue fever only when the fever is approximately one week old.
- 3.The IgG antibodies develop at least two weeks after exposure to primary infection and subsequently remain positive for the rest of the life. A positive result is incapable of differentiating a current infection from a past infection.
- 4.The Dengue NS-1 antigen test is most suited for early diagnosis (within the first week of exposure).



  
 DR.VINAY CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR.YUGAM CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS , MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. AARTI	<b>PATIENT ID</b>	: 1671300
<b>AGE/ GENDER</b>	: 35 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012411130059
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 13/Nov/2024 05:08 PM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 13/Nov/2024 05:10PM
<b>BARCODE NO.</b>	: 01520751	<b>REPORTING DATE</b>	: 13/Nov/2024 06:24PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

**RHEUMATOID FACTOR (RA): QUANTITATIVE - SERUM**

RHEUMATOID (RA) FACTOR QUANTITATIVE: SERUM by NEPHLOMETRY	0.31	IU/mL	NEGATIVE: < 18.0 BORDERLINE: 18.0 - 25.0 POSITIVE: > 25.0
---	------	-------	---

**INTERPRETATION:-**

**RHEUMATOID FACTOR (RA):**

1. Rheumatoid factors (RF) are antibodies that are directed against the Fc fragment of IgG altered in its tertiary structure.
2. Over 75% of patients with rheumatoid arthritis (RA) have an IgM antibody to IgG immunoglobulin. This autoantibody (RF) is diagnostically useful although it may not be etiologically related to RA.
3. Inflammatory Markers such as ESR & C-Reactive protein (CRP) are normal in about 60 % of patients with positive RA.
4. The titer of RF correlates poorly with disease activity, but those patients with high titers tend to have more severe disease course.
5. The test is useful for diagnosis and prognosis of rheumatoid arthritis.

**RHEUMATOID ARTHRITIS:**

1. Rheumatoid Arthritis is a systemic autoimmune disease that is multi-functional in origin and is characterized by chronic inflammation of the membrane lining (synovium) joints which leads to progressive joint destruction and in most cases to disability and reduction of quality life.
2. The disease spreads from small to large joints, with greatest damage in early phase.
3. The diagnosis of RA is primarily based on clinical, radiological & immunological features. The most frequent serological test is the measurement of RA factor.

**CAUTION (FALSE POSTIVE):-**

1. RA factor is not specific for Rheumatoid arthritis, as it is often present in healthy individuals with other autoimmune diseases and chronic infections.
2. Non rheumatoid and rheumatoid arthritis (RA) populations are not clearly separate with regard to the presence of rheumatoid factor (RF) (15% of RA patients have a nonreactive titer and 8% of nonrheumatoid patients have a positive titer).
3. Patients with various nonrheumatoid diseases characterized by chronic inflammation may have positive tests for RF. These diseases include systemic lupus erythematosus, polymyositis, tuberculosis, syphilis, viral hepatitis, infectious mononucleosis, and influenza.
4. Anti-CCP have been discovered in joints of patients with RA, but not in other form of joint disease. Anti-CCP2 is HIGHLY SENSITIVE (71%) & more specific (98%) than RA factor.
5. Upto 30 % of patients with Seronegative Rheumatoid arthritis also show Anti-CCP antibodies.
6. The positive predictive value of Anti-CCP antibodies for Rheumatoid Arthritis is far greater than Rheumatoid factor.



  
 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

Dr. Yugam Chopra  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. AARTI	<b>PATIENT ID</b>	: 1671300
<b>AGE/ GENDER</b>	: 35 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012411130059
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 13/Nov/2024 05:08 PM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 13/Nov/2024 05:10PM
<b>BARCODE NO.</b>	: 01520751	<b>REPORTING DATE</b>	: 13/Nov/2024 06:44PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## CLINICAL PATHOLOGY

### URINE ROUTINE & MICROSCOPIC EXAMINATION

#### PHYSICAL EXAMINATION

QUANTITY RECEIVED	10	ml	
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
COLOUR	AMBER YELLOW		PALE YELLOW
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
TRANSPARANCY	HAZY		CLEAR
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
SPECIFIC GRAVITY	1.01		1.002 - 1.030
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			

#### CHEMICAL EXAMINATION


REACTION	ACIDIC		
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
PROTEIN	Negative		NEGATIVE (-ve)
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
SUGAR	Negative		NEGATIVE (-ve)
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
pH	<=5.0		5.0 - 7.5
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
BILIRUBIN	Negative		NEGATIVE (-ve)
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
NITRITE	Negative		NEGATIVE (-ve)
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
UROBILINOGEN	Normal	EU/dL	0.2 - 1.0
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
KETONE BODIES	Negative		NEGATIVE (-ve)
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
BLOOD	TRACE		NEGATIVE (-ve)
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
ASCORBIC ACID	NEGATIVE (-ve)		NEGATIVE (-ve)
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			

#### MICROSCOPIC EXAMINATION

RED BLOOD CELLS (RBCs)	0-2	/HPF	0 - 3
------------------------	-----	------	-------



  
 DR.VINAY CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR.YUGAM CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)



**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mrs. AARTI	<b>PATIENT ID</b>	: 1671300
<b>AGE/ GENDER</b>	: 35 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 012411130059
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 13/Nov/2024 05:08 PM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 13/Nov/2024 05:10PM
<b>BARCODE NO.</b>	: 01520751	<b>REPORTING DATE</b>	: 13/Nov/2024 06:44PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
PUS CELLS	10-15	/HPF	0 - 5
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
EPITHELIAL CELLS	6-8	/HPF	ABSENT
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
CRYSTALS	NEGATIVE (-ve)		NEGATIVE (-ve)
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
CASTS	NEGATIVE (-ve)		NEGATIVE (-ve)
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
BACTERIA	NEGATIVE (-ve)		NEGATIVE (-ve)
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
OTHERS	NEGATIVE (-ve)		NEGATIVE (-ve)
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
TRICHOMONAS VAGINALIS (PROTOZOA)	ABSENT		ABSENT
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			

\*\*\* End Of Report \*\*\*



  
 DR.VINAY CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
 DR.YUGAM CHOPRA  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)

