



	Dr. Vinay ChopraDr. Yugam ChopraMD (Pathology & Microbiology)MD (Pathology)Chairman & Consultant PathologistCEO & Consultant Pathologist			
NAME	: Mr. VED PARKASH			
AGE/ GENDER	: 63 YRS/MALE	PATIENT ID	: 1673596	
<b>COLLECTED BY</b>	:	REG. NO./LAB NO.	: 012411160051	
<b>REFERRED BY</b>	:	<b>REGISTRATION DATE</b>	: 16/Nov/2024 01:07 PM	
BARCODE NO.	: 01520927	COLLECTION DATE	: 16/Nov/2024 01:07PM	
CLIENT CODE.	: KOS DIAGNOSTIC LAB	<b>REPORTING DATE</b>	: 18/Nov/2024 06:40PM	
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, A	MBALA CANTT		
Test Name		Value Unit	<b>Biological Reference interval</b>	
	CULTURE AEROBIC B	MICROBIOLOGY ACTERIA AND ANTIBIOTIC SEN	SITIVITY: URINE	
CULTURE AND SUS	SCEPTIBILITY: URINE			
DATE OF SAMPLE		16-11-2024		
SPECIMEN SOURCE		URINE		
INCUBATION PERIOD by AUTOMATED BROTH CULTURE		48 HOURS		
CULTURE by AUTOMATED BROTH CULTURE		STERILE		
ORGANISM by AUTOMATED BROTH CULTURE		NO AEROBIC PYOGENIC ORGAN INCUBATION AT 37*C	NO AEROBIC PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF INCUBATION AT 37*C	
AEROBIC SUSCEPT	TIBILITY: URINE			

<u>INTERPRETATIONS</u> 1. In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection. 2. Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out" catheterization or from patients with indwelling catheters.

## SUSCEPTIBILITY:

TEST PERFORMED AT KOS DIAGNOSTIC LAB. AMBALA CANTT

 A test interpreted as SENSTITIVE implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent recommended for that type of infection and infecting species, unless otherwise indicated..
A test interpreted as INTERMEDIATE implies that the" Infection due to the isolate may be appropriately treated in body sites where the drugs are

2. A test interpreted as **INTERMEDIATE** implies that the" Infection due to the isolate may be appropriately treated in body sites where the drugs are physiologically concentrated or when a high dosage of drug can be used". 3.A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal

3.A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

## CAUTION:

Conditions which can cause a false Negative culture:

1. Patient is on antibiotics. Please repeat culture post therapy.

2. Anaerobic bacterial infection.

- 3. Fastidious aerobic bacteria which are not able to grow on routine culture media.
- 4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.
- 5. Renal tuberculosis to be confirmed by AFB studies.

\*\*\* End Of Report \*\*\*





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: Ilnd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 care@koshealthcare.com www.koshealthcare.com

