Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mr. BIRDEVINDER SINGH**		
AGE/ GENDER	: 31 YRS/MALE	**PATIENT ID**	: 1684359
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012411270068**
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	**REGISTRATION DATE**	: 27/Nov/2024 07:05 PM
BARCODE NO.	: 01521570	**COLLECTION DATE**	: 27/Nov/2024 07:08PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Nov/2024 07:44PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

HAEMATOLOGY

BLOOD GROUP (ABO) AND RH FACTOR TYPING

ABO GROUP *by SLIDE AGGLUTINATION*	O		
RH FACTOR TYPE *by SLIDE AGGLUTINATION*	POSITIVE		

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mr. BIRDEVINDER SINGH**		
AGE/ GENDER	: 31 YRS/MALE	**PATIENT ID**	: 1684359
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012411270068**
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	**REGISTRATION DATE**	: 27/Nov/2024 07:05 PM
BARCODE NO.	: 01521570	**COLLECTION DATE**	: 27/Nov/2024 07:08PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Nov/2024 10:50PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

IMMUNOPATHOLOGY/SEROLOGY

HEPATITIS C VIRUS (HCV) ANTIBODY: TOTAL

HEPATITIS C ANTIBODY (HCV) TOTAL: SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	0.06	S/CO	NEGATIVE: < 1.00 POSITIVE: > 1.00
HEPATITIS C ANTIBODY (HCV) TOTAL RESULT *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	NON - REACTIVE		

INTERPRETATION:-

RESULT (INDEX)	REMARKS
< 1.00	NON - REACTIVE/NOT - DETECTED
> =1.00	REACTIVE/ASYMPTOMATIC/INFECTIVE STATE/CARRIER STATE.

Hepatitis C (HCV) is an RNA virus of Favivirus group transmitted via blood transfusions, transplantation, injection drug abusers, accidental needle punctures in healthcare workers, dialysis patients and rarely from mother to infant. 10 % of new cases show sexual transmission. As compared to HAV & HBV , chronic infection with HCV occurs in 85 % of infected individuals. In high risk population, the predictive value of Anti HCV for HCV infection is > 99% whereas in low risk populations it is only 25 %.

USES:

1. Indicator of past or present infection, but does not differentiate between Acute/ Chronic/Resolved Infection.
2. Routine screening of low and high prevelance population including blood donors.

NOTE:

1. False positive results are seen in Auto-immune disease, Rheumatoid Factor, HYpergammaglobulinemia, Paraproteinemia, Passive antibody transfer, Anti-idiotypes and Anti-superoxide dismutase.
2. False negative results are seen in early Acute infection, Immunosuppression and Immuno—incompetence.
3. *HCV-RNA PCR recommended in all reactive results to differentiate between past and present infection.*

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mr. BIRDEVINDER SINGH**		
AGE/ GENDER	: 31 YRS/MALE	**PATIENT ID**	: 1684359
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012411270068**
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	**REGISTRATION DATE**	: 27/Nov/2024 07:05 PM
BARCODE NO.	: 01521570	**COLLECTION DATE**	: 27/Nov/2024 07:08PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Nov/2024 10:50PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

ANTI HUMAN IMMUNODEFICIENCY VIRUS (HIV) DUO ULTRA WITH (P-24 ANTIGEN DETECTION)

HIV 1/2 AND P24 ANTIGEN: SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	0.07	S/CO	NEGATIVE: < 1.00 POSITIVE: > 1.00
HIV 1/2 AND P24 ANTIGEN RESULT *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	NON - REACTIVE		

INTERPRETATION:-

RESULT (INDEX)	REMARKS
< 1.00	NON - REACTIVE
> = 1.00	PROVISIONALLY REACTIVE

Non-Reactive result implies that antibodies to HIV 1/ 2 have not been detected in the sample . This menas that patient has either not been exposed to HIV 1/ 2 infection or the sample has been tested during the “window phase” i.e. before the development of detectable levels of antibodies. Hence a Non Reactive result does not exclude the possibility of exposure or infection with HIV 1/ 2.

RECOMMENDATIONS:

1. Results to be clinically correlated
2. Rarely falsenegativity/positivity may occur.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mr. BIRDEVINDER SINGH**		
AGE/ GENDER	: 31 YRS/MALE	**PATIENT ID**	: 1684359
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012411270068**
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	**REGISTRATION DATE**	: 27/Nov/2024 07:05 PM
BARCODE NO.	: 01521570	**COLLECTION DATE**	: 27/Nov/2024 07:08PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Nov/2024 10:50PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

HEPATITIS B SURFACE ANTIGEN (HBsAg) ULTRA

HEPATITIS B SURFACE ANTIGEN (HBsAg): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	0.27	S/CO	NEGATIVE: < 1.0 POSITIVE: > 1.0
HEPATITIS B SURFACE ANTIGEN (HBsAg) RESULT *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	NON REACTIVE		

INTERPRETATION:

RESULT IN INDEX VALUE	REMARKS
< 1.30	NEGATIVE (-ve)
>=1.30	POSITIVE (+ve)

Hepatitis B Virus (HBV) is a member of the Hepadna virus family causing infection of the liver with extremely variable clinical features. Hepatitis B is transmitted primarily by body fluids especially serum and also spread effectively sexually and from mother to baby. In most individuals HBV hepatitis is self limiting, but 1-2 % normal adolescent and adults develop Chronic Hepatitis. Frequency of chronic HBV infection is 5-10% in immunocompromised patients and 80 % neonates. The initial serological marker of acute infection is HBsAg which typically appears 2-3 months after infection and disappears 12-20 weeks after onset of symtoms. Persistence of HBsAg for more than 6 months indicates carrier state or Chronic Liver disease.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mr. BIRDEVINDER SINGH**		
AGE/ GENDER	: 31 YRS/MALE	**PATIENT ID**	: 1684359
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012411270068**
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	**REGISTRATION DATE**	: 27/Nov/2024 07:05 PM
BARCODE NO.	: 01521570	**COLLECTION DATE**	: 27/Nov/2024 07:08PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 27/Nov/2024 08:38PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

VDRL

VDRL *by IMMUNOCHROMATOGRAPHY*	NON REACTIVE		NON REACTIVE

INTERPRETATION:

1.Does not become positive until 7 - 10 days after appearance ofchancre.
2.***High titer (>1:16) - active disease.***
3.***Low titer (<1:8) - biological falsepositive test in 90% cases or due to late or late latent syphillis.***
4.Treatment of primary syphillis causes progressive decline tonegative VDRL within 2 years.
5.Rising titer (4X) indicates relapse,reinfection, or treatment failure and need for retreatment.
6.May benonreactive in early primary, late latent, and late syphillis (approx. 25% ofcases).
7.Reactive and weakly reactive tests should always be confirmedwith FTA-ABS (fluorescent treponemal antibody absorptiontest).

SHORTTERM FALSE POSITIVE TEST RESULTS (<6 MONTHS DURATION) MAY OCCURIN:

1.Acute viral illnesses (e.g., hepatitis, measles, infectious mononucleosis)
2.M. pneumoniae; Chlamydia; Malaria infection.
3.Some immunizations
4.Pregnancy (rare)

LONGTERM FALSE POSITIVE TEST RESULTS (>6 MONTHS DURATION) MAY OCCUR IN:

1.Serious underlying disease e.g., collagen vascular diseases, leprosy ,malignancy.
2.Intravenous drug users.
3.Rheumatoid arthritis, thyroiditis, AIDS, Sjogren's syndrome.
4.<l0 % of patients older thanage 70 years.
5.Patients taking some anti-hypertensive drugs.

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)