

## **KOS Diagnostic Lab**

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Dog PASHA

**AGE/ GENDER** : 11 YRS/Male **PATIENT ID** : 1684801

**COLLECTED BY** REG. NO./LAB NO. :012411280049

REFERRED BY **REGISTRATION DATE** : 28/Nov/2024 02:00 PM BARCODE NO. :01521622 **COLLECTION DATE** : 28/Nov/2024 02:04PM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 28/Nov/2024 06:18PM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

## CLINICAL CHEMISTRY/BIOCHEMISTRY **AMYLASE**

550.9<sup>H</sup> IU/L 0 - 90AMYLASE - SERUM

by CNPG 3, SPECTROPHOTOMETRY

## INTERPRETATION COMMENTS

1. Amylase is produced in the Pancreas and most of the elevation in serum is due to increased rate of Amylase entry into the blood stream / decreased rate of clearance or both.

2.Serum Amylase rises within 6 to 48 hours of onset of Acute pancreatitis in 80% of patients, but is not proportional to the severity of the disease.

3.Activity usually returns to normal in 3-5 days in patients with milder edematous form of the disease.

4.Values persisting longer than this period suggest continuing necrosis of pancreas or Pseudocyst formation.

5.Approximately 20% of patients with Pancreatitis have normal or near normal activity.

6. Hyperlipemic patients with Pancreatitis also show spuriously normal Amylase levels due to suppression of Amylase activity by triglyceride. 7. Low Amylase levels are seen in Chronic Pancreatitis, Congestive Heart failure, 2nd & 3rd trimesters of pregnancy, Gastrointestinal cancer & bone fractures.

Rechecked



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





## **KOS Diagnostic Lab**

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Dog PASHA

**AGE/ GENDER** : 11 YRS/Male **PATIENT ID** : 1684801

**COLLECTED BY** REG. NO./LAB NO. :012411280049

REFERRED BY **REGISTRATION DATE** : 28/Nov/2024 02:00 PM BARCODE NO. :01521622 **COLLECTION DATE** : 28/Nov/2024 02:04PM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 28/Nov/2024 06:18PM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

LIPASE

LIPASE - SERUM  $164.38^{H}$ U/L 0 - 60

by METHYL RESORUFIN, SPECTROPHOTOMETRY

INTERPRETATION

- 1. Pancreas is the major and primary source of serum lipase though lipases are also present in liver, stomach, intestine, WBC, fat cells and milk.
  2. In acute pancreatitis, serum lipase becomes elevated at the same time as amylase and remains high for 7-10 days.

3. Increased lipase activity rarely lasts longer than 14 days

- 4. Prolonged increase suggests poor prognosis or presence of a cyst.
- The combined use of serum lipase and serum amylase is effective in ruling out acute pancreatitis.

**INCREASED LEVEL:** 

- Acute & Chronic pancreatitis
   Obstruction of pancreatic duct
   Non pancreatic conditions like renal diseases, acute cholecystitis, intestinal obstruction, duodenal ulcer, alcoholism, diabetic ketoacidosis and following endoscopic retrograde cholangiopancreatography
- 1. Elevations 2 to 50 times the upper reference have been reported. The increase in serum lipase is not necessarily proportional to the severity of the attack. Normalization is not necessarily a sign of resolution.

Concomitant testing of serum amylase and lipase is highly recommended to establish a diagnosis of pancreatic injury

End Of Report



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

