NAME : Mr. J.P ABROL

AGE/ GENDER : 80 YRS/MALE **PATIENT ID** : 1687941

COLLECTED BY : SURJESH : 012412020027 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 02/Dec/2024 10:22 AM BARCODE NO. :01521844 **COLLECTION DATE** : 02/Dec/2024 10:36AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 02/Dec/2024 11:06AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Value Unit **Biological Reference interval Test Name**

CLINICAL CHEMISTRY/BIOCHEMISTRY URIC ACID

3.60 - 7.70 URIC ACID: SERUM 3.71 mg/dL

by URICASE - OXIDASE PEROXIDASE

INTERPRETATION:-

1.GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint.

2. Uric Acid is the end product of purine metabolism. Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation.

INCREASED:-

(A).DUE TO INCREASED PRODUCTION:-

1. Idiopathic primary gout.

2. Excessive dietary purines (organ meats, legumes, anchovies, etc).

3. Cytolytic treatment of malignancies especially leukemais & lymphomas.

4. Polycythemai vera & myeloid metaplasia.

5. Psoriasis.

6.Sickle cell anaemia etc. (B).DUE TO DECREASED EXCREATION (BY KIDNEYS)

1. Alcohol ingestion.

2. Thiazide diuretics.

3.Lactic acidosis.

4. Aspirin ingestion (less than 2 grams per day).

5. Diabetic ketoacidosis or starvation.

6. Renal failure due to any cause etc.

DECREASED:-

(A).DUE TO DIETARY DEFICIENCY

- 1. Dietary deficiency of Zinc, Iron and molybdenum.
- 2.Fanconi syndrome & Wilsons disease.
- 3. Multiple sclerosis.
- 4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc.

(B). DUE TO INCREASED EXCREATION

1. Drugs:-Probenecid, sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estrogens etc.

*** End Of Report ***



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

