

**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

|                       |  |                          |                        |
|-----------------------|--|--------------------------|------------------------|
| <b>NAME</b>           | : Mr. RAM RAJ                          | <b>PATIENT ID</b>        | : 1624722              |
| <b>AGE/ GENDER</b>    | : 38 YRS/MALE                          | <b>REG. NO./LAB NO.</b>  | : 012412030052         |
| <b>COLLECTED BY</b>   | :                                      | <b>REGISTRATION DATE</b> | : 03/Dec/2024 05:30 PM |
| <b>REFERRED BY</b>    | :                                      | <b>COLLECTION DATE</b>   | : 03/Dec/2024 05:34PM  |
| <b>BARCODE NO.</b>    | : 01521923                             | <b>REPORTING DATE</b>    | : 03/Dec/2024 06:49PM  |
| <b>CLIENT CODE.</b>   | : KOS DIAGNOSTIC LAB                   |                          |                        |
| <b>CLIENT ADDRESS</b> | : 6349/1, NICHOLSON ROAD, AMBALA CANTT |                          |                        |

| Test Name | Value | Unit | Biological Reference interval |
|-----------|-------|------|-------------------------------|
|-----------|-------|------|-------------------------------|

## CLINICAL CHEMISTRY/BIOCHEMISTRY

### URIC ACID

|   |      |       |             |
|---|------|-------|-------------|
| URIC ACID: SERUM<br>by URICASE - OXIDASE PEROXIDASE | 6.07 | mg/dL | 3.60 - 7.70 |
|---|------|-------|-------------|

#### INTERPRETATION:-

1. GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint.  
 2. Uric Acid is the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation.

#### INCREASED:-

##### (A).DUE TO INCREASED PRODUCTION:-

1. Idiopathic primary gout.
2. Excessive dietary purines (organ meats, legumes, anchovies, etc).
3. Cytolytic treatment of malignancies especially leukemias & lymphomas.
4. Polycythemia vera & myeloid metaplasia.
5. Psoriasis.
6. Sick cell anaemia etc.

##### (B).DUE TO DECREASED EXCRETION (BY KIDNEYS)

1. Alcohol ingestion.
2. Thiazide diuretics.
3. Lactic acidosis.
4. Aspirin ingestion (less than 2 grams per day ).
5. Diabetic ketoacidosis or starvation.
6. Renal failure due to any cause etc.

#### DECREASED:-

##### (A).DUE TO DIETARY DEFICIENCY

1. Dietary deficiency of Zinc, Iron and molybdenum.
2. Fanconi syndrome & Wilsons disease.
3. Multiple sclerosis .
4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc.

##### (B).DUE TO INCREASED EXCRETION

1. Drugs:-Probenecid , sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosteroids and ACTH, anti-coagulants and estrogens etc.

\*\*\* End Of Report \*\*\*





**DR. VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)



**DR. YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)

