

KOS Diagnostic Lab

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. MADAN LAL

AGE/ GENDER : 82 YRS/MALE **PATIENT ID** : 1696296

COLLECTED BY REG. NO./LAB NO. :012412110003

REFERRED BY **REGISTRATION DATE** : 11/Dec/2024 08:16 AM BARCODE NO. :01522283 **COLLECTION DATE** : 11/Dec/2024 08:23AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE :11/Dec/2024 12:16PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Value Unit **Biological Reference interval Test Name**

CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F) AND POST PRANDIAL (PP)

GLUCOSE FASTING (F): PLASMA mg/dL NORMAL: < 100.0 173.65^H

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0

DIABETIC: > 0R = 126.0

NORMAL: < 140.00 GLUCOSE POST PRANDIAL (PP): PLASMA mg/dL 191.59^H

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 140.0 - 200.0

DIABETIC: > 0R = 200.0

INTERPRETATION:

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose below 100 mg/dL and post-prandial plasma glucose level below 140 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl and post-prandial plasma glucose level between 140 - 200 mg/dL is considered as glucose intolerant or pre diabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients

3. A fasting plasma glucose level of above 125 mg/dL and post-prandial plasma glucose level above 200 mg/dL is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

** End Of Report ***



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

