

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.



	Dr. Vinay Chopra MD (Pathology & Micr Chairman & Consultan	obiology)	Dr. Yugam (MD (P CEO & Consultant P	athology)
NAME	: Mr. RAHUL			
AGE/ GENDER	: 39 YRS/MALE	PA	TIENT ID	: 1618063
COLLECTED BY	:		G. NO./LAB NO.	: 012412130023
REFERRED BY	:		GISTRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO. CLIENT CODE.	: 01522398 : KOS DIAGNOSTIC LAB		LLECTION DATE PORTING DATE	: 13/Dec/2024 11:01AM : 13/Dec/2024 11:33AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMB/		I ORIEND DATE	. 13/ Dec/ 2024 11.33AM
Test Name		Value	Unit	Biological Reference interval
	SWASTI	HYA WELL	NESS PANEL: 1.0	
	СОМР	LETE BLOO	D COUNT (CBC)	
RED BLOOD CELLS	(RBCS) COUNT AND INDICES			
HAEMOGLOBIN (HB)	14.2	gm/dL	12.0 - 17.0
RED BLOOD CELL (R	BC) COUNT	4.29	Millions/cr	mm 3.50 - 5.00
PACKED CELL VOLU	ME (PCV) ITOMATED HEMATOLOGY ANALYZER	43.6	%	40.0 - 54.0
MEAN CORPUSCULA	R VOLUME (MCV) ITOMATED HEMATOLOGY ANALYZER	101.7 ^H	fL	80.0 - 100.0
	R HAEMOGLOBIN (MCH)	33.1	pg	27.0 - 34.0
MEAN CORPUSCULA	R HEMOGLOBIN CONC. (MCHC)	32.5	g/dL	32.0 - 36.0
	TION WIDTH (RDW-CV) ITOMATED HEMATOLOGY ANALYZER	15.4	%	11.00 - 16.00
	TION WIDTH (RDW-SD)	58.8 ^H	fL	35.0 - 56.0
MENTZERS INDEX		23.71	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDI by CALCULATED		36.51	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CEL				1000 11000
FOTAL LEUCOCYTE	COUNT (TLC) by sf cube & microscopy	7550	/cmm	4000 - 11000
	LOOD CELLS (nRBCS) THEMATOLOGY ANALYZER	NIL		0.00 - 20.00
	OOD CELLS (nRBCS) %	NIL	%	< 10 %



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

 KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana

 KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana

 0171-2643898, +91 99910 43898
 care@koshealthcare.com

 www.koshealthcare.com
 www.koshealthcare.com



Page 1 of 14







Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME	: Mr. RAHUL		
AGE/ GENDER	: 39 YRS/MALE	PATIENT ID	: 1618063
COLLECTED BY	:	REG. NO./LAB NO.	: 012412130023
REFERRED BY	:	REGISTRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO.	: 01522398	COLLECTION DATE	: 13/Dec/2024 11:01AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPORTING DATE	: 13/Dec/2024 11:33AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name		Value	Unit	Biological Reference interval
DIFFERENTIAL LEUCOCY	TE COUNT (DLC)			
NEUTROPHILS by FLOW CYTOMETRY BY SF C	UBE & MICROSCOPY	64	%	50 - 70
LYMPHOCYTES by FLOW CYTOMETRY BY SF C		28	%	20 - 40
EOSINOPHILS by FLOW CYTOMETRY BY SF C	UBE & MICROSCOPY	2	%	1 - 6
MONOCYTES by FLOW CYTOMETRY BY SF C	UBE & MICROSCOPY	6	%	2 - 12
BASOPHILS by FLOW CYTOMETRY BY SF C	UBE & MICROSCOPY	0	%	0 - 1
ABSOLUTE LEUKOCYTES	(WBC) COUNT			
ABSOLUTE NEUTROPHIL by FLOW CYTOMETRY BY SF C		4832	/cmm	2000 - 7500
ABSOLUTE LYMPHOCYTE by FLOW CYTOMETRY BY SF C		2114	/cmm	800 - 4900
ABSOLUTE EOSINOPHIL C by FLOW CYTOMETRY BY SF C		151	/cmm	40 - 440
ABSOLUTE MONOCYTE CC by FLOW CYTOMETRY BY SF C		453	/cmm	80 - 880
ABSOLUTE BASOPHIL COU by FLOW CYTOMETRY BY SF C		0	/cmm	0 - 110
PLATELETS AND OTHER	PLATELET PREDICTIVI	E MARKERS.		
PLATELET COUNT (PLT) by HYDRO DYNAMIC FOCUSING	G, ELECTRICAL IMPEDENCE	276000	/cmm	150000 - 450000
PLATELETCRIT (PCT) by HYDRO DYNAMIC FOCUSING	G, ELECTRICAL IMPEDENCE	0.32	%	0.10 - 0.36
MEAN PLATELET VOLUME by HYDRO DYNAMIC FOCUSING		12	fL	6.50 - 12.0
PLATELET LARGE CELL CO by HYDRO DYNAMIC FOCUSING		102000 ^H	/cmm	30000 - 90000
PLATELET LARGE CELL R. by HYDRO DYNAMIC FOCUSING		36.9	%	11.0 - 45.0
PLATELET DISTRIBUTION by HYDRO DYNAMIC FOCUSING NOTE: TEST CONDUCTED C	G, ELECTRICAL IMPEDENCE	16.4	%	15.0 - 17.0





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)







	Dr. Vinay Chopra MD (Pathology & Microl Chairman & Consultant	G/ /	(Pathology)
NAME	: Mr. RAHUL		
AGE/ GENDER	: 39 YRS/MALE	PATIENT ID	: 1618063
COLLECTED BY	:	REG. NO./LAB NO.	: 012412130023
REFERRED BY	:	REGISTRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO.	: 01522398	COLLECTION DATE	: 13/Dec/2024 11:01AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPORTING DATE	: 13/Dec/2024 11:33AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBAL	LA CANTT	
Test Name		/alue Unit	Biological Reference interval

Test Name	Value	Unit	Biological Reference interval





DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.



	MD (Pathology & Microbiology)		MD	Dr. Yugam Chopra MD (Pathology) & Consultant Pathologist	
AME	: Mr. RAHUL				
GE/ GENDER	: 39 YRS/MALE	l	PATIENT ID	: 1618063	
OLLECTED BY	:	1	REG. NO./LAB NO.	: 012412130023	
EFERRED BY	:	l	REGISTRATION DATE	: 13/Dec/2024 11:00 AM	
ARCODE NO.	: 01522398	(COLLECTION DATE	: 13/Dec/2024 11:01AM	
LIENT CODE.	: KOS DIAGNOSTIC LAB]	REPORTING DATE	: 13/Dec/2024 11:42AM	
LIENT ADDRESS	: 6349/1, NICHOLSON ROAI	D, AMBALA CANTT			
est Name		Value	Unit	Biological Reference interval	
bolycythaemia), sigr s sickle cells in sickl (OTE: . ESR and C - reactiv . Generally, ESR doe . CRP is not affected . If the ESR is elevat . Women tend to ha . Drugs such as dext	hificantly high white blood cell e cell anaemia) also lower the e protein (C-RP) are both mark is not change as rapidly as doe by as many other factors as is ed, it is typically a result of two ve a higher ESR, and menstrua	count (leucocytosis) ESR. s CRP, either at the s ESR, making it a bett o types of proteins, g tion and pregnancy c	, and some protein abno tart of inflammation or as er marker of inflammatior lobulins or fibrinogen. an cause temporary eleva	n.	





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)







		Chopra / & Microbiology) onsultant Pathologist	Dr. Yugam MD CEO & Consultant	(Pathology)
NAME	: Mr. RAHUL			
AGE/ GENDER	: 39 YRS/MALE	PATIE	NT ID	: 1618063
COLLECTED BY	:	REG. N	IO./LAB NO.	: 012412130023
REFERRED BY	:	REGIS	TRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO.	: 01522398	COLLE	CTION DATE	: 13/Dec/2024 11:01AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPO	RTING DATE	: 13/Dec/2024 01:58PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROA	D, AMBALA CANTT		
Test Name		Value	Unit	Biological Reference interval
	CLIN	ICAL CHEMISTRY/	BIOCHEMIST	'RY
	CLIN	ICAL CHEMISTRY/ GLUCOSE FAST		'nY

KOS Diagnostic Lab (A Unit of KOS Healthcare)

A fasting plasma glucose level below 100 mg/dl is considered normal.
 A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood

test (after consumption of 75 gms of glucose) is recommended for all such patients. 3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.





	Dr. Vinay Cl MD (Pathology Chairman & Co			(Pathology)
IAME	: Mr. RAHUL			
GE/ GENDER	: 39 YRS/MALE		PATIENT ID	: 1618063
OLLECTED BY	:		REG. NO./LAB NO.	: 012412130023
REFERRED BY	:		REGISTRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO.	: 01522398		COLLECTION DATE	: 13/Dec/2024 11:01AM
LIENT CODE.	: KOS DIAGNOSTIC LAB		REPORTING DATE	: 13/Dec/2024 01:25PM
LIENT ADDRESS	: 6349/1, NICHOLSON ROAD	, AMBALA CANTT		
Fest Name		Value	Unit	Biological Reference interval
		LIPID PRO	FILE : BASIC	
CHOLESTEROL TO	TAL: SERUM	245.28 ^H	mg/dL	OPTIMAL: < 200.0
by CHOLESTEROL OX	(IDASE PAP	~ 10.20	0	BORDERLINE HIGH: 200.0 -
				239.0 HIGH CHOLESTEROL: > OR =
				240.0
RIGLYCERIDES: S		234.76 ^H	mg/dL	OPTIMAL: < 150.0
by GLYCEROL PHOSE	PHATE OXIDASE (ENZYMATIC)			BORDERLINE HIGH: 150.0 - 199.0
				HIGH: 200.0 - 499.0
		177.0.1		VERY HIGH: $> OR = 500.0$
IDL CHOLESTERO by SELECTIVE INHIBIT	L (DIRECT): SERUM	47.34	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0
				60.0
			()7	HIGH HDL: $> OR = 60.0$
DL CHOLESTERO		150.99 ^H	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.
				BORDERLINE HIGH: 130.0 -
				159.0 HIGH: 160.0 - 189.0
				VERY HIGH: $> OR = 190.0$
NON HDL CHOLES		197.94 ^H	mg/dL	OPTIMAL: < 130.0
by CALCULATED, SPE	CTROPHOTOMETRY			ABOVE OPTIMAL: 130.0 - 159. BORDERLINE HIGH: 160.0 -
				189.0
				HIGH: 190.0 - 219.0
LDL CHOLESTER	OI · SERIIM	46.95 ^H	mg/dL	VERY HIGH: > OR = 220.0 0.00 - 45.00
by CALCULATED, SPE	CTROPHOTOMETRY	40.93**		
OTAL LIPIDS: SEF by calculated, spe		725.32 ^H	mg/dL	350.00 - 700.00
CHOLESTEROL/HI		5.18 ^H	RATIO	LOW RISK: 3.30 - 4.40
	CTROPHOTOMETRY			AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0
by CALCULATED, SPE				



DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.





		hopra & Microbiology) nsultant Pathologist	Dr. Yugam MD CEO & Consultant	(Pathology)
NAME	: Mr. RAHUL			
AGE/ GENDER	: 39 YRS/MALE	PATI	ENT ID	: 1618063
COLLECTED BY	:	REG.	NO./LAB NO.	: 012412130023
REFERRED BY	:	REGI	STRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO.	: 01522398	COLL	ECTION DATE	: 13/Dec/2024 11:01AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPO	DRTING DATE	: 13/Dec/2024 01:25PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD	, AMBALA CANTT		
Test Name		Value	Unit	Biological Reference interval
LDL/HDL RATIO: S by Calculated, spe		3.19 ^H	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0
TRIGLYCERIDES/H	IDL RATIO: SERUM	4.96	RATIO	3.00 - 5.00

by CALCULATED, SPECTROPHOTOMETRY

INTERPRETATION:

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL & Non HDL

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)









Dr. Vinay Chopra Dr. Yugam Chopra MD (Pathology) MD (Pathology & Microbiology) Chairman & Consultant Pathologist **CEO & Consultant Pathologist** NAME : Mr. RAHUL AGE/ GENDER : 39 YRS/MALE **PATIENT ID** :1618063 **COLLECTED BY** REG. NO./LAB NO. :012412130023 : **REFERRED BY REGISTRATION DATE** : 13/Dec/2024 11:00 AM : **BARCODE NO.** :01522398 **COLLECTION DATE** :13/Dec/2024 11:01AM CLIENT CODE. : KOS DIAGNOSTIC LAB **REPORTING DATE** :13/Dec/2024 01:25PM **CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT Test Name Value Unit LIVER FUNCTION TEST (COMPLETE) BILIBUBIN TOTAL · SERUM 0.84 mg/dI

BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY	0.84	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY	0.18	mg/dL	0.00 - 0.40
BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY	0.66	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	27.3	U/L	7.00 - 45.00
SGPT/ALT: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	33.4	U/L	0.00 - 49.00
AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	0.82	RATIO	0.00 - 46.00
ALKALINE PHOSPHATASE: SERUM by para nitrophenyl phosphatase by amino methyl propanol	115.26	U/L	40.0 - 130.0
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY	27.44	U/L	0.00 - 55.0
TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY	7.01	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL GREEN	4.65	gm/dL	3.50 - 5.50
GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.36	gm/dL	2.30 - 3.50
A : G RATIO: SERUM	1.97	RATIO	1.00 - 2.00

by CALCULATED, SPECTROPHOTOMETRY

INTERPRETATION

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range. USE: - Differential diagnosis of diseases of hepatobiliary system and pancreas.

INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com



Biological Reference interval





	Dr. Vinay Chopr MD (Pathology & Micr Chairman & Consultar	robiology) MD	n Chopra 9 (Pathology) t Pathologist
NAME	: Mr. RAHUL		
AGE/ GENDER	: 39 YRS/MALE	PATIENT ID	: 1618063
COLLECTED BY	:	REG. NO./LAB NO.	: 012412130023
REFERRED BY	:	REGISTRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO.	: 01522398	COLLECTION DATE	: 13/Dec/2024 11:01AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPORTING DATE	: 13/Dec/2024 01:25PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMB	ALA CANTT	
Tost Namo		Value Unit	Rialogical Reference interval

0	Test Name Value Unit Biological Reference
---	---

DECREASED:

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

PROGNOSTIC SIGNIFICANO	:Е:

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)







	Dr. Vinay Chop MD (Pathology & M Chairman & Consul	licrobiology)		Pathology)
NAME	: Mr. RAHUL			
AGE/ GENDER	: 39 YRS/MALE]	PATIENT ID	: 1618063
COLLECTED BY	:]	REG. NO./LAB NO.	: 012412130023
REFERRED BY	:]	REGISTRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO.	: 01522398		COLLECTION DATE	: 13/Dec/2024 11:01AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB]	REPORTING DATE	: 13/Dec/2024 02:54PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AM	MBALA CANTT		
Test Name		Value	Unit	Biological Reference interv
	KIDNE	Y FUNCTIO	N TEST (COMPLETE)	
UREA: SERUM		52.33 ^H	mg/dL	10.00 - 50.00
by UREASE - GLUTAN	NATE DEHYDROGENASE (GLDH)		Ũ	
CREATININE: SER		2.48 ^H	mg/dL	0.40 - 1.40
	ROGEN (BUN): SERUM	24.45	mg/dL	7.0 - 25.0
by CALCULATED, SPE	ECTROPHOTOMETRY			
BLOOD UREA NITH RATIO: SERUM	ROGEN (BUN)/CREATININE	9.86^L	RATIO	10.0 - 20.0
	ECTROPHOTOMETRY			
UREA/CREATININ		21.1	RATIO	
URIC ACID: SERUM	ECTROPHOTOMETRY 1	10.27 ^H	mg/dL	3.60 - 7.70
by URICASE - OXIDAS			-	
CALCIUM: SERUM by ARSENAZO III, SPE		9.95	mg/dL	8.50 - 10.60
PHOSPHOROUS: SH		3.1	mg/dL	2.30 - 4.70
	DATE, SPECTROPHOTOMETRY		0	
ELECTROLYTES				
SODIUM: SERUM by ISE (ION SELECTIV	/F FL FCTRODF)	146.3	mmol/L	135.0 - 150.0
POTASSIUM: SERU		5.07 ^H	mmol/L	3.50 - 5.00
by ISE (ION SELECTIV			1/1	00.0 110.0
CHLORIDE: SERUN by ISE (ION SELECTIV		109.73	mmol/L	90.0 - 110.0
	MERULAR FILTERATION RATE			
	ERULAR FILTERATION RATE	33		
(eGFR): SERUM by CALCULATED				
INTERPRETATION:				
To differentiate bate				

To differentiate between pre- and post renal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

2. Catabolic states with increased tissue breakdown.

3. GI haemorrhage.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

 KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana

 KOS Molecular Lab: Ilnd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana

 0171-2643898, +91 99910 43898
 care@koshealthcare.com

 www.koshealthcare.com



PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANT





		Dr. Vinay Chopra MD (Pathology & Micro Chairman & Consultant	obiology)	Dr. Yı CEO & Cons	ugam C MD (Pat sultant Pat	hology)				
NAME	: Mr. RAHUL									
AGE/ GENDER	: 39 YRS/MAI	.E	P	ATIENT ID	:	1618063				
COLLECTED BY	•			EG. NO./LAB NO.		012412130	093			
REFERRED BY	:			EGISTRATION DA		13/Dec/2024				
BARCODE NO.	:01522398			DLLECTION DATE		13/Dec/2024				
CLIENT CODE.	: KOS DIAGN	OSTIC LAB	R	EPORTING DATE		13/Dec/2024	4 02:54P	М		
CLIENT ADDRESS	: 6349/1, NIC	HOLSON ROAD, AMBA	LA CANTT							
Test Name			Value	Uni	it	Biolo	ogical R	eferenc	e interv	al
 Reduced muscle m Certain drugs (e.g. INCREASED RATIO (>2 Postrenal azotemia Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr 	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises dis superimposed 0:1) WITH DECI osis.	creatinine production) ucocorticoids) ATED CREATININE LEVE I proportionately more th on renal disease.	LS:) (e.g. obstructive	uropathy)					
 P. Certain drugs (e.g., INCREASED RATIO (>2 Postrenal azotemia Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Inherited hyperam SIADH (syndrome c Pregnancy. DECREASED RATIO (<1 Phenacimide thera Rabdomyolysis (r 	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises dis superimposed 0:1) WITH DECI osis. d starvation. creased urea sy urea rather tha monemias (ure f inappropiate 0:1) WITH INCR py (accelerates eleases muscle	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the on renal disease. REASED BUN : In creatinine diffuses of a is virtually absent in the antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine).	L S: han creatinine ut of extracell blood). due to tubular	ular fluid). secretion of urea.						
 Reduced muscle m Certain drugs (e.g. INCREASED RATIO (>2 Postrenal azotemia Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Inherited hyperam SIADH (syndrome c Pregnancy. DECREASED RATIO (<1 Phenacimide thera Rhabdomyolysis (r Muscular patients 	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises disp superimposed 0:1) WITH DECI osis. d starvation. creased urea sy urea rather tha monemias (ure f inappropiate 0:1) WITH INCR py (accelerates eleases muscle who develop re	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the on renal disease. REASED BUN : In creatinine diffuses of a is virtually absent in the antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine).	L S: han creatinine ut of extracell blood). due to tubular	ular fluid). secretion of urea.						
 B. Reduced muscle m Certain drugs (e.g., INCREASED RATIO (>2 Postrenal azotemia Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Inherited hyperam SIADH (syndrome c Pregnancy. DECREASED RATIO (<1 Phenacimide thera Rabdomyolysis (r Muscular patients INAPPROPIATE RATIO 	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises disp superimposed 0:1) WITH DECI osis. Id starvation. 2: creased urea sy urea rather tha monemias (urea f inappropiate 0:1) WITH INCR py (accelerates eleases muscle who develop re- sis (acetoaceta	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the on renal disease. REASED BUN : In creatinine diffuses of a is virtually absent in the antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine). enal failure.	L S: han creatinine ut of extracell blood). due to tubular to creatinine)	ular fluid). secretion of urea.			normal r	atio whe	n dehyd	ratic
 Reduced muscle m Certain drugs (e.g. NCREASED RATIO (>2 Postrenal azotemia Prerenal azotemia Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Inherited hyperam SIADH (syndrome c Pregnancy. PCREASED RATIO (<1 Phenacimide thera Rhabdomyolysis (r Muscular patients NAPPROPIATE RATIO Diabetic ketoacido should produce an in Cephalosporin ther 	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises dis superimposed 0:1) WITH DECI osis. d starvation. creased urea sy urea rather tha monemias (urea f inappropiate 0:1) WITH INCR py (accelerates eleases muscle who develop re- sis (acetoaceta creased BUN/cl apy (interferes	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the on renal disease. REASED BUN : The sis. In creatinine diffuses of a is virtually absent in the antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine). enal failure. the causes false increase reatinine ratio). with creatinine measur	LS: han creatinine ut of extracell blood). due to tubular to creatinine; e in creatinine	ular fluid). secretion of urea.			normal r	atio whe	n dehyd	ratio
B. Reduced muscle m Certain drugs (e.g. NCREASED RATIO (>2 Postrenal azotemia Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Inherited hyperam SIADH (syndrome c Pregnancy. DECREASED RATIO (<1 Phenacimide thera Rhabdomyolysis (r Muscular patients NAPPROPIATE RATIO Diabetic ketoacido hould produce an in Cephalosporin ther STIMATED GLOMERL	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises dis superimposed 0:1) WITH DECI osis. d starvation. creased urea sy urea rather tha monemias (urea f inappropiate 0:1) WITH INCR py (accelerates eleases muscle who develop re- sis (acetoaceta creased BUN/cl apy (interferes	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the on renal disease. REASED BUN : The sis. In creatinine diffuses of a is virtually absent in the antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine). Inal failure. the causes false increase reatinine ratio). with creatinine measur DN RATE:	LS: nan creatinine ut of extracell blood). due to tubular to creatinine) e in creatinine rement).	ular fluid). secretion of urea. with certain meth	nodologies	,resulting in r		atio whe	n dehyd	ratic
B. Reduced muscle m Certain drugs (e.g. NCREASED RATIO (>2 Postrenal azotemia Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Inherited hyperam SIADH (syndrome c Pregnancy. DECREASED RATIO (<1 Phenacimide thera Rhabdomyolysis (r Muscular patients NAPPROPIATE RATIO Diabetic ketoacido hould produce an in Cephalosporin ther	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises dis superimposed 0:1) WITH DECI osis. d starvation. 2: creased urea sy urea rather tha monemias (urea f inappropiate 0:1) WITH INCR py (accelerates eleases muscle who develop re- sis (acetoaceta creased BUN/cl apy (interferes UAR FILTERATIC	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the on renal disease. REASED BUN : The sis. In creatinine diffuses of a is virtually absent in the antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine). enal failure. te causes false increase reatinine ratio). with creatinine measur	LS: nan creatinine ut of extracell blood). due to tubular to creatinine) e in creatinine rement).	ular fluid). secretion of urea.	nodologies	,resulting in r		atio whe	n dehyd	ratic
B. Reduced muscle m Certain drugs (e.g. NCREASED RATIO (>2 Postrenal azotemia Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Diherited hyperam SIADH (syndrome of Pregnancy. DECREASED RATIO (<1 Phenacimide thera Rhabdomyolysis (r Nuscular patients NAPPROPIATE RATIO Diabetic ketoacido hould produce an in Cephalosporin ther STADE	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises dis superimposed 0:1) WITH DECI osis. Id starvation. 2: creased urea sy urea rather tha monemias (urea f inappropiate 0:1) WITH INCR py (accelerates eleases muscle who develop re- sis (acetoaceta creased BUN/cl apy (interferes LAR FILTERATIO	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the proportionately more the proportionately more the proportionately more the proportionately more the proportionately more the proportionately more the antidiures. EASED BUN : Conversion diffuses of antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine). enal failure. the causes false increase reatinine ratio). with creatinine measure on RATE: DESCRIPTION rmal kidney function idney damage with	LS: nan creatinine ut of extracell blood). due to tubular to creatinine) e in creatinine rement).	ular fluid). secretion of urea. with certain meth min/1.73m2)	nodologies ASSOC	,resulting in r ATED FINDIN proteinuria nce of Protein	GS	atio whe	n dehyd	ratic
A. Reduced muscle m Certain drugs (e.g. NCREASED RATIO (>2 Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Inherited hyperam SIADH (syndrome c Pregnancy. DECREASED RATIO (<1 Nhenacimide thera Rhabdomyolysis (r Muscular patients NAPPROPIATE RATIO Diabetic ketoacido should produce an in Cephalosporin ther ESTIMATED GLOMERL CKD STAGE G1 G2	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises dis superimposed 0:1) WITH DECI osis. Id starvation. 2: creased urea sy urea rather tha monemias (urea f inappropiate 0:1) WITH INCR py (accelerates eleases muscle who develop re- sis (acetoaceta creased BUN/cl apy (interferes LAR FILTERATION No	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the on renal disease. REASED BUN : The sis. In creatinine diffuses of a is virtually absent in the antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine). enal failure. the causes false increase reatinine ratio). with creatinine measure on RATE: DESCRIPTION rmal kidney function idney damage with formal or high GFR	LS: han creatinine ut of extracell blood). due to tubular to creatinine; e in creatinine rement). GFR (mL/	ular fluid). secretion of urea. with certain meth <u>min/1.73m2)</u> >90 >90	nodologies ASSOC	,resulting in r ATED FINDING proteinuria	GS	atio whe	n dehyd	ratio
A. Reduced muscle m Certain drugs (e.g. NCREASED RATIO (>2 Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Inherited hyperam SIADH (syndrome of Pregnancy. DECREASED RATIO (<1 Phenacimide thera Rhabdomyolysis (r Muscular patients NAPPROPIATE RATIO Diabetic ketoacido should produce an in Cephalosporin ther ESTIMATED GLOMERL CKD STAGE G1 G2 G3a	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises disp superimposed 0:1) WITH DECI osis. Id starvation. 2. creased urea sy urea rather tha monemias (urea f inappropiate 0:1) WITH INCR py (accelerates eleases muscle who develop re- sis (acetoaceta creased BUN/ca apy (interferes LAR FILTERATION NO NO NO NO NO NO NO NO NO NO NO NO N	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the on renal disease. REASED BUN : The creatinine diffuses of a is virtually absent in the antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine). anal failure. the causes false increase reatinine ratio). with creatinine measure N RATE: DESCRIPTION rmal kidney function idney damage with iormal or high GFR ild decrease in GFR	LS: han creatinine ut of extracell blood). due to tubular to creatinine; e in creatinine ement). GFR (mL/	ular fluid). secretion of urea. with certain meth <u>min/1.73m2)</u> >90 >90 0 -89	nodologies ASSOC	,resulting in r ATED FINDIN proteinuria nce of Protein	GS	atio whe	n dehyd	ratic
B. Reduced muscle m Certain drugs (e.g. INCREASED RATIO (>2 Prerenal azotemia DECREASED RATIO (<1 Acute tubular necr Low protein diet ar Severe liver disease Other causes of de Repeated dialysis (Inherited hyperam SIADH (syndrome c Pregnancy. DECREASED RATIO (<1 Nhenacimide thera Rhabdomyolysis (r Muscular patients INAPPROPIATE RATIO Diabetic ketoacido should produce an in Cephalosporin ther ESTIMATED GLOMERL CKD STAGE G1 G2	ass (subnormal tetracycline, gl 0:1) WITH ELEV (BUN rises disp superimposed 0:1) WITH DECI osis. Id starvation. 2: creased urea sy urea rather tha monemias (urea f inappropiate 0:1) WITH INCR py (accelerates eleases muscle who develop re- sis (acetoaceta creased BUN/cr apy (interferes LAR FILTERATION NO NO NO NO NO	creatinine production) ucocorticoids) ATED CREATININE LEVEL proportionately more the on renal disease. REASED BUN : The sis. In creatinine diffuses of a is virtually absent in the antidiuretic harmone) of EASED CREATININE: conversion of creatine creatinine). enal failure. the causes false increase reatinine ratio). with creatinine measure on RATE: DESCRIPTION rmal kidney function idney damage with formal or high GFR	LS: han creatinine ut of extracell blood). due to tubular to creatinine; e in creatinine ement). GFR (mL/	ular fluid). secretion of urea. with certain meth <u>min/1.73m2)</u> >90 >90	nodologies ASSOC	,resulting in r ATED FINDIN proteinuria nce of Protein	GS	atio whe	n dehyd	ratic





DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)









	Dr. Vinay Chopra MD (Pathology & Microbic Chairman & Consultant Pa		(Pathology)
NAME	: Mr. RAHUL		
AGE/ GENDER	: 39 YRS/MALE	PATIENT ID	: 1618063
COLLECTED BY	:	REG. NO./LAB NO.	: 012412130023
REFERRED BY	:	REGISTRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO.	: 01522398	COLLECTION DATE	: 13/Dec/2024 11:01AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPORTING DATE	: 13/Dec/2024 02:54PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA	CANTT	
Test Name	Va	lue Unit	Biological Reference interval

COMMENTS:

Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.
 eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012
 In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure of CFD with the commended to measure

3. In patients, with eGFR cleaning between 45-59 minimit 1.73 m2 (G3) and without any marker of Kidney damage, it is recommended to measure eGFR with Cystatin C for confirmation of CKD
4. eGFR category G1 OR G2 does not fulfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

ADVICE:

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

MBBS, MD (PATHOLOGY)







	Dr. Vinay Ch MD (Pathology & Chairman & Cons		Dr. Yugan MD CEO & Consultant	(Pathology)
NAME	: Mr. RAHUL			
AGE/ GENDER	: 39 YRS/MALE	PAT	FIENT ID	: 1618063
COLLECTED BY	:	REG	G. NO./LAB NO.	: 012412130023
REFERRED BY	:	REG	GISTRATION DATE	: 13/Dec/2024 11:00 AM
BARCODE NO.	: 01522398	COL	LLECTION DATE	: 13/Dec/2024 11:01AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REI	PORTING DATE	: 13/Dec/2024 12:44PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, A	AMBALA CANTT		
Test Name		Value	Unit	Biological Reference interval
		CLINICAL PA	THOLOGY	
	URINE RO	UTINE & MICRO	SCOPIC EXAMIN	ATION
PHYSICAL EXAMI				
QUANTITY RECIEV		10	ml	
COLOUR		PALE YELLO	N	PALE YELLOW
TRANSPARANCY	TANCE SPECTROPHOTOMETRY	HAZY		CLEAR
SPECIFIC GRAVITY	TANCE SPECTROPHOTOMETRY	>=1.030		1.002 - 1.030
CHEMICAL EXAMI				
REACTION	TANCE SPECTROPHOTOMETRY	ACIDIC		
PROTEIN		3+		NEGATIVE (-ve)
by DIP STICK/REFLEG	CTANCE SPECTROPHOTOMETRY	Negative		NEGATIVE (-ve)
	TANCE SPECTROPHOTOMETRY	Negative		
pH	TANCE SPECTROPHOTOMETRY	<=5.0		5.0 - 7.5
BILIRUBIN		Negative		NEGATIVE (-ve)
by DIP STICK/REFLEC	TANCE SPECTROPHOTOMETRY			
	TANCE SPECTROPHOTOMETRY.	Negative		NEGATIVE (-ve)
UROBILINOGEN by DIP STICK/REFLEC	TANCE SPECTROPHOTOMETRY	Normal	EU/dL	0.2 - 1.0
KETONE BODIES		Negative		NEGATIVE (-ve)
BLOOD	TANCE SPECTROPHOTOMETRY	2+		NEGATIVE (-ve)
by DIP STICK/REFLEC	CTANCE SPECTROPHOTOMETRY	NEGATIVE (-v		NEGATIVE (-ve)
	TANCE SPECTROPHOTOMETRY	MEGATIVE (-)		NEGATIVE (-VC)
MICROSCOPIC EX				
RED BLOOD CELLS	(RBCs) CENTRIFUGED URINARY SEDIMENT	15-18	/HPF	0 - 3





DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





NANCE





Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME	: Mr. RAHUL					
AGE/ GENDER	: 39 YRS/MALE	PATIENT 1	ID	: 1618063		
COLLECTED BY	:	REG. NO./LAB NO.		: 012412130023		
REFERRED BY:BARCODE NO.: 01522398CLIENT CODE.: KOS DIAGNOSTIC LAB		REGISTRATION DATE COLLECTION DATE REPORTING DATE		: 13/Dec/2024 11:00 AM : 13/Dec/2024 11:01AM		
				CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AM	MBALA CANTT
Test Name		Value	Unit	Biological Reference interval		
PUS CELLS	CENTRIFUGED URINARY SEDIMENT	2-3	/HPF	0 - 5		
EPITHELIAL CELL		1-2	/HPF	ABSENT		
-						

by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT NEGATIVE (-ve) NEGATIVE (-ve) CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT BACTERIA NEGATIVE (-ve) NEGATIVE (-ve) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT OTHERS NEGATIVE (-ve) NEGATIVE (-ve) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT TRICHOMONAS VAGINALIS (PROTOZOA) ABSENT ABSENT

by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT

DATIT

** End Of Report ***



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) Thopsa Hopsa

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

 KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana

 KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana

 0171-2643898, +91 99910 43898
 care@koshealthcare.com
 www.koshealthcare.com



Page 14 of 14