

TEST PERFORMED AT KOS DIAGNOSTIC LAB. AMBALA CANTT



		& Microbiology)	Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist	
NAME	: Mr. NARESH SHARMA			
AGE/ GENDER	: 66 YRS/MALE	PATIEN	T ID	: 1702226
COLLECTED BY	:	REG. NO)./LAB NO.	: 012412180027
REFERRED BY	:	REGIST	RATION DATE	: 18/Dec/2024 11:19 AM
BARCODE NO.	:01522626	COLLEC	TION DATE	: 18/Dec/2024 11:21AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPOR	FING DATE	:18/Dec/2024 12:34PM
Test Name		Value	Unit	Biological Reference interv
	MICROALBU	CLINICAL PATH MIN/CREATININE R		MURINE
MICROALBUMIN: RANDOM URINE by SPECTROPHOTOMETRY			17	0.05
		93.9 ^H	mg/L	0 - 25
	IETRY DOM URINE	93.9ⁿ 51.4	mg/L mg/dL	0 - 25 20 - 320
by SPECTROPHOTOM CREATININE: RAN by SPECTROPHOTOM MICROALBUMIN/(RANDOM URINE by SPECTROPHOTOM	IETRY DOM URINE IETRY CREATININE RATIO -		C	
by SPECTROPHOTOM CREATININE: RAN by SPECTROPHOTOM MICROALBUMIN/(RANDOM URINE by SPECTROPHOTOM	IETRY DOM URINE IETRY CREATININE RATIO -	51.4	mg/dL	20 - 320
by SPECTROPHOTOM CREATININE: RAN by SPECTROPHOTOM MICROALBUMIN/(RANDOM URINE by SPECTROPHOTOM INTERPRETATION:-	IETRY DOM URINE IETRY CREATININE RATIO - IETRY NORMAL: mg/L	51.4	mg/dL mg/g	20 - 320

Long standing un-treated Diabetes and Hypertension can lead to renal dysfunction. 2. Diabetic nephropathy or kidney disease is the most common cause of end stage renal disease(ERSD) or kidney failure. 3. Presence of Microalbuminuria is an early indicator of onset of compromised renal function in these patients. 4.Microalbuminuria is the condition when urinary albumin excre tion is between 30-300 mg & above this it is called as macroalbuminuria, the presence of which indicates serious kidney disease. 5.Microalbuminuria is not only associated with kidney disease but of cardiovascular disease in patients with dibetes & hypertension. 6. Microalbuminuria of only associated with kidney disease but of cardiovascular disease in patients with dibetes & hypertension.

6.Microalbuminuria reflects vascular damage & appear to be a marker of of early arterial disease & endothelial dysfunction. **NOTE:-** IF A PATIENT HAS = 1+ PROTEINURIA (30 mg/dl OR 300 mg/L) BY URINE DIPSTICK (URINEANALYSIS), OVERT PROTEINURIA IS PRESENT AND TESTING FOR MICROALBUMIN IS INAPPROPIATE. IN SUCH A CASE, URINE PROTEIN:CREATININE RATIO OR 24 HOURS TOTAL URINE MICROPROTEIN IS APPROPIATE.

*** End Of Report ***





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 care@koshealthcare.com www.koshealthcare.com

