

NAME	: Mr. VIRENDER NATH KUMARIA	PATIENT ID	: 1710579
AGE/ GENDER	: 82 YRS/MALE	REG. NO./LAB NO.	: 012412280011
COLLECTED BY	: SURJESH	REGISTRATION DATE	: 28/Dec/2024 09:06 AM
REFERRED BY	:	COLLECTION DATE	: 28/Dec/2024 09:35AM
BARCODE NO.	: 01523109	REPORTING DATE	: 30/Dec/2024 02:53PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## MICROBIOLOGY

### CULTURE AEROBIC BACTERIA AND ANTIBIOTIC SENSITIVITY: URINE

#### CULTURE AND SUSCEPTIBILITY: URINE

DATE OF SAMPLE 28-12-2024

SPECIMEN SOURCE URINE

INCUBATION PERIOD 48 HOURS

by AUTOMATED BROTH CULTURE

CULTURE STERILE

by AUTOMATED BROTH CULTURE

ORGANISM NO AEROBIC PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF  
INCUBATION AT 37°C

by AUTOMATED BROTH CULTURE

#### AEROBIC SUSCEPTIBILITY: URINE

##### INTERPRETATION:

1. In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection.  
2. Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out" catheterization or from patients with indwelling catheters.

##### SUSCEPTIBILITY:

1. A test interpreted as **SENSITIVE** implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent recommended for that type of infection and infecting species, unless otherwise indicated..  
2. A test interpreted as **INTERMEDIATE** implies that the "infection due to the isolate may be appropriately treated in body sites where the drugs are physiologically concentrated or when a high dosage of drug can be used".  
3. A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.


##### CAUTION:


Conditions which can cause a false Negative culture:

1. Patient is on antibiotics. Please repeat culture post therapy.
2. Anaerobic bacterial infection.
3. Fastidious aerobic bacteria which are not able to grow on routine culture media.
4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.
5. Renal tuberculosis to be confirmed by AFB studies.

\*\*\* End Of Report \*\*\*



  
DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)

