

Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. RISHAVJOT KAUR	PATIENT ID	: 1711877
AGE/ GENDER	: 29 YRS/FEMALE	REG. NO./LAB NO.	: 012412300034
COLLECTED BY	:	REGISTRATION DATE	: 30/Dec/2024 03:20 PM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	COLLECTION DATE	: 30/Dec/2024 03:20PM
BARCODE NO.	: 01523222	REPORTING DATE	: 30/Dec/2024 03:37PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

HAEMATOLOGY

HAEMOGLOBIN (HB)

HAEMOGLOBIN (HB) by CALORIMETRIC	12.1	gm/dL	12.0 - 16.0
-------------------------------------	------	-------	-------------

INTERPRETATION:-

Hemoglobin is the protein molecule in red blood cells that carries oxygen from the lungs to the bodys tissues and returns carbon dioxide from the tissues back to the lungs.

A low hemoglobin level is referred to as ANEMIA or low red blood count.

ANEMIA (DECREASED HAEMOGLOBIN):

- 1) Loss of blood (traumatic injury, surgery, bleeding, colon cancer or stomach ulcer)
- 2) Nutritional deficiency (iron, vitamin B12, folate)
- 3) Bone marrow problems (replacement of bone marrow by cancer)
- 4) Suppression by red blood cell synthesis by chemotherapy drugs
- 5) Kidney failure
- 6) Abnormal hemoglobin structure (sickle cell anemia or thalassemia).


POLYCYTHEMIA (INCREASED HAEMOGLOBIN):

- 1) People in higher altitudes (Physiological)
- 2) Smoking (Secondary Polycythemia)
- 3) Dehydration produces a falsely rise in hemoglobin due to increased haemoconcentration
- 4) Advanced lung disease (for example, emphysema)
- 5) Certain tumors
- 6) A disorder of the bone marrow known as polycythemia rubra vera,
- 7) Abuse of the drug erythropoetin (Epogen) by athletes for blood doping purposes (increasing the amount of oxygen available to the body by chemically raising the production of red blood cells).

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD




DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS , MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. RISHAVJOT KAUR	PATIENT ID	: 1711877
AGE/ GENDER	: 29 YRS/FEMALE	REG. NO./LAB NO.	: 012412300034
COLLECTED BY	:	REGISTRATION DATE	: 30/Dec/2024 03:20 PM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	COLLECTION DATE	: 30/Dec/2024 03:20PM
BARCODE NO.	: 01523222	REPORTING DATE	: 30/Dec/2024 03:38PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		


Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

BLOOD GROUP (ABO) AND RH FACTOR TYPING

ABO GROUP
 by SLIDE AGGLUTINATION
 RH FACTOR TYPE
 by SLIDE AGGLUTINATION

O
 POSITIVE




 DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. RISHAVJOT KAUR	PATIENT ID	: 1711877
AGE/ GENDER	: 29 YRS/FEMALE	REG. NO./LAB NO.	: 012412300034
COLLECTED BY	:	REGISTRATION DATE	: 30/Dec/2024 03:20 PM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	COLLECTION DATE	: 30/Dec/2024 03:20PM
BARCODE NO.	: 01523222	REPORTING DATE	: 30/Dec/2024 05:08PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

BLEEDING TIME (BT)

BLEEDING TIME (BT) by DUKE METHOD	1 MIN 45 SEC	MINS	1 - 5
--------------------------------------	--------------	------	-------




DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist


NAME	: Mrs. RISHAVJOT KAUR	PATIENT ID	: 1711877
AGE/ GENDER	: 29 YRS/FEMALE	REG. NO./LAB NO.	: 012412300034
COLLECTED BY	:	REGISTRATION DATE	: 30/Dec/2024 03:20 PM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	COLLECTION DATE	: 30/Dec/2024 03:20PM
BARCODE NO.	: 01523222	REPORTING DATE	: 30/Dec/2024 05:08PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

CLOTTING TIME (CT)

CLOTTING TIME (CT)	5 MIN 15 SEC	MINS	4 - 9
by CAPILLARY TUBE METHOD			




 DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. RISHAVJOT KAUR	PATIENT ID	: 1711877
AGE/ GENDER	: 29 YRS/FEMALE	REG. NO./LAB NO.	: 012412300034
COLLECTED BY	:	REGISTRATION DATE	: 30/Dec/2024 03:20 PM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	COLLECTION DATE	: 30/Dec/2024 03:20PM
BARCODE NO.	: 01523222	REPORTING DATE	: 30/Dec/2024 04:11PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

CLINICAL CHEMISTRY/BIOCHEMISTRY

GLUCOSE RANDOM (R)


GLUCOSE RANDOM (R): PLASMA by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)	90.94	mg/dL	NORMAL: < 140.00 PREDIABETIC: 140.0 - 200.0 DIABETIC: > OR = 200.0
---	-------	-------	--


INTERPRETATION

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A random plasma glucose level below 140 mg/dl is considered normal.
2. A random glucose level between 140 - 200 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
3. A random glucose level of above 200 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.




DR. VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR. YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. RISHAVJOT KAUR	PATIENT ID	: 1711877
AGE/ GENDER	: 29 YRS/FEMALE	REG. NO./LAB NO.	: 012412300034
COLLECTED BY	:	REGISTRATION DATE	: 30/Dec/2024 03:20 PM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	COLLECTION DATE	: 30/Dec/2024 03:20PM
BARCODE NO.	: 01523222	REPORTING DATE	: 30/Dec/2024 04:20PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

ENDOCRINOLOGY

BETA HCG - TOTAL (QUANTITATIVE): MATERNAL

BETA HCG TOTAL, PREGNANCY MATERNAL: **952.87^H** mIU/mL < 5.0
 SERUM
 by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

INTERPRETATION:

MEN:	mIU/ml	< 2.0
NON PREGNANT PRE-MENOPAUSAL WOMEN:	mIU/ml	< 5.0
MENOPAUSAL WOMEN:	mIU/ml	< 7.0
BETA HCG EXPECTED VALUES IN ACCORDANCE TO WEEKS OF GESTATIONAL AGE		
WEEKS OF GESTATION	Unit	Value
4-5	mIU/ml	1500 -23000
5-6	mIU/ml	3400 - 135300
6-7	mIU/ml	10500 - 161000
7-8	mIU/ml	18000 - 209000
8-9	mIU/ml	37500 - 219000
9-10	mIU/ml	42800 - 218000
10-11	mIU/ml	33700 - 218700
11-12	mIU/ml	21800 - 193200
12-13	mIU/ml	20300 - 166100
13-14	mIU/ml	15400 - 190000
2rd TRIMESTER	mIU/ml	2800 - 176100
3rd TRIMESTER	mIU/ml	2800 - 144400




DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. RISHAVJOT KAUR	PATIENT ID	: 1711877
AGE/ GENDER	: 29 YRS/FEMALE	REG. NO./LAB NO.	: 012412300034
COLLECTED BY	:	REGISTRATION DATE	: 30/Dec/2024 03:20 PM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	COLLECTION DATE	: 30/Dec/2024 03:20PM
BARCODE NO.	: 01523222	REPORTING DATE	: 30/Dec/2024 04:20PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

1.hCG is a Glycoprotein with alpha and beta chains. Beta subunit is specific to hCG.

2.It is largely secreted by trophoblastic tissue. Small amounts may be secreted by fetal tissues and by the adult ant pituitary.

INCREASED :

1.Pregnancy

2.Gestationalsite & Non gestational trophoblastic neoplasia.

3.In mixed germ cell tumors.

SIGNIFICANTLY HIGHER THAN EXPECTED LEVEL:

1.Multiple pregnancies & High risk molar pregnancies are usually associated with levels in excess of one lac mIU/ml.

2.Erythroblastosis fetalis & Down's syndrome.

DECREASED:

1.Ectopic pregnancy.

2.Intra-uterine fetal death.

NOTE:

1.The test becomes positive 7-9 days after the midcycle surge that precedes ovulation (time of blastocyst implantation). Blood levels rise rapidly after this and double every 1.4 - 2 days.

2.Peak values are usually seen at 60-80 days of LMP. The levels then begin to taper and ebb out around the 20th week. These low levels are then maintained throughout pregnancy.

3.Doubling time: In intra-uterine pregnancy, serum hCG levels increase by approximately 66% every 48 hrs.Inappropriately rising serum hCG levels are suggestive of dying or ectopic pregnancy.

CAUTION:

Spuriously high levels (Phantom hCG) may be seen in presence of heterophilic antibodies (found in some normal people). If persistently raised levels are seen in a non-pregnant patient with no evidence of other obvious causes for such an increase a urine hCG assay may help confirm presence of the heterophile antibodies.




 DR.VINAY CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mrs. RISHAVJOT KAUR
AGE/ GENDER : 29 YRS/FEMALE
COLLECTED BY :
REFERRED BY : LOOMBA HOSPITAL (AMBALA CANTT)
BARCODE NO. : 01523222
CLIENT CODE. : KOS DIAGNOSTIC LAB
CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

PATIENT ID : 1711877
REG. NO./LAB NO. : 012412300034
REGISTRATION DATE : 30/Dec/2024 03:20 PM
COLLECTION DATE : 30/Dec/2024 03:20PM
REPORTING DATE : 30/Dec/2024 04:20PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

IMMUNOPATHOLOGY/SEROLOGY

HEPATITIS C VIRUS (HCV) ANTIBODY: TOTAL

HEPATITIS C ANTIBODY (HCV) TOTAL: SERUM 0.06 S/CO
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)
NEGATIVE: < 1.00
POSITIVE: > 1.00

HEPATITIS C ANTIBODY (HCV) TOTAL NON - REACTIVE
RESULT
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

INTERPRETATION:-

RESULT (INDEX)	REMARKS
< 1.00	NON - REACTIVE/NOT - DETECTED
> =1.00	REACTIVE/ASYMPTOMATIC/INFECTIVE STATE/CARRIER STATE.

Hepatitis C (HCV) is an RNA virus of Favivirus group transmitted via blood transfusions, transplantation, injection drug abusers, accidental needle punctures in healthcare workers, dialysis patients and rarely from mother to infant. 10 % of new cases show sexual transmission. As compared to HAV & HBV , chronic infection with HCV occurs in 85 % of infected individuals. In high risk population, the predictive value of Anti HCV for HCV infection is > 99% whereas in low risk populations it is only 25 %.

USES:

- Indicator of past or present infection, but does not differentiate between Acute/ Chronic/Resolved Infection.
- Routine screening of low and high prevalence population including blood donors.

NOTE:

- False positive results are seen in Auto-immune disease, Rheumatoid Factor, HYpergammaglobulinemia, Paraproteinemia, Passive antibody transfer, Anti-idiotypes and Anti-superoxide dismutase.
- False negative results are seen in early Acute infection, Immunosuppression and Immuno— incompetence.
- HCV-RNA PCR recommended in all reactive results to differentiate between past and present infection.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. RISHAVJOT KAUR	PATIENT ID	: 1711877
AGE/ GENDER	: 29 YRS/FEMALE	REG. NO./LAB NO.	: 012412300034
COLLECTED BY	:	REGISTRATION DATE	: 30/Dec/2024 03:20 PM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	COLLECTION DATE	: 30/Dec/2024 03:20PM
BARCODE NO.	: 01523222	REPORTING DATE	: 30/Dec/2024 04:20PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

ANTI HUMAN IMMUNODEFICIENCY VIRUS (HIV) DUO ULTRA WITH (P-24 ANTIGEN DETECTION)

HIV 1/2 AND P24 ANTIGEN: SERUM	0.05	S/CO	NEGATIVE: < 1.00 POSITIVE: > 1.00
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			
HIV 1/2 AND P24 ANTIGEN RESULT	NON - REACTIVE		
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			

INTERPRETATION:-

RESULT (INDEX)	REMARKS
< 1.00	NON - REACTIVE
> = 1.00	PROVISIONALLY REACTIVE

Non-Reactive result implies that antibodies to HIV 1/ 2 have not been detected in the sample . This means that patient has either not been exposed to HIV 1/ 2 infection or the sample has been tested during the "window phase" i.e. before the development of detectable levels of antibodies. Hence a Non Reactive result does not exclude the possibility of exposure or infection with HIV 1/ 2.

RECOMMENDATIONS:

1. Results to be clinically correlated
2. Rarely falsenegativity/positivity may occur.




 DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mrs. RISHAVJOT KAUR
AGE/ GENDER : 29 YRS/FEMALE
COLLECTED BY :
REFERRED BY : LOOMBA HOSPITAL (AMBALA CANTT)
BARCODE NO. : 01523222
CLIENT CODE. : KOS DIAGNOSTIC LAB
CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

PATIENT ID : 1711877
REG. NO./LAB NO. : 012412300034
REGISTRATION DATE : 30/Dec/2024 03:20 PM
COLLECTION DATE : 30/Dec/2024 03:20PM
REPORTING DATE : 30/Dec/2024 04:20PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

HEPATITIS B SURFACE ANTIGEN (HBsAg) ULTRA

HEPATITIS B SURFACE ANTIGEN (HBsAg): 0.25 S/CO
SERUM
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)
NEGATIVE: < 1.0
POSITIVE: > 1.0

HEPATITIS B SURFACE ANTIGEN (HBsAg) NON REACTIVE
RESULT
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

INTERPRETATION:

RESULT IN INDEX VALUE	REMARKS
< 1.30	NEGATIVE (-ve)
>=1.30	POSITIVE (+ve)

Hepatitis B Virus (HBV) is a member of the Hepadna virus family causing infection of the liver with extremely variable clinical features. Hepatitis B is transmitted primarily by body fluids especially serum and also spread effectively sexually and from mother to baby. In most individuals HBV hepatitis is self limiting, but 1-2 % normal adolescent and adults develop Chronic Hepatitis. Frequency of chronic HBV infection is 5-10% in immunocompromised patients and 80 % neonates. The initial serological marker of acute infection is HBsAg which typically appears 2-3 months after infection and disappears 12-20 weeks after onset of symptoms. Persistence of HBsAg for more than 6 months indicates carrier state or Chronic Liver disease.




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)



Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. RISHAVJOT KAUR	PATIENT ID	: 1711877
AGE/ GENDER	: 29 YRS/FEMALE	REG. NO./LAB NO.	: 012412300034
COLLECTED BY	:	REGISTRATION DATE	: 30/Dec/2024 03:20 PM
REFERRED BY	: LOOMBA HOSPITAL (AMBALA CANTT)	COLLECTION DATE	: 30/Dec/2024 03:20PM
BARCODE NO.	: 01523222	REPORTING DATE	: 30/Dec/2024 03:44PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

VDRL

VDRL by IMMUNOCHROMATOGRAPHY	NON REACTIVE	NON REACTIVE
---------------------------------	--------------	--------------

INTERPRETATION:

- Does not become positive until 7 - 10 days after appearance of chancre.
- High titer (>1:16) - active disease.**
- Low titer (<1:8) - biological falsepositive test in 90% cases or due to late or late latent syphilis.**
- Treatment of primary syphilis causes progressive decline tonegative VDRL within 2 years.
- Rising titer (4X) indicates relapse, reinfection, or treatment failure and need for retreatment.
- May benonreactive in early primary, late latent, and late syphilis (approx. 25% ofcases).
- Reactive and weakly reactive tests should always be confirmedwith FTA-ABS (fluorescent treponemal antibody absorptiontest).**

SHORTTERM FALSE POSITIVE TEST RESULTS (<6 MONTHS DURATION) MAY OCCURIN:


- Acute viral illnesses (e.g., hepatitis, measles, infectious mononucleosis)
- M. pneumoniae; Chlamydia; Malaria infection.
- Some immunizations
- Pregnancy (rare)

LONGTERM FALSE POSITIVE TEST RESULTS (>6 MONTHS DURATION) MAY OCCUR IN:

- Serious underlying disease e.g., collagen vascular diseases, leprosy ,malignancy.
- Intravenous drug users.
- Rheumatoid arthritis, thyroiditis, AIDS, Sjogren's syndrome.
- <10 % of patients older thanage 70 years.
- Patients taking some anti-hypertensive drugs.

*** End Of Report ***




 DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS , MD (PATHOLOGY)

