

Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. SUSHMA	PATIENT ID	: 1713526
AGE/ GENDER	: 24 YRS/FEMALE	REG. NO./LAB NO.	: 012501010021
COLLECTED BY	:	REGISTRATION DATE	: 01/Jan/2025 12:34 PM
REFERRED BY	:	COLLECTION DATE	: 01/Jan/2025 12:36PM
BARCODE NO.	: 01523295	REPORTING DATE	: 01/Jan/2025 01:29PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

IMMUNOPATHOLOGY/SEROLOGY
WIDAL SLIDE AGGLUTINATION TEST

SALMONELLA TYPHI O <i>by SLIDE AGGLUTINATION</i>	1 : 160	TITRE	1 : 80
SALMONELLA TYPHI H <i>by SLIDE AGGLUTINATION</i>	1 : 40	TITRE	1 : 160
SALMONELLA PARATYPHI AH <i>by SLIDE AGGLUTINATION</i>	1 : 20	TITRE	1 : 160
SALMONELLA PARATYPHI BH <i>by SLIDE AGGLUTINATION</i>	NIL	TITRE	1 : 160

INTERPRETATION:

1. Titres of 1:80 or more for "O" agglutinin is considered significant.
2. Titres of 1:160 or more for "H" agglutinin is considered significant.

LIMITATIONS:


1. Agglutinins usually appear by 5th to 6th day of illness of enteric fever, hence a negative result in early stage is inconclusive. The titre then rises till 3rd or 4th week, after which it declines gradually.
2. Lower titres may be found in normal individuals.
3. A single positive result has less significance than the rising agglutination titre, since demonstration of rising titre four or more in 1st and 3rd week is considered as a definite evidence of infection.
4. A simultaneous rise in H agglutinins is suggestive of paratyphoid infection.


NOTE:

1. Individuals with prior infection or immunization with TAB vaccine may develop an ANAMNESTIC RESPONSE (False-Positive) during an unrelated fever i.e High titres of antibodies to various antigens. This may be differentiated by repetition of the test after a week.
2. The anamnestic response shows only a transient rise, while in enteric fever rise is sustained.
3. H agglutinins tend to persist for many months after vaccination but O agglutinins tend to disappear sooner i.e within 6 months. Therefore rise in O agglutinins indicate recent infection.

*** End Of Report ***




DR. VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR. YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.