NAME : Mr. AVTAR SINGH

AGE/ GENDER : 52 YRS/MALE **PATIENT ID** :1729974

: 012501210034 **COLLECTED BY** REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 21/Jan/2025 01:15 PM BARCODE NO. :01524194 **COLLECTION DATE** : 21/Jan/2025 01:17PM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 21/Jan/2025 05:44PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Value Unit **Biological Reference interval** Test Name

CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE POST PRANDIAL (PP)

GLUCOSE POST PRANDIAL (PP): PLASMA by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) 435.5H

mg/dL

NORMAL: < 140.00

PREDIABETIC: 140.0 - 200.0 DIABETIC: > 0R = 200.0

INTERPRETATION
IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A post-prandial plasma glucose level below 140 mg/dl is considered normal.

2. A post-prandial glucose level between 140 - 200 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood

test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A post-prandial plasma glucose level of above 200 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

*** End Of Report ***



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

